



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.


We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





UNIVERSITY

STANFORD



LIBRARIES

UNIVERSITY



STANFORD

STANFORD



LIBRARIES

LIBRARIES









II.

a

THE
OLUTION OF MAN:

POPULAR EXPOSITION OF THE PRINCIPALS
S OF HUMAN ONTOGENY AND PHYLOGENY.

FROM THE GERMAN OF
ERNST HAECKEL,

PROFESSOR IN THE UNIVERSITY OF JENA,
AUTHOR OF "THE HISTORY OF CREATION," ETC.

IN TWO VOLUMES.
VOL. II.



LONDON:
GAN PAUL, TRENCH & CO., 1, PATERNOSTER SQUARE.
1883.



By the same Author.

THE HISTORY OF CREATION. Translation revised by Professor E. RAY LANKESTER, M.A., F.R.S. With Coloured Plates and Genealogical Trees of the various groups of both Plants and Animals. 2 vols. Second Edition. Post 8vo., cloth, 82s.

A VISIT TO CEYLON. Translated by CLARA BELL. Post 8vo., cloth, 7s. 6d.

FREEDOM IN SCIENCE AND TEACHING. A Reply to Professor VIRCHOW on the Freedom of Science in the Modern State. Authorised Translation, with Introduction by Professor HUXLEY. Crown 8vo., cloth, 5s.

LONDON : KEGAN PAUL, TRENCH & Co.

THE
VOLUTION OF MAN:

*POPULAR EXPOSITION OF THE PRINCIPAL
NTS OF HUMAN ONTOGENY AND PHYLOGENY.*

FROM THE GERMAN OF
ERNST HAECKEL,

PROFESSOR IN THE UNIVERSITY OF JENA,
AUTHOR OF "THE HISTORY OF CREATION," ETC.

IN TWO VOLUMES.
VOL. II.



LONDON:
EGAN PAUL, TRENCH & CO., 1, PATERNOSTER SQUARE.
1883.

575

H133e

v. 2



A 26830
FEB 24 1899

(The rights of translation and of reproduction are reserved.)

CONTENTS OF VOL. II.

	PAGE
List of Plates 	xiii
List of Woodcuts 	xiv
List of Genetic Tables 	xvii

CHAPTER XV.

THE DURATION OF HUMAN TRIBAL HISTORY.

Comparison of Ontogenetic and Phylogenetic Periods of Time.—Duration of Germ-history in Man and in Different Animals.—Extreme Brevity of the Latter in Comparison with the Immeasurable Long Periods of Tribal History.—Relation of this Rapid Ontogenetic Modification to the Slow Phylogenetic Metamorphosis.—Estimate of the Past Duration of the Organic World, founded on the Relative Thickness of Sedimentary Rock-strata, or Neptunian Formations.—The Five Main Divisions in the Latter: I. Primordial, or Archilithic Epoch. II. Primary, or Palæolithic Epoch. III. Secondary, or Mesolithic Epoch. IV. Tertiary, or Cæolithic Epoch. V. Quaternary, or Anthropolithic Epoch.—The Relative Duration of the Five Epochs.—The Results of Comparative Philology as Explaining the Phylogeny of Species.—The Inter-relations of the Main Stems and Branches of the Indo-Germanic Languages are Analogous to the Inter-relations of the Main Stems and Branches of the Vertebrate Tribe.—The Parent Forms in both Cases are Extinct.—The Most Important Stages among the Human Ancestral Forms.—Monera originated by Spontaneous Generation.—Necessity of Spontaneous Generation 1

CHAPTER XVI.

THE ANCESTRY OF MAN.

I. FROM THE MONERA TO THE GASTRÆA.

Relation of the General Inductive Law of the Theory of Descent to the Special Deductive Laws of the Hypotheses of Descent.—Incompleteness of the Three Great Records of Creation: Palæontology, Ontogeny, and Comparative Anatomy.—Unequal Certainty of the Various Special Hypotheses of Descent.—The Ancestral Line of Men in Twenty-two Stages: Eight Invertebrate and Fourteen Vertebrate Ancestors.—Distribution of these Twenty-two Parent-forms in the Five Main Divisions of the Organic History of the Earth.—First Ancestral Stage: Monera.—The Structureless and Homogeneous Plasson of the Monera.—Differentiation of the Plasson into Nucleus, and the Protoplasm of the Cells.—Cytods and Cells as Two Different Plastid-forms.—Vital Phenomena of Monera.—Organisms without Organs.—Second Ancestral Stage: Amœbæ.—One-celled Primitive Animals of the Simplest and most Undifferentiated Nature.—The Amœboid Egg-cells.—The Egg is Older than the Hen.—Third Ancestral Stage: Syn-Amœba, Ontogenetically reproduced in the Morula.—A Community of Homogeneous Amœboid Cells.—Fourth Ancestral Stage: Planœa, Ontogenetically reproduced in the Blastula or Planula.—Fifth Ancestral Stage: Gastrœa, Ontogenetically reproduced in the Gastrula and the Two-layered Germ-disc.—Origin of the Gastrœa by Inversion (*invagination*) of the Planœa.—Haliphysema and Gastrophysema.—Extant Gastrœads

CHAPTER XVII.

THE ANCESTRAL SERIES OF MAN.

II. FROM THE PRIMITIVE WORM TO THE SKULLED ANIMAL.

The Four Higher Animal Tribes are descended from the Worm Tribe.—The Descendants of the Gastrœa; in one direction the Parent Form of Plant-Animals (Sponges and Sea-Nettles), in the other the Parent Form of Worms.—Radiate form of the former, Bilateral form of the latter.—The Two Main Divisions of the Worms, Acoelomi and Coelomati: the former without, the latter with, a Body Cavity and Blood-vessel System.—Sixth Ancestral Stage: Archelminthes, most nearly allied to Turbellaria.—Descent of the

CONTENTS.

vii

	PAGE
Coelomati from the Acoelomi.—Mantled Animals (<i>Tunicata</i>) and Chorda-Animals (<i>Chordonia</i>).—Seventh Stage: Soft-Worms (<i>Scolecida</i>).—A Side Branch of the latter: the Acorn-Worm (<i>Balanoglossus</i>).—Differentiation of the Intestinal Tube into Gill-intestine and Stomach-intestine.—Eighth Stage: Chorda-Animals (<i>Chordonia</i>).—Ascidian Larva exhibits the Outline of a Chorda-Animal.—Construction of the Notochord.—Mantled Animals and Vertebrates as Diverging Branches of Chorda-Animals.—Separation of Vertebrates from the other Higher Animal Tribes (Articulated Animals, Star-Animals, Soft-bodied Animals).—Significance of the Metamerie Formation.—Skull-less Animals (<i>Acrania</i>) and Skulled Animals (<i>Craniota</i>).—Ninth Ancestral Stage: Skull-less Animals.— <i>Amphioxus</i> and Primitive Vertebrate.—Development of Skulled Animals (Construction of the Head, Skull, and Brain).—Tenth Ancestral Stage: Skulled Animals, allied to the Cyclostomi (<i>Myxinoidea</i> and <i>Petromyzonidae</i>) 71	

CHAPTER XVIII.

THE PEDIGREE OF MAN.

III. FROM THE PRIMITIVE FISH TO THE AMNIOTIC ANIMAL.

Comparative Anatomy of the Vertebrates.—The Characteristic Qualities of the Double-nostrilled and Jaw-mouthed: the Double-Nostrils, the Gill-arch Apparatus, with the Jaw-arches, the Swimming-bladder, the Two Pairs of Limbs.—Relationship of the Three Groups of Fishes: the Primitive Fishes (*Selachii*), the Ganoids (*Ganoides*), the Osseous Fishes (*Teleostei*).—Dawn of Terrestrial Life on the Earth.—Modification of the Swimming-bladder into the Lungs.—Intermediate Position of the Dipneusta between the Primitive Fishes and Amphibia.—The Three Extant Dipneusta (*Protopterus*, *Lepidosiren*, *Ceratodus*).—Modification of the Many-toed Fin of the Fish into the Five-toed Foot.—Causes and Effects of the latter.—Descent of all Higher Vertebrates from a Five-toed Amphibian.—Intermediate Position of the Amphibians between the Lower and Higher Vertebrates.—Modification or Metamorphosis of Frogs.—Different Stages in Amphibian Metamorphosis.—The Gilled Batrachians (*Proteus* and *Axolotl*).—The Tailed Batrachians (*Salamanders* and *Mud-fish*).—Frog Batrachians (*Frogs* and *Toads*).—Chief Group of the Amnion Animals, or *Amniota* (*Reptiles*, *Birds*, and *Mammals*).—Descent of all the *Amniota* from a Common

Lizard-like Parent-form (*Protamnion*).—First Formation of Allantois and of the Amnion.—Branching of the Amnion Anin in Two Lines: on the one side, Reptiles (and Birds), on the other side, Mammals

CHAPTER XIX.

THE PEDIGREE OF MAN.

IV. FROM THE PRIMITIVE MAMMAL TO THE APE.

The Mammalian Character of Man.—Common Descent of all Mammals from a Single Parent-form (*Promammalian*).—Bifurcation of Amnion Animals into Two Main Lines: on the one side, Reptiles and Birds, on the other, Mammals.—Date of the Origin of Mammals: the Trias Period.—The Three Main Groups or Classes of Mammals: their Genealogical Relations.—Sixteenth Ancestral Stage: Cloacal Animals (*Monotremata*, or *Ornithodelphia*).—The Extinct Primitive Mammals (*Promammalia*) and the Extinct Beaked Animals (*Ornithostoma*).—Seventeenth Ancestral Stage: Pouched Animals (*Marsupialia*, or *Didelphia*).—Extinct and Extant Pouched Animals.—Their Intermediate Position between Monotremes and Placental Animals.—Origin and Structure of Placental Animals (*Placentalia*, or *Monodelphia*).—Formation of the Placenta.—The Deciduous Embryonic Membrane (*Decidua*).—Grading of the *Indecidua* and of the *Deciduata*.—The Formation of the *Decidua (vera, serotina, reflexa)* in Man and in Apes.—Eighteenth Stage: Semi-apes (*Prosimia*).—Nineteenth Stage: Tailed Ape (*Monocerca*).—Twentieth Stage: Man-like Apes (*Anthropoides*).—Speechless and Speaking Men (*Mali. Homines*) ...

CHAPTER XX.

THE HISTORY OF THE EVOLUTION OF THE EPIDERMIS AND OF THE NERVOUS SYSTEM.

Animal and Vegetative Organ-systems.—Original Relations of these to the Two Primary Germ-layers.—Sensory Apparatus.—Constitution of Sensory Apparatus: originally only the Exoderm, or Skin-layer; afterwards, the Skin-covering specialized from the Nerve-system.—Double Function of the Skin (as a Covering and as Organ).

CONTENTS.

ix

PAGE

Touch).—Outer Skin (<i>Epidermis</i>) and Leather-skin (<i>Corium</i>).— Appendages of the Epidermis: Skin-glands (Sweat-glands, Tear- glands, Sebaceous Glands, Milk-glands); Nails and Hair.—The Embryonic Wool-covering.—Hair of the Head and of the Beard.— Influence of Sexual Selection.—Arrangement of the Nerve-system. —Motor and Sensory Nerves.—Central Marrow: Brain and Dorsal Marrow.—Constitution of the Human Brain: Large Brain (<i>Cere- brum</i>) and Small Brain (<i>Cerebellum</i>).—Comparative Anatomy of the Central Marrow.—Germ-history of the Medullary-tube.—Sepa- ration of the Medullary-tube into Brain and Dorsal Marrow.— Modification of the Simple Brain-bladder into Five Consecutive Brain-bladders: Fore-brain (Large Brain, or <i>Cerebrum</i>), Twixt- brain ("Centre of Sight"), Mid-brain ("Four Bulbs"), Hind-brain (Small Brain, or <i>Cerebellum</i>), After-brain (Neck Medulla).—Various Formation of the Five Brain-bladders in the various Vertebrate Classes.—Development of the Conductive Marrow, or "Peripheric Nervous System" 190
--

CHAPTER XXI.

DEVELOPMENT OF THE SENSE-ORGANS.

Origin of the most highly Purposive Sense-organs by no Preconceived Purpose, but simply by Natural Selection.—The Six Sense-organs and the Seven Sense-functions.—All the Sense-organs originally Developed from the Outer Skin-covering (from the Skin-sensory Layer).—Organs of the Pressure Sense, the Heat Sense, the Sexual Sense, and the Taste Sense.—Structure of the Organ of Scent.—The Blind Nose-pits of Fishes.—The Nasal Furrows change into Nasal Canals.—Separation of the Cavities of the Nose and Mouth by the Palate Roof.—Structure of the Eye.—The Primary Eye Vesicles (Stalked Protuberances from the Twixt-brain).— Inversion of this Eye Vesicle by the Crystalline Lens, separated from the Horn-plate.—Inversion of the Vitreous Body.—The Vas- cular Capsule and the Fibrous Capsule of the Eyeball.—Eyelids. —Structure of the Ear.—The Apparatus for Perception of Sound: Labyrinth and Auditory Nerve.—Origin of the Labyrinth from the Primitive Ear Vesicles (by Separation from the Horn-plate).— Conducting Apparatus of Sound: Drum Cavity, Ear Bonelets, and Drum Membrane.—Origin of these from the First Gill-opening and the Parts immediately round it (the First and Second Gill- arch).—Rudimentary Outer Ear.—Rudimentary Muscles of the Ear-shell 233

CHAPTER XXII.

DEVELOPMENT OF THE ORGANS OF MOTION.

The Motive Apparatus of Vertebrates.—These are constituted by the Passive and Active Organs of Motion (Skeleton and Muscles).—The Significance of the Internal Skeleton of Vertebrates.—Structure of the Vertebral Column.—Formation and Number of the Vertebrae.—The Ribs and Breast-bone.—Germ-history of the Vertebral Column.—The Notochord.—The Primitive Vertebral Plates.—The Formation of the Metamers.—Cartilaginous and Bony Vertebrae.—Intervertebral Disca.—Head-skeleton (Skull and Gill-arches).—Vertebral Theory of the Skull (Goethe and Oken, Huxley and Gegenbaur).—Primitive Skull, or Primordial Cranium.—Its Formation from Nine or Ten Coalescent Metamers.—The Gill-arches (Ribs of the Head).—Bones of the Two Pairs of Limbs.—Development of the Five-toed Foot, adapted for Walking, from the Many-toed Fin of the Fish.—The Primitive Fin of the Selachian (*Archipterygium* of Gegenbaur).—Transition of the Pinnate into the Semi-pinnate Fin.—Atrophy of the Rays or Toes of the Fins.—Many-fingered and Five-fingered Vertebrates.—Comparison of the Anterior Limbs (Pectoral Fins) and the Posterior Limbs (Ventral Fins).—Shoulder Girdle and Pelvis Girdle.—Germ-history of the Limbs.—Development of the Muscles

CHAPTER XXIII.

DEVELOPMENT OF THE INTESTINAL SYSTEM.

The Primitive Intestine of the Gastrula.—Its Homology, or Morphological Identity in all Animals (excepting the Protozoa).—Survey of the Structure of the Developed Intestinal Canal in Man.—The Mouth-cavity.—The Throat (*pharynx*).—The Gullet (*oesophagus*).—The Wind-pipe (*trachea*) and Lungs.—The Larynx.—The Stomach.—The Small Intestine.—The Liver and Gall-bladder.—The Ventral Salivary Gland (*pancreas*).—The Large Intestine.—The Rectum.—The First Rudiment of the Simple Intestinal Tube.—The Gastrula of the Amphioxus and of Mammals.—Separation of the Germ from the Intestinal Germ Vesicle (*Gastrocystis*).—The Primitive Intestine (Protagaster) and the After Intestine (Metagaster).—Secondary Formation of the Mouth and Anus from the Outer Skin.—Development of the Intestinal Epithelium from the Intestinal-glandular Layer, and of all other parts of the Intestine from the Intestinal fibrous Layer.—Simple Intestinal Pouch of the Lower Worms.—

CONTENTS.

xi

	PAGE
Differentiation of the Primitive Intestinal Tube into a Respiratory and a Digestive Intestine.—Gill-intestine and Stomach-intestine of the Amphioxus and Ascidian.—Origin and Significance of the Gill-openings.—Their Disappearance.—The Gill-arches and the Jaw-Skeleton.—Formation of the Teeth.—Development of the Lungs from the Swim-bladder of Fish.—Differentiation of the Stomach.—Development of the Liver and Pancreas.—Differentiation of the Small and Large Intestines.—Formation of the Cloaca ...	... 311

CHAPTER XXIV.

DEVELOPMENT OF THE VASCULAR SYSTEM.

Application of the Fundamental Law of Biogeny.—The Two Sides.—Hereditv of Conservative Organs.—Adaptation of Progressive Organs.—Ontogeny and Comparative Anatomy complementary of each other.—New "Theories of Evolution" of His.—The "Envelope Theory" and the "Waste-rag Theory."—Main Germ and Supplementary Germ.—Formative Yolk and Nutritive Yolk.—Phylogenetic Origin of the latter from the Primitive Intestine.—Origin of the Vascular System from the Vascular Layer, or Intestinal-fibrous Layer.—Phylogenetic Significance of the Ontogenetic Succession of the Organ-systems and Tissues.—Deviation from the Original Sequence; Ontogenetic Heterochronism.—Covering Tissue.—Connective Tissue.—Nerve-muscle Tissue.—Vascular Tissue.—Relative Age of the Vascular System.—First Commencement of the latter; Coeloma.—Dorsal Vessel and Ventral Vessel of Worms.—Simple Heart of Ascidia.—Atrophy of the Heart in the Amphioxus.—Two-chambered Heart of the Cyclostoma.—Arterial Arches of the Selachii.—Double Auricle in Dipneusta and Amphibia.—Double Ventricle in Birds and Mammals.—Arterial Arches in Birds and Mammals.—Germ-history (Ontogeny) of the Human Heart.—Parallelism of the Tribal-history (Phylogeny) ...	... 348
---	---------

CHAPTER XXV.

DEVELOPMENT OF THE URINARY AND SEXUAL ORGANS.

Importance of Reproduction.—Growth.—Simplest Forms of Asexual Reproduction: Division and the Formation of Buds (Gemmation).—Simplest Forms of Sexual Reproduction: Amalgamation of Two Differentiated Cells; the Male Sperm-cell and the Female Egg-cell.—Fertilization.—Source of Love.—Original Hermaphroditism;
--

Later Separation of the Sexes (Gonochorism).—Original Development of the Two Kinds of Sexual Cells from the Two Primary Germ-layers.—The Male Exoderm and Female Entoderm.—Development of the Testes and Ovaries.—Passage of the Sexual Cells into the Coelom.—Hermaphrodite Rudiment of the Embryonic Epithelium, or Sexual Plate.—Channels of Exit, or Sexual Ducts.—Egg-duct and Seed-duct.—Development of these from the Primitive Kidney Ducts.—Excretory Organs of worms.—“Coiled Canals” of Ringed Worms (*Annelida*).—Side Canals of the *Amphioxus*.—Primitive Kidneys of the *Myxinoidea*.—Primitive Kidneys of Skulled Animals (*Craniota*).—Development of the Permanent Secondary Kidneys in *Amniota*.—Development of the Urinary Bladder from the Allantois.—Differentiation of the Primary and Secondary Primitive Kidney Ducts.—The Müllerian Duct (Egg-duct) and the Wolffian Duct (Seed-duct).—Change of Position of the Germ-glands in Mammals.—Formation of the Egg in Mammals (Graafian Follicle).—Origin of the External Sexual Organs.—Formation of the Cloaca.—Hermaphroditism in Man 38

CHAPTER XXVI.

RESULTS OF ANTHROPOGENY.

Review of the Germ-history as given.—Its Explanation by the Fundamental Law of Biogeny.—Its Causal Relation to the History of the Tribe.—Rudimentary Organs of Man.—Dysteleology, or the Doctrine of Purposelessness.—Inheritances from Apes.—Man's place in the Natural System of the Animal Kingdom.—Man as a Vertebrate and a Mammal.—Special Tribal Relation of Men and Apes.—Evidences regarding the Ape Question.—The <i>Catarrhina</i> and the <i>Platyrrhina</i> .—The Divine Origin of Man.—Adam and Eve.—History of the Evolution of the Mind.—Important Mental Differences within a Single Class of Animals.—The Mammalian Mind and the Insect Mind.—Mind in the Ant and in the Scale-louse (<i>Coccus</i>).—Mind in Man and in Ape.—The Organ of Mental Activity: the Central Nervous System.—The Ontogeny and Phylogeny of the Mind.—The Monistic and Dualistic Theories of the Mind.—Heredity of the Mind.—Bearing of the Fundamental Law of Biogeny on Psychology.—Influence of Anthropogeny on the Victory of the Monistic Philosophy and the Defeat of the Dualistic.—Nature and Spirit.—Natural Science and Spiritual Science.—Conception of the World reformed by Anthropogeny 43
NOTES. Remarks and References to Literature 45
INDEX 49

LIST OF PLATES.

		PAGE
Plate XII. (between p. 130 and p. 131). The Australian Mud- fish (<i>Ceratodus Fosteri</i>)	Explanation	118
Plate XIII. (between p. 130 and p. 131). The Mexican Axolotl (<i>Siredon pisciformis</i>) and the European Land-salamander (<i>Salamandra maculata</i>)	Explanation	129
Plate XIV. (between p. 180 and p. 181). Four Catarhines (Chimpanzee, Gorilla, Orang, Negro) ...	Explanation	181
Plate XV. (between p. 188 and p. 189). Pedigree of Man	Explanation	184

LIST OF WOODCUTS.

FIGURE	PAGE	FIGURE	PAGE
163. Moneron (<i>Protamoeba</i>) . . .	46	195, 196. Beaked Animal (<i>Ornithorhynchus</i>) and its skeleton . . .	1
164. Bathybius, primitive slime . . .	49	197. Pouched Animal (Marsupial) with young . . .	1
165. Monerula of Mammal . . .	51	198. Human egg-membranes . . .	1
166. Cytula of Mammal . . .	51	199. Semi-ape (Lori) . . .	1
167. Amœba	53	200. Human germ with its membranes	1
168. Amœboid egg-cell . . .	53	201. Human uterus, navel-cord, and embryo	1
169. Original egg-cleavage . . .	55	202. Head of Nose-ape . . .	1
170. Mulberry-germ (<i>Morula</i>) . . .	55	203. Tailed Ape (Sea-cat) . . .	1
171. Germination of Monoxenia . . .	57	204. Skeleton of Gibbon . . .	1
172, 173. Magosphœra . . .	60	205. Skeleton of Orang-outang . . .	1
174-179. Gastrula of various animals	65	206. Skeleton of Chimpanzee . . .	1
180, 181. Haliphysema . . .	67	207. Skeleton of Gorilla . . .	1
182, 183. Ascula of a Sponge . . .	68	208. Skeleton of Man . . .	1
184, 185. A Gliding-worm (<i>Rhabdocœlum</i>) . . .	80	209. Gastrula of Gastrophysa	1
186. Acorn-worm (<i>Balanoglossus</i>)	86	210. Germ-layers of Earth-worms	1
187. Appendicularia . . .	90	211. Nerve-system of Gliding-worm	1
188. Ascidia	90	212. Human skin-covering . . .	1
189. Amphioxus	91		
190. Lamprey (<i>Petromyzon</i>) . . .	103		
191, 192. Shark (<i>Selachii</i>) . . .	113		
193. Larval Salamander . . .	127		
194. Larval Frog (Tadpole) . . .	127		

LIST OF WOODCUTS.

XV

FIGURE	PAGE	FIGURE	PAGE
3. Epidermis cells . . .	201	257. Portion of notochord . . .	286
4. Tear-glands . . .	202	258-260. Growth of the primi-	
5, 216. Milk-glands . . .	203	tive vertebral series in	
7, 218. Central marrow of		embryo Chick . . .	288
human embryo . . .	210	261. Longitudinal section of	
9. Human brain . . .	212	breast-vertebra . . .	290
10. " " . . .	213	262. Transverse section of same	291
11-223. Lyre-shaped embryo		263. Intervertebral disc . . .	291
Chick	218	264. Human skull . . .	292
24-226. The five brain-blad-		265. Head skeleton of Primi-	
ders of the human germ	220	tive Fish	296
27. The five brain-bladders of		266. Primitive skull of Man . . .	297
<i>Craniota</i>	222	267. Skeleton of fin of <i>Ceratodus</i>	302
28. Brain of Shark . . .	222	268. Skeleton of fin of <i>Acon-</i>	
29. Brain of Frog . . .	222	<i>thias</i>	302
30. Brain of Rabbit . . .	224	269. Skeleton of fin of Primi-	
31. Nose of Shark . . .	241	tive Fish	302
32-236. Development of the		270. Skeleton of hand of Frog	302
face in embryo Chick . . .	243	271. Skeleton of hand of Gorilla	302
37. Nose and mouth cavities . . .	246	272. Skeleton of human hand . . .	302
38-240. The face in the		273. Skeleton of hand of Mam-	
human embryo	247	mal	306
41. Human eye	250	274. Gastrula of <i>Olynthus</i> . . .	313
42. Development of the eyes	253	275. Human stomach . . .	317
43. " "	256	276. Gastrula of <i>Amphioxus</i> . . .	321
44. Human auditory passage	260	277. Gastrula of Mammal . . .	321
45. Human auditory labyrinth	263	278, 279. Human germ with	
46-248. Development of the		yolk-sac and allantois . . .	324
ear	264	280. Intestine of <i>Turbellaria</i> . . .	327
49. Primitive skull with ear-		281. Intestine of <i>Ascidia</i> . . .	327
vesicles	264	282. Intestine of <i>Amphioxus</i> . . .	328
50. Rudimentary ear-muscles	270	283. Scales of Shark . . .	332
51, 252. Human skeleton . . .	279	284, 285. Intestine of embryo	
53. Human vertebral column	280	Dog with the intestinal	
54. Neck-vertebra	281	glands	334
55. Breast-vertebra	281	286. Intestine with allantois . . .	338
56. Lumbar-vertebra	281	287. Intestine of human germ	339

FIGURE	PAGE	FIGURE	PAGE
288. Liver of human germ	342	316. Rudiments of <i>Urogenitalia</i>	400
289. Nail-tissue	362	317. Primitive kidney of <i>Bdello-</i>	
290. Intestinal epithelium	362	<i>stoma</i>	406
291. Jelly-like tissue	363	318. Earliest primitive kidney	
292. Cartilaginous tissue	363	rudiments	406
293. Neuro-muscular cells	364	319, 320. Primitive kidneys of	
294. Nerve-tissue	364	Mammals	406
295. Muscle-tissue	364	321. Development of urogeni-	
296. Vascular tissue	365	tal system	414
297. Blood-cells	365	322, 323. " "	415
298. Blood-vessels of a Worm .	371	324-326. " "	416
299. Head with blood-vessels		327. Female sexual organs of	
of Fish	375	Beaked Animal (<i>Orni-</i>	
300-302. Arterial arches	377	<i>thorhynchus</i>)	418
303-306. " "	378	328. Change of position of both	
307-310. Development of the		kinds of sexual glands	
heart	380	in human beings	420
311-314. Development of the		329. Development of the human	
heart	382	external sexual organs	423
315. Transverse section		330. Human egg-follicles	426
through Haliphysema	393		

LIST OF GENETIC TABLES.



TABLE	PAGE
XII. Systematic Survey of palæontological periods ...	11
XIII. Systematic Survey of palæontological formations ...	12
XIV. Systematic Survey of the thickness of the formations	19
XV. Pedigree of Indo-Germanic languages	23
XVI. Systematic Survey of the most important stages in the animal ancestral line of Man	44
XVII. Systematic Survey of the five first stages in the evolution of Man (phylogenetic, ontogenetic, systematic)	70
XVIII. Systematic Survey of the phylogenetic system of the animal kingdom	92
XIX. Monophyletic pedigree of the animal kingdom ...	93
XX. Systematic Survey of the phylogenetic system of Vertebrates	120
XXI. Monophyletic pedigree of Vertebrates	121
XXII. Systematic Survey of the periods of human tribal history	184
XXIII. Systematic Survey of the phylogenetic system of Mammals, founded on the Gastræa Theory ...	187
XXIV. Monophyletic pedigree of Mammals	188
XXV. Pedigree of Apes	189
XXVI. Systematic Survey of the organ-systems of the human body	194
XXVII. Systematic Survey of the phylogenetic history of the human skin-covering	229
XXVIII. Systematic Survey of the phylogenetic history of the human nerve-system	230

TABLE	PAGE
XXIX. Systematic Survey of the ontogeny of the skin and nerve systems	233
XXX. Systematic Survey of the phylogeny of the human nose	249
XXXI. Systematic Survey of the ontogeny of the human eye	258
XXXII. Systematic Survey of the phylogeny of the human ear	267
XXXIII. Systematic Survey of the ontogeny of the human ear	268
XXXIV. Systematic Survey of the constitution of the human skeleton	278
XXXV. Systematic Survey of the phylogeny of the human skeleton	300
XXXVI. Systematic Survey of the constitution of the human intestinal system	330
XXXVII. Systematic Survey of the phylogeny of the human intestinal system	340
XXXVIII. Systematic Survey of the sequence, according to age, of the human tissue-groups (phylogenetic sequence of the tissues)	366
XXXIX. Systematic Survey of the sequence, according to age, of the human organ-systems (phylogenetic sequence of the organs)	367
XL. Systematic Survey of the phylogeny of the human vascular system	394
XLI. Systematic Survey of the phylogeny of the human heart	396
XLII. Systematic Survey of the homologies of Worms, Articulated Animals (<i>Arthropoda</i>), Soft-bodied Animals (<i>Mollusca</i>), and Vertebrates	397
XLIII. Systematic Survey of the phylogeny of the human urinary and sexual organs	428
XLIV. Systematic Survey of the homologies of the sexual organs in the two sexes of Mammals	431

THE EVOLUTION OF MAN.

CHAPTER XV.

THE DURATION OF HUMAN TRIBAL HISTORY.

Comparison of Ontogenetic and Phylogenetic Periods of Time.—Duration of Germ-history in Man and in Different Animals.—Extreme Brevity of the Latter in Comparison with the Immeasurable Long Periods of Tribal History.—Relation of this Rapid Ontogenetic Modification to the Slow Phylogenetic Metamorphosis.—Estimate of the Past Duration of the Organic World, founded on the Relative Thickness of Sedimentary Rock-strata, or Neptunian Formations.—The Five Main Divisions in the Latter: I. Primordial, or Archilithic Epoch. II. Primary, or Palæolithic Epoch. III. Secondary, or Mesolithic Epoch. IV. Tertiary, or Cænelithic Epoch. V. Quaternary, or Anthropolithic Epoch.—The Relative Duration of the Five Epochs.—The Results of Comparative Philology as Explaining the Phylogeny of Species.—The Inter-relations of the Main Stems and Branches of the Indo-Germanic Languages are Analogous to the Inter-relations of the Main Stems and Branches of the Vertebrate Tribe.—The Parent Forms in both Cases are Extinct.—The Most Important Stages among the Human Ancestral Forms.—Monera originated by Spontaneous Generation.—Necessity of Spontaneous Generation.

"In vain as yet has it been attempted to draw an exact line of demarcation between historic and prehistoric times; the origin of man and the period of his first appearance pass back into indefinable time; the so-called archaic age cannot be sharply distinguished from the present age. This is the fate of all geological, as of all historical periods. The periods which we distinguish are, therefore, more or less arbitrarily defined, and, like the divisions

in systematic natural history, can only serve to bring the subject of our study better before us and to render it more manageable; but not to make real distinctions between different things."—BERNHARD COTTA (1866).

OUR comparative study of the Anatomy and Ontogeny of the Amphioxus and the Ascidian has afforded us aid, the value of which can hardly be over-estimated, toward acquiring a knowledge of human Ontogeny. For in the first place we have in this way filled up, as regards Anatomy, the wide chasm which in all previous systems of the animal kingdom existed between Vertebrates and Invertebrates; and in the second place, in the germ-history of the Amphioxus we have recognized primordial phases of development, which have long disappeared from the Ontogeny of Man, and which have been lost in accordance with the law of abridged heredity. Of special importance among these phases of development is the Archigastrula, the original, genuine Gastrula-form which the Amphioxus has retained up to the present time, and which re-appears in the same form in low invertebrate animals of the most diverse classes.

The germ-history of the Amphioxus and the Ascidian has, therefore, so far perfected our direct knowledge of human genealogy, that, notwithstanding the incompleteness of our empiric knowledge, there is no essential gap of any great moment in the pedigree. We may, therefore, at once proceed to our task, and, aided by the ontogenetic and comparative-anatomical materials at our command, may reconstruct the main outlines of human Phylogeny. The immense importance of the direct application of the fundamental biogenetic law of the causal connection between Ontogeny and Phylogeny now becomes evident. But, before

beginning this task, it will be well to note a few other general facts which may enable us better to understand the phenomena we are about to study.

Firstly, it may not be out of place to insert a few remarks as to the duration of time during which Man was developing from the animal kingdom. The first thought that occurs to the mind when we consider the facts in question, is of the immense difference between the duration of the germ-history of Man on the one hand, and of his tribal history on the other. The brief period in which the Ontogeny of the human individual takes place, bears no proportion to the infinitely long period required for the Phylogeny of the human tribe. The human individual requires nine months for its perfect development from the fertilized egg-cell to the moment at which it is born and quits the mother's body. The human embryo, therefore, passes through the whole course of its development in the brief space of 40 weeks (usually in exactly 280 days). Each man is really older by this period than is usually assumed. When, for example, a child is said to be $9\frac{1}{4}$ years old, he is in reality 10 years old. For individual existence does not begin at the moment of birth, but at the moment of fertilization. In many other Mammals the duration of the embryonic development is the same as in Man, *e.g.*, the Ox. In the Horse and the Ass it is somewhat longer, viz., from 43 to 45 weeks; in the Camel it is 13 months. In the largest Mammals the embryo requires a much longer time for its complete formation within the maternal body; in the Rhinoceros, for instance, $1\frac{1}{2}$ year, in the Elephant 90 weeks. In the latter case, therefore, gestation lasts more than twice as long as in Man—for

nearly a year and three quarters. In the smaller M the duration of embryonic development is, on the c much shorter. The smallest Mammals, the Harve develop fully in 3 weeks ; Rabbits and Hares in 4 Rats and Marmots in 5 weeks ; the Dog in 9, the P the Sheep in 21, and the Stag in 36 weeks. Develop yet more rapid in Birds. The Chick, under nor conditions of incubation, requires only 3 weeks, or just for its full development. The Duck, on the othe takes 25, the Turkey 27, the Peacock 31, the Swan the New Holland Cassowary 65 days. The smalle Birds, the Humming-bird, quits the egg after the twe It is, therefore, evident that in Mammals and in F duration of development within the egg-membrane in a definite relation to the size of body attained vertebrate species. But the latter is not the sole d ing cause of the former. There are many other stances which influence the duration of individual ment within the membranes of the egg.¹²⁶

In all cases, however, the duration of the O appears infinitely brief when compared with the er the infinitely long period during which the Phyl gradual development of the ancestral series, too This period is not to be measured by years and c but by thousands and millions of years. Many mi years must indeed have elapsed while the most vertebrate organism, Man, gradually developed f primæval one-celled ancestral organism. The oppo the development theory, who regard this gradual ment of Man from lower animal forms, and his descent from a one-celled primitive animal as in

do not reflect that the very same marvel actually recurs before our eyes in the short space of nine months during the embryonic development of each human individual. The same series of multifariously diverse forms, through which our brute ancestors passed in the course of many millions of years, has been traversed by every Man during the first 40 weeks of his individual existence within the maternal body.

All changes in organic forms, all metamorphoses of animal and plant forms, appear to us all the more remarkable and all the more wonderful in proportion as they occur more rapidly. When, therefore, our opponents pronounce that the past development of the human race from lower animal forms is incredible, they must regard the embryonic development of the human individual from the simple egg-cell as far more wonderful in comparison. This latter process—the ontogenetic modification—which takes place before our eyes, must appear more wonderful than the phylogenetic modification, in proportion as the duration of the tribal history exceeds that of the germ-history. For the human embryo must pass through the whole process of individual development, from the simple cell up to the many-celled perfect Man, with all his organs, in the brief space of 40 weeks. On the other hand, we may assign many millions of years for the accomplishment of the analogous process of phylogenetic development—the development of Man's ancestors from the simplest one-celled form.

As regards these phylogenetic periods, it is impossible to fix approximately their length in hundreds or in thousands of years, or to establish any absolute measure of their duration. But the researches of geologists have long since

enabled us to estimate and compare the relative durations of the various periods of the earth's organic history. The most direct standard for determining the relative duration of geological periods is afforded by the thickness of the so-called Neptunian strata or sedimentary rock, i.e., all those strata which have been deposited, as mud, at the bottom of the ocean, or under fresh water. These stratified sedimentary rocks—limestone, clay, marl, sandstone, slate, etc.—which constitute the great mass of mountain-chains, and which are often several thousand feet in thickness, afford us data for estimating the relative lengths of the various periods of the earth's history.

For the sake of completeness, I must say a few words as to the development of the earth as a whole, briefly indicating a few of the more prominent facts relating to this matter. At the very outset we are confronted with the weighty fact, that life originated on our planet at a certain definite period. This is a proposition that is no longer gainsaid by any competent geologist. We now know for certain that organic life upon our planet actually began at a certain time, and that it did not exist there from eternity, as some have supposed. The indisputable proofs of this are furnished, on the one hand, by physico-astronomical cosmogeny; on the other, by the Ontogeny of organisms. Species and tribes, like individuals, do not enjoy a perpetual life.¹⁸⁷ They also had a beginning. The time which has elapsed since the origin of life upon the earth up to the present time (and with this period of time alone we are here concerned) we call the "history of the organic earth," as distinguished from the "history of the inorganic earth" which embraces the period before the origin of organic life.

regard to the latter, we first obtained clear ideas from natural philosophical researches and computations of great critical philosopher, Immanuel Kant, and on this

I must refer the reader to Kant's "*Allgemeine Naturlichte und Theorie des Himmels*" and to the numerous *logienies* which treat the subject in a popular style. I cannot here dwell upon questions of this kind.

The organic history of the earth could begin only when water in fluid drops existed upon its surface. For the very existence of all organisms, without any exception, depends upon water in the fluid state, their bodies containing a considerable amount of the same. Our own body, in its fully developed state, contains in its tissues 70 per cent. of water and only 30 per cent. of solid matter. The amount of water is still greater in the body of the child, and is greatest of all in the embryo. In early stages of development the human embryo contains more than 90 per cent. of water, and not over 10 per cent. of solid matter. In low marine animals, especially in the *Medusæ*, the body contains even more than 95 per cent. of water, and not even one per cent. of solid matter. No organism can exist and perform its vital functions without water. Without water there is no life.

Water in the fluid state, which is, therefore, indispensable for the existence of life, could not, however, exist upon the earth until after the temperature of the surface of the fiery globe had sunk to a certain point. Before this it existed only in the form of steam. As soon as, however, as the first drop of water in a fluid state was precipitated by cooling from the envelope of steam, it began geological action, and from that time to this it has effected continual changes in the modification of the hard

crust of the earth. The result of this unceasing work of the water, which in the form of rain and hail, of snow and ice, of rushing torrent and surging wave crumbles and dissolves the rocks, is the formation of ooze. As Huxley says, in his excellent "Lectures on the Causes of the Phenomena of Organic Nature," the most important fact in the past history of our earth is ooze, and the question as to the history of the past ages of the world resolves itself into a question as to the formation of ooze. All the stratified rocks of our mountainous formations were originally deposited as ooze at the bottom of the waters, and only afterwards hardened into solid stone.

As has already been said, it is possible, by bringing together and comparing the various rock-strata from many places on the surface of the earth, to obtain an approximate conception of the relative ages of these various strata. Geologists have long agreed that there is an entirely definite historical sequence of the various formations. The various groups of strata which lie one over another correspond to successive periods in the earth's organic history, during which they were deposited in the shape of mud at the bottom of the sea. Gradually this mud was hardened into solid rock. The latter, by alternate upheaval and depression of the surface of the earth, was lifted above the water, and assumed the form of mountains. Four or five main periods in the earth's organic history, answering to the larger and smaller groups of these sedimentary rock-strata, are usually distinguished. These main periods are subdivided into numerous subordinate or lesser periods. From twelve to fifteen of the latter are usually assumed. (Cf. Tables XII. and XIII., pp. 11, 12.) The relative thick-

of the various groups of strata affords the means of approximately estimating the relative length of these various periods of time. Of course we cannot say, "In a hundred years on the average a stratum of a certain thickness (say two inches) is deposited, and therefore a rock-column of a thousand feet in thickness is 600,000 years old." For different rock-formations of equal thickness may have occupied periods of very various length during their deposition and consolidation. From the thickness of a formation we may, however, approximately judge of the *relative* length of the period during which it was deposited.

Of the four or five main periods of the earth's organic history, our acquaintance with which is indispensable for the Phylogeny of the human race, the first and oldest is known as the Primordial, Archizoic, or Archilithic Epoch. We estimate the total thickness of all the sedimentary rocks as averaging about 130,000 feet, then 70,000 feet belong to this first epoch—more than one half. From this and other circumstances we may conclude that the corresponding Primordial or Archilithic Epoch must alone have been considerably longer than the whole long period between the beginning of the Archilithic and the present time. Probably the Primordial Epoch was much longer than might appear from the ratio of 7 : 6, which we have given. This Epoch is divided into three sub-periods, known as the Laurentian, Cambrian, and Silurian, corresponding to the three principal groups of sedimentary rock-strata which constitute the Archilithic Era. The enormous length of time required for the formation at the bottom of the primordial sea of these gigantic strata, of over 70,000 feet in thickness, must, at all events,

have been many millions of years. During that time there came into existence by spontaneous generation the oldest and simplest organisms—those in which life began upon our planet—viz., the Monera. From these, one-celled plants and animals first developed—the Amœbæ and many kinds of Protista. During this same Archilithic Epoch, also, all the invertebrate ancestors of the human race developed from these one-celled organisms. We draw this conclusion from the fact that towards the close of the Silurian period a few remains of fossil Fishes are already to be found, viz., Selachians and Ganoids. These are, however, much more highly organized and of later origin than the lowest Vertebrates (the Amphioxus), or than the various skull-less Vertebrates allied to Amphioxus, which must have lived during this time. The latter must necessarily have been preceded by all the invertebrate ancestors of man. Hence we may characterize this entire epoch as the “age of man’s invertebrate ancestors;” or, with special reference to the oldest representatives of the Vertebrate tribe, as the “age of Skull-less Animals.” During the whole Archilithic Epoch the inhabitants of our planet consisted exclusively of aquatic forms; at least, no remains of terrestrial animals or plants dating from this period have as yet been found. A few remains of land-dwelling organisms which are sometimes referred to the Silurian Period, are Devonian.

The Primordial Epoch was followed by the Palæolithic, Palæozoic, or Primary Epoch, which is also separable into three sub-periods: the Devonian, the Carboniferous, and the Permian. During the Devonian Period the Old Red Sandstone, or Devonian system was formed; during the Carboniferous, those great beds of coal were deposited which

TABLE XII.

Systematic Survey of the Palæontological Periods, or the Greater Divisions
in the History of the Organic Earth.

I. First Epoch : The Archilithic, or Primordial Epoch.

(Age of Skull-less Animals and Seaweed Forests.)

Older Archilithic Epoch	or	Laurentian Period.
Middle Archilithic Epoch	"	Cambrian Period.
Later Archilithic Epoch	"	Silurian Period.

II. Second Epoch : The Palæolithie, or Primary Epoch.

(Age of Fishes and Fern Forests.)

Older Palæolithie Epoch	or	Devonian Period.
Middle Palæolithie Epoch	"	Coal Period.
Later Palæolithie Epoch	"	Permian Period.

III. Third Epoch : The Mesolithie, or Secondary Epoch.

(Age of Reptiles and Pine Forests, *Coniferae*.)

Older Mesolithie Epoch	or	Triassic Period.
Middle Mesolithie Epoch	"	Jurassic Period.
Later Mesolithie Epoch	"	Chalk Period.

IV. Fourth Epoch : The Cæolithie, or Tertiary Epoch.

(Age of Mammals and Leaf Forests.)

Older Cæolithie Epoch	or	Eocene Period.
Middle Cæolithie Epoch	"	Miocene Period.
Later Cæolithie Epoch	"	Pliocene Period.

V. Fifth Epoch : The Anthropolithie, or Quaternary Epoch.

(Age of Man and Cultivated Forests.)

Older Anthropolithie Epoch	or	Ice Age, Glacial Period.
Middle Anthropolithie Epoch	"	Post Glacial Period.
Later Anthropolithie Epoch	"	Period of Culture.
Period of Culture is the Historic Period, or Period of Tradition.)		

TABLE XIII.

Systematic Survey of the Palæontological Formations, or the Fossiliferous Strata of the Earth's Crust.

<i>Rock-Groups.</i>	<i>Systems.</i>	<i>Formations.</i>	<i>Synonyms of Formations.</i>
V. <i>Quaternary Group, or Anthropolithic (Anthropozoic) groups of strata</i>	XIV. Recent (Alluvium) XIII. Pleistocene (Diluvium)	36. Present 35. Recent 34. Post glacial 33. Glacial	Upper alluvial Lower alluvial Upper diluvial Lower diluvial
IV. <i>Tertiary Group, or Cænoolithic (Cænozoic) groups of strata</i>	XII. Pliocene (New tertiary) XI. Miocene (Middle tertiary) X. Eocene (Old tertiary)	32. Arvernian 31. Sub-Appenine 30. Falunian 29. Limburgian 28. Gypsum 27. Nummulitic 26. London clay	Upper pliocene Lower pliocene Upper miocene Lower miocene Upper eocene Middle eocene Lower eocene
III. <i>Secondary Group, or Mesolithic (Mesozoic) groups of strata</i>	IX. Cretaceous. VIII. Jura VII. Trias	25. White chalk 24. Green sand 23. Neocomian 22. Wealden 21. Portlandian 20. Oxfordian 19. Bath 18. Lias 17. Keuper 16. Muschelkalk 15. Bunter sand	Upper cretaceous Middle cretaceous Lower cretaceous The Kentish Weald Upper oolite Middle oolite Lower oolite Lias formation Upper trias Middle trias Lower trias
II. <i>Primary Group, or Palæolithic (Palæozoic) groups of strata</i>	VI. Permian (New red sandstone) V. Carboniferous (Coal) IV. Devonian (Old red sandstone)	14. Mountain limestone (Zechstein) 13. Red sandstone 12. Carboniferous sandstone 11. Carboniferous limestone 10. Pilton 9. Ilfracombe 8. Linton	Upper Permian Lower Permian Upper carboniferous Lower carboniferous Upper Devonian Middle Devonian Lower Devonian
I. <i>Primordial Group, or Archilithic (Archizoic) groups of strata</i>	III. Silurian II. Cambrian I. Laurentian	7. Ludlow 6. Wenlock 5. Llandeilo 4. Potsdam 3. Longmynd 2. Labrador 1. Ottawa	Upper Silurian Middle Silurian Lower Silurian Upper Cambrian Lower Cambrian Upper Laurentian Lower Laurentian

supply us with our principal fuel; in the Permian, the new Red Sandstone, the Magnesian Limestone (Zechstein), and the Cupriferous Slate were formed. The approximate thickness of this entire group of strata is estimated at 42,000 feet at most; some geologists make it somewhat more, others considerably less. In any case, this Palæolithic Epoch, taken as a whole, is considerably shorter than the Archilithic, but yet is considerably longer than all the following Epochs taken together. The strata deposited during this Primary Epoch supply fossil remains in great abundance; besides numerous species of Invertebrates we find also very many Vertebrates—Fishes preponderating. As early as the Devonian, and even during the Carboniferous and the Permian periods, there existed so great a number of Fishes, especially Primitive Fishes (Sharks) and Ganoids, that we may designate the entire Palæolithic Period as the *Age of Fishes*. The Palæozoic Ganoids especially are represented by a large number of forms.

But even during this period some Fishes began to custom themselves to living upon the land, and thus gave rise to the Amphibian class. Even in the carboniferous system we find fossil remains of Amphibia—the earliest terrestrial and air-breathing animals. In the Permian period the variety of these Amphibia becomes greater. Towards its close the first Amnion-animals, the tribal ancestors of the true higher Vertebrate classes, seem first to appear. These are a few lizard-like animals, of which the *Proteronotus* from the Cupriferous Slate at Eisenach is the best known. The appearance of the most ancient Amnion animals (*Amniota*), to which the common parent-form of

Reptiles, Birds, and Mammals must have belonged, not in fact to be referred by these oldest reptilian remains to the close of the Palæolithic Epoch. During this Epoch the ancestors of the human race must accordingly have been represented, first by true Fishes, then by Mud-Fish (*Dipneusta*) and Amphibia, and finally by the old Amnion Animals, the Protamnia.

After the Palæolithic Epoch comes a third main division of the earth's organic history, known as the Mesolithic or Secondary Epoch. This is again distinguished into three subdivisions—the Triassic, the Jurassic, and the Cretaceous Periods. The approximate thickness of the strata formed during these three Periods from the beginning of the Triassic down to the end of the Cretaceous Period amounts in all to about 15,000 feet, not one half the thickness of the Palæolithic deposits. During this Epoch a great and varied development took place in all divisions of the animal kingdom. In the vertebrate tribe especially a number of new and interesting forms developed. Among the Fishes the Osseous Fishes (*Teleostei*) now first appear. Among the Reptiles surpass all others both in numbers and diversity of species—the most remarkable and the most familiar forms being the gigantic extinct Dragons (Dinosaurs), the Sea-Dragons (Halisaurians), and the Flying Lizards (Pterosaurians). In reference to this predominant reptilian class this time is known as the *age of reptiles*. But the class of Birds also developed during this period undoubtedly originated from a branch of the lizard-Reptiles. This is shown by the similar embryology of Birds and of Reptiles, by their Comparative Anatomy, and also by the fact that we know of fossil birds with toothed jaws.

ed with lizard's tail, belonging to this period (*Odon-
nis Archæopteryx*). Finally, it was during this period
at there appeared upon the scene that most perfect and,
us, most important vertebrate class, the mammalian
a. The oldest fossil remains of these have been found
the most recent Triassic strata, viz., molar teeth of
small insectivorous Pouched Animal (Marsupial). Numer-
remains occur somewhat later in the Jura system, and
w in the chalk. All the remains of Mammals from this
olitic Epoch with which we are acquainted belong to
low Pouched Animal division; and among these were
ubtly the ancestors of Man. On the other hand,
a single undisputed relic has yet been discovered
ughout all this period of one of the higher Mammals
centalia). This last division, of which Man is a member,
not develop till later, in the immediately subsequent
ary Epoch.

The fourth main division of the history of the organic
a, the Tertiary, Cænozoic, or Cænolithic Epoch, was of
a shorter duration than the preceding. For the strata
sited during this period are in all only about 3000 feet
ickness. This Epoch, also, is divided into three sub-
ions, known as the Eocene, Miocene, and Pliocene
ods. During these periods the most diverse develop-
t of the higher classes of plants and animals took place
the fauna and flora of our globe now approached
er and nearer to their present character. The most
ly developed class of animals, that of Mammals, now
ned pre-eminence. This Tertiary Epoch may, therefore,
alled the *age of Mammals*. The most perfect section
his class, the Placental Animals, among which is Man,

now first appeared. The first appearance of Man, speak more correctly—the development of man from most nearly allied ape-form, dates probably either from Miocene or the Pliocene Period,—from the middle latest section of the Tertiary Epoch. Perhaps, as is said by others, Man strictly so-called, *i.e.*, Man gifted with language, first developed from the speechless man-like in the subsequent Anthropolithic Age.

At all events, the perfect development and distribution of the various races of Man dates from the fifth and last main division of the organic history of the earth, and this Epoch has been called the Anthropolithic, or Anthropozoic, and also the Quaternary Epoch. It is true that in the present imperfect state of our palæontological and prehistoric knowledge, we cannot solve the problem whether the development of Man from the nearest Ape-forms took place in the beginning of this Anthropozoic Epoch, or as early as the middle or towards the close of the preceding Tertiary Epoch. This much, however, is certain, that the true development of human culture dates from the Anthropolithic Epoch, and that this latter constitutes only an insignificantly small section of the enormous period of time occupied in the development of the organic earth. When we reflect upon this, it is absurd to speak of the brief span of man's period of culture as "the world's history." This so-called History of the World does not amount approximately to even half per cent. of the length of those enormous periods which have passed away from the beginning of the organic history down to the present time. For this V

History, or more correctly, History of People, is itself only the latter half of the Anthropolithic Epoch, while even the first half of this Epoch must be reckoned as a prehistoric period. Hence this last main period, reaching from the close of the Cærolithic Epoch to the present time, can only be called the "age of man," inasmuch as the diffusion and differentiation of the different species and races of man, which have so powerfully influenced all the rest of the organic population of the globe, took place during its course.

Since the awakening of the human consciousness, human vanity and human arrogance have delighted in regarding Man as the real main-purpose and end of all earthly life, and as the centre of terrestrial Nature, for whose use and service all the activities of the rest of creation were from the first defined or predestined by a "wise providence." How utterly baseless these presumptuous anthropocentric conceptions are, nothing could evince more strikingly than comparison of the duration of the Anthropolzoic or Quaternary Epoch with that of the preceding Epochs. For even though the Anthropolithic Epoch may embrace several hundreds of thousands of years, how small is this time when compared with the millions of years that have elapsed since the beginning of the world's organic history down to the first appearance of the human race!

If the entire duration of the organic history of the earth, from the generation of the first Monera down to the present day, is divided into a hundred equal parts, and if then, corresponding with the relative average thickness of the intervening strata-systems, the respective percentages are

assigned to the relative durations of the five main divisions or Epochs, the latter will be found to be about as follows:—

I. Archilithic, or archizoic (primordial) Epoch	. .	53.6
II. Palæolithic, or palæozoic (primary) Epoch	. .	32.1
III. Mesolithic, or mesozoic (secondary) Epoch	. .	11.5
IV. Cæolithic, or cæozoic (tertiary) Epoch	. .	2.3
V. Anthropolithic, or anthropozoic (quaternary) Epoch	. .	0.5
Total ...		100.0

The relative durations of the five main epochs of the earth's organic history, are yet more clearly seen in the opposite Table (XIV.), in which the relative thicknesses of the strata systems deposited within these Epochs is represented on a scale corresponding to their actual depths.

This table shows that the period of the so-called History of the World forms but an inconsiderable span in comparison with the immeasurable duration of those earlier epochs during which Man did not exist upon this planet. Even the great Cæozoic Epoch, the so-called Tertiary Epoch, during which the Placental Animals, the higher Mammals, developed includes but little more than two per cent. of the whole enormous duration of the organic history of the world.¹²⁸

And now before we turn to our proper phylogenetic task; before, guided by our knowledge of ontogenetic facts and by the fundamental law of Biogeny, we attempt to trace step by step the history of the palæontological evolution of our animal ancestors, let us turn aside for a short time into another and apparently very different and very remote department of science, a general review of which will make the solution of the difficult problems which now rise before us very much easier. The science is the

TABLE XIV.

matic Survey of the Neptunian fossiliferous strata of the earth with
reference to their relative sectional thickness (130,000 feet circa).

IV. Cenolithic Strata, circa 3000 feet.	Pliocene, Miocene, Eocene.
III. Mesolithic Strata. Deposits of the Secondary Epoch, circa 15,000 feet.	IX. Chalk System. VIII. Jurassic System. VII. Triassic System
II. Palæolithic Strata. Deposits of the Primary Epoch, circa 42,000 feet.	VI. Permian System. V. Coal System. IV. Devonian System.
I. Archilithic Strata. Deposits of the Primordial Epoch, circa 70,000 feet.	III. Silurian System, circa 22,000 feet. II. Cambrian System, circa 18,000 feet. I. Laurentian System, circa 30,000 feet.

of Comparative Philology. Ever since Darwin, by the theory of Natural Selection, infused new life into Biology, and raised the fundamental question of development in every branch of science, attention has frequently and from very different quarters been called to the remarkable parallelism, which exists between the evolution of the various human languages and the evolution of organic species. The comparison is quite justifiable and very instructive. Indeed it is hardly possible to find an analogy better adapted to throw a clear light on many obscure and difficult facts in the evolution of species, which is governed and directed by the same natural laws which guide the course of the evolution of language.

All philologists who have made any progress in their science, now unanimously agree that all human languages have developed slowly and by degrees from the simplest rudiments. On the other hand, the strange proposition which till thirty years ago was defended by eminent authorities, that language is a divine gift, is now universally rejected, or at best defended only by theologians and by people who have no conception of natural evolution. Such brilliant results have been attained in Comparative Philology that only one who is wilfully blind can fail to recognize the natural evolution of language. The latter is necessarily evident to the student of nature. For speech is a physiological function of the human organism, developing simultaneously with its special organs, the larynx and the tongue, and simultaneously with the functions of the brain. It is, therefore, quite natural that in the history of the evolution of languages, and in their whole system, we should find the same correlations as in the history of the evolution of organic species and in their whole system. The various


larger and smaller groups of speech-forms, which are distinguished in Comparative Philology as primitive languages, fundamental languages, parent languages, derived languages, dialects, patois, etc., correspond perfectly in their mode of development with the various larger and smaller groups of organisms classed in systems of Zoology and Botany as tribes, classes, orders, families, genera, species, and varieties of the animal and vegetable kingdoms. The relations between these various systematic groups, or categories, are in both cases identical; moreover, evolution follows the same course in one case as in the other. This instructive comparison was first elaborated by one of the most eminent of German philologists, one who, unfortunately, died prematurely—August Schleicher, not only a philologist but also a learned botanist. In his more important works, the Comparative Anatomy and evolutionary history of languages is treated by the same phylogenetic method which we employ in the Comparative Anatomy and evolutionary history of animal forms. He has especially applied this method to the Indo-Germanic family of languages; and in his little treatise on "The Darwinian Theory and the Science of Language" ("Die Darwin'sche Theorie und die Sprachwissenschaft"), he illustrated it by means of a synoptical pedigree of the Indo-Germanic family of languages.¹²⁰

If the formation of the various branch languages which have developed from the common root of the primitive Indo-Germanic tongue is studied with the aid of this pedigree, a very clear idea of their Phylogeny will be acquired. At the same time it becomes evident how entirely analogous is the evolution of the greater and lesser groups of the Vertebrates, which have sprung from the one common root.

form of the primitive Vertebrates. The primitive Indo-Germanic root-tongue first separated into two chief stems: the Slavo-Germanic and the Ario-Romanic main-trunks. The Slavo-Germanic then branched into a primitive German and a primitive Slavo-Lettic tongue. Similarly, the Ario-Romanic split up into a primitive Arian and a primitive Græco-Romanic language (p. 23). If we continue our examination of this pedigree of the four primitive Indo-Germanic languages, we find that the primitive Germanic tongue divided into three chief branches—a Scandinavian, a Gothic, and a Teutonic branch. From the Teutonic branch proceeded, on the one hand, High German, and, on the other hand, Low German, to which latter belong the various Friesian, Saxon, and Low German dialects. Similarly, the Slavo-Lettic tongue developed first into a Baltic and a Slavonic language. From the Baltic spring the Lettic, Lithuanian, and Old Prussian dialects. The Slavic, on the other hand, give rise, in the South-east, to the Russian and the South Slavic dialects, and, in the West, to the Polish and Czech dialects.

Turning now to the other main stem of the Indo-Germanic languages and its branches—the primitive Ario-Romanic—it is found to develop with the same luxuriance. The primitive Græco-Romanic language gave rise, on the one hand, to the Thracian language (Albanian Greek), and on the other, to the Italo-Keltic. From the latter in turn sprung two divergent branches—in the South, the Italian branch (Romanic and Latin), and in the North, the Keltic, from which arose all the different British (Old British, Old Scottish, and Irish) and Gallic tongues. The numerous Iranian and Indian dialects branched out in the same way from the primitive Arian language.

Pedigree of the Indo-Germanic Languages.



A close study of this pedigree of the Indo-Germanic languages is, in many respects, of great interest. Comparative Philology, to which we are indebted for our knowledge of this subject, thus shows itself to be a true science—a *natural science*. It, indeed, long ago anticipated in its own province the phylogenetic method with the aid of which we now attain the highest results in Zoology and in Botany. And here I cannot refrain from remarking how much to the advantage of our general culture it would be if the study of languages (which is undoubtedly one of the most powerful means of culture) were *comparatively* prosecuted; and if our cut and dried Philology were replaced by a living, many-sided, *comparative* study of languages. The latter stands in the same relation to the former as the living history of organic evolution does to the lifeless classification of species. How much deeper would the interest taken in the study of language by the students in our schools be, and how much more vivid would be the results if even the first elements of Comparative Philology were taught instead of the distasteful composition of Latin exercises in Ciceronian style!

I have entered with this detail into the "Comparative Anatomy" and the history of the evolution of languages, because it is unsurpassed as a means of illustrating the Phylogeny of organic species. We find that in structure and in development these primitive languages, parent languages, derived languages, and dialects, correspond exactly like the classes, orders, genera, and species of the animal kingdom. The "natural system" is in both cases phylogenetic. Just as Comparative Anatomy, Ontogeny, and Palæontology afford certain proof that all Vertebrates,

er extinct or extant, are descended from a common
ral form, so does the comparative study of the dead
iving Indo-Germanic language absolutely convince
at all these languages have sprung from a common
. This monophyletic view is unanimously adopted by
guists of importance who have studied the question,
who are capable of passing a critical judgment upon it.¹³⁰
The point, however, to which I would specially call your
attention in this comparison between the various branches,
on the one hand, of the Indo-Germanic language, and, on the
other, of the vertebrate tribe, is that the direct descendants
never be confounded with the collateral lines, nor the
extinct with the extant forms. This mistake is often
made, and results in the formation of erroneous notions of
descent. Our opponents often take advantage in order to
disprove the whole theory of descent. When, for instance, it
is said that human beings are descended from Apes, the
from Semi-apes, and the Semi-apes from Pouched
Mammals (*Marsupialia*), very many people think only of the
familiar living species of these different mammalian orders,
as are to be found stuffed in our museums. Now, our
opponents attribute this erroneous view to us, and, with
more craft than judgment, declare the thing impossible; or
they ask us as a physiological experiment to transform
a kangaroo into a Semi-ape, this into a Gorilla, and the
Gorilla into a Man. Their demand is as childish as the
supposition on which it is founded is erroneous; for all
extant forms have varied more or less from their
parent-form, and none of them are capable of pro-
ducing the same divergent posterity which were really
produced thousands of years ago by that parent-form.¹³¹

There is no doubt that Man is descended from an extinct mammalian form, which, if we could see it, we should certainly class with the Apes. It is equally certain that this primitive Ape in turn descended from an unknown Semi-ape, and the latter from an extinct Pouched Ape. But then it is beyond a doubt that it is only in respect of essential internal structure, and on account of their similarity in the distinctive anatomical characters of the order, that these extinct ancestral forms can be spoken of as members of the yet extant mammalian orders. In external form, generic and specific characters, they must have been more or less—perhaps even greatly—different from all living representatives of the orders to which they belonged. For it must be accepted as a quite universal and natural fact of phylogenetic evolution that the parent-forms themselves, with their specific characters, became extinct at a more or less distant period. Those extant forms which come nearest to them, yet differ from them more or less, perhaps even very essentially. Hence in our phylogenetic research and in our comparative view of the still living divergent descendants all we can undertake to do is to determine how far the latter depart from the parent-form. We can quite confidently assume that no single older parent-form has reproduced itself without modification down to our times.

We find this same state of things on comparing various extinct and living languages, which have sprung from a common primitive tongue. If, from this point of view, we examine the genealogical tree of the Indo-Germanic languages, we may conclude, on *a priori* grounds, that the earlier primitive languages, fundamental languages, or ancestral languages, from which the living dialects

ended in the first or second degree, have been extinct longer or shorter period. And this is the case. The Romanic and the Sclavo-Germanic tongues have long altogether extinct, as are also the primitive Arian and o-Romanic, the Sclavo-Lettic, and primitive Germanic ages. Some even of the languages descended from have also long been dead, and all those of the Indo-anic branch which are yet extant, are akin only in as they are divergent descendants of common parent. Some have diverged from this ancestral form more, less.

This easily demonstrable fact very well illustrates analogous facts in the descent of vertebrate species. Phyletic "Comparative Philology," as a powerful ally, supplementary "Comparative Zoology." The former can, however, adduce far more direct evidence than the latter, as the palæontological materials of Philology, the great monuments of extinct tongues, have been far better preserved than the palæontological materials of Comparative Zoology, the fossil bones of vertebrates. The more these analogous conditions are considered, the more convincing is their force.

We shall presently find that we can trace back the logical line of Man, not only to the lower Mammals, even to the Amphibia, to the shark-like Primitive Fishes, even far below these, to the skull-less Vertebrates allied to *Amphioxus*. It must be remembered this does not mean that the living *Amphioxus*, Shark, or Amphibian accurately represent the outward appearance of the parent-forms which we speak. Still less does it mean that the *Amphioxus*, or the Shark of our day, or any extant species

Here arises another question : " Whence, in the beginning of the organic history of the earth, at the commencement of the Laurentian period, came the earliest Amœbæ ? " To this there is but one reply. Like all one-celled organisms, the Amœbæ have originally developed only from the simplest organisms known to us, the Monera. These Monera, which we have already described, are also the simplest conceivable organisms. Their body has no definite form, and is but a particle of primitive slime (plasson)—a little mass of living albumen, performing all the essential functions of life, and everywhere met with as the material basis of life. This brings us to the last, or perhaps the first question in the history of evolution—the question as to the origin of the Monera. And this is the momentous question as to the prime origin of life—the question of spontaneous generation (*generatio spontanea* or *œquivoca*).

We have neither time, nor indeed have we any occasion, to discuss at length the weighty question of spontaneous generation. On this subject I must refer you to my "History of Creation," and, especially, to the second book of the *Generelle Morphologie*, and to the discussion on Monera and spontaneous generation in my "Studien über Moneren und andere Protista."¹⁸² I have there stated my own views on this important subject in very great detail. Here I will only say a few words on the obscure question as to the first origin of life, and will answer it so far as it concerns our radical conception of the history of organic evolution. In the definite, limited sense in which I maintain spontaneous generation (*generatio spontanea*) and assume it as a necessary hypothesis in explanation of the first beginning of life upon

earth, it merely implies the origin of Monera from organic carbon compounds. When animated bodies first appeared on our planet, previously without life, there must, in the first place, have been formed, by a process purely chemical, from purely inorganic carbon combinations, that very complex nitrogenized carbon compound which we call murex, or "primitive slime," and which is the oldest material substance in which all vital activities are embodied. In the lowest depths of the sea such homogeneous amorphous protoplasm probably still lives, in its simplest character, under the name of Bathybius.¹³⁷ Each individual living particle of this structureless mass is called a Moneron. The oldest Monera originated in the sea by spontaneous generation, just as crystals form in the matrix. This assumption is required by the demand of the human understanding for causality. For when, on the one hand, we reflect that the whole inorganic history of the earth proceeds in accordance with mechanical laws and without any intervention by creative power, and when, on the other hand, we consider that the entire organic history of the world is also determined by similar mechanical laws; when we see that no supernatural interference by a creative power is needed for the production of the various organisms, then it is certainly quite inconsistent to assume such supernatural creative interference for the first production of life upon our globe. In all events we, as investigators of nature, are bound at least to attempt a natural explanation.

At present, the much agitated question of spontaneous generation appears very intricate, because a large number of very different, and in part quite absurd, conceptions are included under the term "spontaneous generation," and

because some have supposed that the problem could be solved by means of the crudest experiments. The doctrine of spontaneous generation cannot be experimentally refuted. For each experiment with a negative result merely proves that under the conditions (always very artificial) supplied by us, no organism has been produced from inorganic combinations. Neither can the theory of spontaneous generation be experimentally proved unless great difficulties are overcome; and even if in our own time Monera were produced daily by spontaneous generation—as is very possible—yet the absolute empiric proof of this fact would be extremely difficult—indeed, in most cases impossible. He, however, who does not assume a spontaneous generation of Monera, in the sense here indicated, to explain the first origin of life upon our earth, has no other resource but to believe in a supernatural miracle; and this, in fact, is the questionable standpoint still taken by many so-called “exact naturalists,” who thus renounce their own reason.

Sir William Thomson has indeed tried to avoid the necessary hypothesis of spontaneous generation by assuming that the organic inhabitants of our earth originally descended from germs which proceeded from the inhabitants of other planets, and which, with fragments of the latter, with meteorites, accidentally fell on to the earth. This hypothesis has met with much applause, and was even supported by Helmholtz. Friederich Zoellner, an acute physicist, has, however, refuted it in his excellent natural-philosophical work “Ueber die Natur der Cometen,” a critical book containing most valuable contributions to the history and theory of knowledge.¹²⁷ Zoellner has plainly shown that the hypothesis is unscientific in two respects—

rstly, in point of logic, and secondly, in its scientific tenor (p. xxvi). At the same time he rightly shows that the hypothesis of spontaneous generation, in the sense which we have defined, is the "condition necessary to the conceivability of nature in accordance with the laws of causality."

In conclusion, I repeat, with emphasis, that it is only in the case of Monera—of structureless organisms without organs—that we can assume the hypothesis of spontaneous generation. Every differentiated organism, every organism composed of organs, can only have originated from an undifferentiated lower organism by differentiation of its parts, and consequently by Phylogeny. Hence, even in the production of the simplest cell we must not assume the process of spontaneous generation. For even the simplest cell consists of at least two distinct constituent parts; the inner and firmer kernel (nucleus), and the outer and softer cell-substance or protoplasm. These two distinct parts can only have come into being by differentiation of the homogeneous plasson of a moneron and of a cytod. It is for this very reason that the natural history of Monera is of the highest interest; for it alone can remove the principal difficulties which beset the question of spontaneous generation. The extant Monera do afford us organless and structureless organisms, such as must have originated by spontaneous generation at the first beginning of organic life upon the earth.¹³³

CHAPTER XVI.

THE ANCESTRY OF MAN.

I. FROM THE MONERA TO THE GASTRÆA.

Relation of the General Inductive Law of the Theory of Descent to Special Deductive Laws of the Hypotheses of Descent.—Incompleteness of the Three Great Records of Creation: Palæontology, Ontogeny and Comparative Anatomy.—Unequal Certainty of the Various Scientific Hypotheses of Descent.—The Ancestral Line of Men in Twenty-two Stages: Eight Invertebrate and Fourteen Vertebrate Ancestors.—Distribution of these Twenty-two Parent-forms in the Five Main Divisions of the Organic History of the Earth.—First Ancestral Stage: Monera. The Structureless and Homogeneous Plasson of the Monera.—Differentiation of the Plasson into Nucleus, and the Protoplasm of the Cytods and Cells as Two Different Plastid-forms.—Vital Phenomena of Monera.—Organisms without Organs.—Second Ancestral Stage: Amœbæ.—One-celled Primitive Animals of the Simplest and most undifferentiated Nature.—The Amœboid Egg-cells.—The Egg is Older than the Hen.—Third Ancestral Stage: Syn-Amœba, Ontogenetically reproduced in the Morula.—A Community of Homogeneous Amœboid Cells.—Fourth Ancestral Stage: Planæa, Ontogenetically reproduced in the Blastula or Planula.—Fifth Ancestral Stage: Gastræa, Ontogenetically reproduced in the Gastrula and the Two-layered Germ-disc.—Orientation of the Gastræa by Inversion (*invaginatio*) of the Planæa.—Haliphys and Gastrophysæma.—Extant Gastræads.

“Now, very probably, if the course of evolution proves to be so simple, it will be thought that the whole matter is self-evident, and no research is hardly required to establish it. But the story of Columbus and the egg is daily repeated; and it is necessary to perform the exper-

one's self. How slowly progress is made in the knowledge even of self-evident matters, especially when respectable authorities disagree, I myself have experienced sufficiently."—KARL ERNST BAER (1828).

GUIDED by the fundamental law of Biogeny and by the records of creation, we now turn to the interesting task of examining the animal parent-forms of the human race in their proper sequence. To ensure accuracy, we must first become acquainted with the various mental operations which we shall apply in this natural-philosophical research. These operations are partly of an inductive, partly of a deductive nature; partly conclusions from numerous particular experiences to a general law; partly conclusions from this general law back to particular experiences.

Tribal history as a whole is an inductive science; for the whole theory of descent, as an indispensable and most essential part of the whole theory of evolution, is entirely founded on inductions. From all the biological incidents in plant life, in animal life, and in human life, we have derived the certain inductive conception that the whole of the organic inhabitants of our globe originated in accordance with a single law of evolution. To this law of evolution, Lamarck, Darwin, and their successors gave definite form in the theory of descent. All the interesting phenomena exhibited by Ontogeny, Palæontology, Comparative Anatomy, Embryology, Chorology, the Œcology of organisms, all the important general laws, which we infer from multitudinous phenomena of these different sciences, and which are most intimately connected together, are the broad inductive data from which is drawn the most extensive inductive view of Biology. Because the innate connection between all

these infinitely various groups of phenomena in these different departments becomes explicable and comprehensible solely through the theory of descent, therefore this theory of evolution must be regarded as an extensive inductive law. If we now really apply this inductive law, and its help seek to discover the descent of individual organisms or species, we must necessarily form phylogenetic hypotheses which are of an essentially deductive nature, and which are inferences from the general theory of descent back to individual particular cases. These special deductive conclusions are, however, in accordance with the inexorable law of Logic, as justifiable, as necessary, and as indispensable to our department of knowledge as the general inductive conclusions of which the whole theory of evolution is formed. The doctrine of the animal parent-forms of a kind is also a special deductive law of this kind, which is the logical conclusion from the general inductive law of the theory of descent.¹⁸⁴

As is now very generally acknowledged, both by the adherents of and the opponents of the theory of descent, the choice, in the matter of the origin of the human race, lies between two radically different assumptions: We may either accustom ourselves to the idea that all the various species of animals and plants, Man also included, originated independently of each other by the supernatural process of a divine "creation," which as such is entirely removed from the sphere of scientific observation—or we are compelled to accept the theory of descent in its entirety and trace the human race, equally with the various animal and plant species, from an entirely simple primæval parent form. Between these two assumptions there is no t

course. Either a blind belief in creation, or a scientific theory of evolution. By assuming the latter, and this is the only possible natural-scientific conception of the universe, we are enabled, with the help of Comparative Anatomy and Ontogeny, to recognize the human ancestral line with a certain approximate degree of certainty, just as is more or less the case with respect to all other organisms. Our previous study of the Comparative Anatomy and Ontogeny of Man, and of other Vertebrates, has made it quite clear that we must first seek the pedigree of mankind in that of the vertebrate tribe. There can be no doubt that (if the theory of descent is correct) Man has developed as a true Vertebrate, and that he originated from one and the same common parent-form with all other Vertebrates. This special deduction must be regarded as quite certain, correctness of the inductive law of the theory of descent being of course first granted. No single adherent of the latter can raise a doubt about this important deductive conclusion. We can, moreover, name a series of different forms of the vertebrate tribe, which may be safely regarded as the representatives of different successive phylogenetic stages of evolution, or as different members of the human ancestral line. We can also prove with equal certainty that the vertebrate tribe as a whole originated from a group of low invertebrate animal forms; and among these we can again with more or less certainty recognize a series of members of the ancestral chain.

We must, however, at once expressly say that the certainty of the different hypotheses of descent, which are founded entirely on special deductive inferences, is very unequal. Several of these conclusions are already fully

established ; others, on the contrary, are most doubtful ; in yet others, it depends upon the subjective proportion of the knowledge of the naturalist and on his capability of drawing conclusions, what degree of probability he will accord to them. It is, at all events, necessary thoroughly to distinguish between the *absolute* certainty of the general (inductive) theory of descent, and the *relative* certainty of the special (deductive) hypothesis of descent. We can never in any case prove the whole ancestral line of ancestors of an organism with the same certainty with which we regard the theory of descent as the only scientific explanation of the organic forms. On the contrary, the special proof of all separate parent-forms must always remain more or less incomplete and hypothetical. That is quite natural. For all the records of creation upon which we rely are in a great measure incomplete, and will always remain incomplete ; just as in the case of Comparative Philology.

Above all, Palæontology, the most ancient of all records of creation, is in the highest degree incomplete. We know that all the petrifications with which we are acquainted form but an insignificantly small fragment of the whole number of animal forms and plant forms which have ever existed. For each extinct species obtained by us in a petrified condition, there are at least a hundred, probably thousands of extinct species which have left no trace of their existence. This extreme and most deplorable defectiveness of the palæontological record of creation, upon which it is impossible to insist too strongly, is very easily accounted for. The very conditions under which organic remains become petrified necessitate it. It is also partly explicable as the result of

imperfect knowledge in this department. It must be remembered, that far the greater proportion of the rock strata which constitute the mountain masses of the surface of the earth is not yet unfolded to us. Of the countless petrifications which are hidden in the huge mountains in chains of Asia and Africa, we know but a few small examples. Part of Europe and of North America has alone been more minutely explored. The whole of the petrifications accurately known and in our collections do not amount to a hundredth part of those which really exist in the crust of the earth. In this respect we may, therefore, expect a rich harvest of discoveries in the future. But, in spite of this, the palæontological record of creation (for reasons which I have amply explained in Chapter XV. of my "Natural History of Creation") will always remain extremely incomplete.

Not less incomplete is the second, most important record of creation, that of Ontogeny. For the Phylogeny of the individual it is the most important of all. Yet, it also has great defects, and often leaves us in the lurch. In this matter, we must distinguish quite clearly between palæontological and kenogenetic phenomena, between the original, inherited evolution and the later, vitiated evolution. We must never forget that the laws of abridged and vitiated heredity frequently disguise the original course of evolution beyond recognition. The reproduction of the Phylogeny of the Ontogeny is but rarely tolerably complete. The earliest and most important stages of germ-history are usually the most abridged and compressed. The youthful evolutionary forms have in turn often adapted themselves to new conditions, and have thus been modified. The



struggle for existence has excited an equally strong modifying influence upon the various independent and yet undeveloped young forms, as upon the developed and mature forms. Therefore, in the Ontogeny of the higher animal forms, the Phylogeny has been very greatly limited by Keno-genesis ; as a rule, only a blurred and much vitiated picture of the original course of evolution of their ancestors now lies before us in the Ontogeny. Only with great precaution and judgment dare we infer the tribal history directly from the germ-history. Moreover, the germ-history itself is known to us only in the case of very few species.

Lastly, the highly important record of creation afforded by Comparative Anatomy is unfortunately very incomplete, and for the simple reason, that the number of extant animal species forms but a very small fragment of the whole number of different animal forms that have existed from the beginning of the organic history of the world to the present time. The total sum of the latter may safely be estimated at several millions. The number of those animals the organization of which has at present been investigated by Comparative Anatomy is very small in proportion. The more extended investigations of the future will, here also, open up unexpected treasures.

In view of this evident and natural incompleteness of the most important records of creation, we must of course take good care, in the tribal history of Man, not to lay too great weight on single known animal forms, nor with equal certainty to consider all the stages of evolution which come under our consideration, as parent-forms. On the contrary, in hypothetically arranging our ancestral line, we must take good care to remember that the single hypothetical

parent-forms are of very diverse values in relation to the certainty of our knowledge. From the few remarks which, while speaking of the Ontogeny, we made as to the corresponding phylogenetic forms, it will have been understood that some germ-forms may with certainty be regarded as reproductions of corresponding parent-forms. We recognized the human egg-cell and the parent-cell which results from the impregnation of the latter as the first and most important form of this kind.

From the weighty fact that the egg of the human being, like the egg of all other animals, is a simple cell, it may be quite certainly inferred that a one-celled parent-form once existed, from which all the many-celled animals, Man included, developed.

A second very significant germ-form, which evidently reproduces a primæval parent-form, is the germ-vesicle (*Blastula*), a simple hollow sphere, the wall of which consists of a single cell-stratum. A third extremely important form in germ-history, which may be quite safely and directly referred back to the tribal history, is the true *Gastrula*. This most interesting larval form already exhibits the animal body composed of two germ-layers, and furnished with the fundamental primitive organ, the intestinal canal. Now, as the same two-layered germ-condition, with the primitive rudiment of the intestinal canal, is common to all the other animal tribes (with the single exception of the Primitive Animals, *Protozoa*), we may certainly from this infer a common parent-form of similar construction to the *Gastrula*, the *Gastræa*. Equally important in their bearing on the Phylogeny of Man, are the very important ontogenetical form conditions which correspond to certain Worms,

Skull-less Animals (*Acrania*), Fishes, etc., etc. On the other hand, between these quite certain and most valuable phylogenetic points, great gaps in our knowledge unfortunately exist, with which we shall again and again meet, and which are satisfactorily explained by reasons which have already been named, especially by the incompleteness of Palæontology, of Comparative Anatomy, and Ontogeny.

In the first attempts to construct the human ancestral line, which I made in my *Generelle Morphologie*, and in the "Natural History of Creation," I arranged first ten, and, later, twenty-two different animal forms, which, with more or less certainty, may be regarded as the animal ancestors of the human race, and which must be looked upon as in a sense the most important stages of evolution in the long evolutionary series from the one-celled organisms up to Man.¹⁸⁵ Of these twenty to twenty-two animal forms, about eight fall within the older division of the Invertebrates, while twelve to fourteen belong to the more recent Vertebrate division. How these twenty-two most important parent-forms in the human ancestral line are distributed through the five main periods of the organic history of the earth, is shown in the following Table (XVI). At least half of these twenty-two stages of evolution (that is, the eleven oldest ancestral forms) are found within the Archilithic Epoch, within that first main period of the organic history of the earth, which includes the larger half of the latter, and during which probably only aquatic organisms existed. The eleven remaining parent-forms fall within the four remaining main Epochs: three within the Palæolithic Epoch, three within the Mesolithic Epoch, and four within the Cænolithic Epoch. In the last, the Anthropolithic Age, Man already existed.

If we would now undertake the difficult attempt to discover the phylogenetic course of evolution of these twenty-two human ancestral stages from the very commencement of life, and if we venture to lift the dark veil which covers the oldest secrets of the organic history of the earth, we must undoubtedly seek the first beginning of life among those wonderful living beings which, under the name of Monera, we have already frequently pointed out as the simplest known organisms. They are, at the same time, the simplest conceivable organisms; for their entire body, in its fully developed and freely moving condition, consists merely of a small piece of structureless primitive slime or plasson, of a small fragment of that extraordinarily important nitrogenous carbon compound, which is now universally esteemed the most important material substratum of all the active phenomena of life. The experiences of the last ten years particularly have convinced us with more and more certainty that wherever a natural body exhibits the active phenomena of life, nutrition, propagation, spontaneous movement, and sensation, a nitrogenous carbon compound, belonging to the chemical group of albuminous bodies, is always active, and represents the material substance through which these vital activities are effected. Whether, in a monistic sense, we conceive the function as the direct effect of the formed material substance, or, in a dualistic sense, we regard "Matter and Force" as distinct, it is at least certain that, hitherto, no living organism has been observed in which the exercise of vital activities was not inseparably connected with a plasson-body. In the Monera, the simplest conceivable organisms, the whole body consists merely of plasson, corresponding to the "primitive slime" of earlier natural philosophy.

TABLE XVI.

Systematic Survey of the most Important Stages in the Animal
Ancestral Line of Man.

M N = Boundary between the Invertebrate and the Vertebrate Ancestors.

<i>Epochs of the Organic History of the Earth.</i>	<i>Geological Periods of the Organic History of the Earth.</i>	<i>Animal Ancestral Stages of Man.</i>	<i>Nearest Living Relations of the Ancestral Stage.</i>
I. Archilithic or Primordial Epoch	1. Laurentian Period 2. Cambrian Period 3. Silurian Period	1. Monera (<i>Monera</i>)	Rathybus Protomaba
		2. Oldest Amoebae	Simple Amoebae (<i>Amoeba</i>)
		3. Amœboid Societies (<i>Synamœbia</i>)	Mœnula larva
		4. Ciliated planulae (<i>Planulae</i>)	Blastula larva
		5. Primitive Intes- tinal animals (<i>Gastrædæ</i>)	Gastrula larva
		6. Primitive Worms (<i>Archeimintæ</i>)	Girdling Worms (<i>Turbellaria</i>)
		7. Soft-worms (<i>Scolecida</i>)	? Between the gird- ling worms and the Sea-squirrels
		8. Chorda animals (<i>Chordonia</i>)	Sea-squirrels (<i>Ascidia</i>) (<i>Appendicularia</i>)
		M.....N	
		9. Skull-less animals (<i>Acrania</i>)	Lancelets (<i>Amphioxus</i>)
		10. Round-mouths (<i>Cyclostomæ</i>)	Lampreys (<i>Petromyzontes</i>)
II. Palæolithic or Primary Epoch	4. Devonian Period 5. Coal Period 6. Permian Period	11. Primitive Fishes (<i>Selachii</i>)	Sharks (<i>Squalaceæ</i>)
		12. Salamander Fishes (<i>Dipneusta</i>)	Mud fish (<i>Protoptera</i>)
		13. Gilled Amphibia (<i>Saxobranchia</i>)	Siren (<i>Protos</i>) and Axolotl (<i>Sirenia</i>)
		14. Tailed Amphibia (<i>Nosura</i>)	Water-newt (<i>Triton</i>)
III. Mesolithic or Secondary Epoch	7. Triassic Period 8. Jurassic Period 9. Chalk Period	15. Primitive Am- niota (<i>Protamnion</i>)	? Between Tailed Amphibians and Beaked animals
		16. Primitive Mam- mals (<i>Protomammalia</i>)	Beaked animals (<i>Monotremes</i>)
		17. Pouched Animals (<i>Marsupialia</i>)	Pouched Rats (<i>Didelphides</i>)
IV. Cærolithic or Tertiary Epoch	10. Eocene Period 11. Miocene Period 12. Pliocene Period	18. Semi-Apes (<i>Prosimia</i>)	Lori (<i>Streps</i>) Maki (<i>Lemur</i>)
		19. Tailed Narrow- nosed Apes	Nose Apes Holy Apes
		20. Men-like Apes or Tail-less Narrow- nosed Apes.	Gorilla, Chimpan- zee, Orang, Gibbon
		21. Speechless Men or Ape-like Men	Cretins or Micro- cephali
V. Quaternary Epoch	13. Diluvial Period 14. Alluvial Period	22. Men capable of speech	Australians and Papuaans

The soft slimelike plasson-substance of the body of the Moneron is commonly called "*protoplasma*," and identified with the cell-substance of ordinary animal and plant cells. As, however, Eduard van Beneden, in his excellent work upon the Gregarinæ, first clearly pointed out, we must, strictly speaking, distinguish thoroughly between the plasson of cytods and the protoplasm of cells. This distinction is of special importance in its bearing on the history of evolution. As was before incidentally mentioned, we must assume two different stages of evolution in those elementary organisms, which, as formative particles or plastids, represent organic individuality of the first order. The older and lower stage is that of the cytods, in which the whole body consists of but one kind of albuminous substance, of the simplest plasson or formative material. The more recent and higher stage is that of cells, in which a separation or differentiation of the original plasson into two different kinds of albuminous substances, into the inner cell-kernel (*nucleus*), and the outer cell-substance (*protoplasma*), has already taken place.

The Monera are the simplest permanent cytods. Their entire body consists merely of soft, structureless plasson. However thoroughly we examine them with the help of the most delicate chemical reagents and the strongest optical instruments, we yet find that all the parts are completely homogeneous. These Monera are, therefore, in the strictest sense of the word, "organisms without organs;" or even, in a strictly philosophical sense, they might not even be called "organisms," since they possess no organs, since they are not composed of various particles. They can only be called organisms, in so far as they are capable of exercising the

organic phenomena of life, of nutrition, reproduction, sensation, and movement. If we tried to construct, *à priori*, the simplest conceivable organism, we should always be compelled to fall back upon such a Monera.

Although in all real Monera the body consists merely of such a small living piece of plasson, yet, among the Monera, which have been observed in the sea and in fresh water, we have been able to distinguish several different genera and species, varying in the mode in which their tiny bodies move and reproduce. In the ways in which movement is accomplished very noticeable differences exist.



FIG. 163.—A Moneron (*Protamœba*) in the act of reproduction: A, the whole Moneron, which moves, like the ordinary Amœba, by means of variable processes; B, a contraction round its circumference parts it into two halves; C, the two halves separate, and each now forms an independent individual (much enlarged).

In some Monera, especially in the *Protamœba* (Fig. 163), the formless body, during its movements, invariably develops only a few, short, and blunt processes, which project like fingers, slowly altering their form and size, but never branching. In other Monera, on the other hand (*e.g.* *Protomyxa*, *Myxastrum*), very numerous, long, fine, and generally thread-like processes arise from the surface of the movable body, and these branch irregularly, inter-

ing their free moving ends, so as to form a net. Huge
s of such slime-nets crawl upon the deepest bottom
e sea (*Bathybius*, Fig. 164). Within these soft slime-
plasson-nets slow currents continually pass. Such a
ron may be fed with finely pulverized colouring
r (for instance, carmine or indigo powder), if this
er is scattered in the drop of water under the micro-
in which the Moneron is contained. The grains
louring matter at first adhere to the surface of the
body, and then gradually penetrate, and are driven
in irregular directions. The separate smallest par-
or molecules, of the Moneron-body, called "plas-
es,"¹²⁶ displace each other, change their relative
ons, and thus effect a change in the position of the
bed particles of colouring matter. This change of
ion, at the same time, proves positively that a hidden
ate structure does not exist. It might be argued that
Monera are not really structureless, but that their
ization is so minute that, in consequence of the in-
uate power of our magnifying glasses, it is invisible.
objection is, however, invalid, for by the experiment
eding, we can, at any moment, prove the entrance of
gn, formed, small bodies into the different parts of the
of the Moneron, and that these are irregularly driven
t in all directions. At the same time we see that the
geable network of threads, formed by the branching
ie protoplasmic threads and the coalescence of the con-
t branches, alter their configuration every moment;
as has long been known to occur in the thread-nets
ie protoplasm in the interior of the plant-cells. The
era are, therefore, really homogeneous and structureless;

each part of the body is every other part. Each part can absorb and digest nourishment; each part is excitable and sensitive; each part can move itself independently; and lastly, each part is capable of reproduction and regeneration.

The reproduction of Monera always occurs asexually. In the *Protamoeba* (Fig. 163), each individual, after it has grown to a certain size, simply separates into two pieces. Round the circumference of the body a contraction arises, as in cell-division. The connection between the two halves continually becomes more slender (*B*), and finally parts in the middle. Thus, in the simplest possible way, two new individuals proceed by self-division from one quite simple individual (*C*). Other Monera, after they have grown to a certain size, gather themselves together into a spherical form. The globular protoplasmic body exudes a jelly-like protecting envelope, and a breaking-up of the whole plasmon-ball takes place within this covering; it breaks either into four pieces (*Vampyrella*), or into a large number of smaller globules (*Protomonas*, *Protomyxa*; cf. Plate I. in the "Natural History of Creation"). After a time, these globules begin to move, split the integument by their movement, and emerge; after which they float about by means of a long, thin, thread-shaped process. Each again passes by simple growth into the mature form. Thus, it is possible to distinguish different genera and species of Monera, on one hand, by the form of the different processes of the body, and, on the other hand, by the different kind and manner of reproduction. In the appendix to my monograph of the Monera I enumerated eight genera and sixteen species ("Biol. Studien," vol. i. p. 182). The

most remarkable of all Monera is the Bathybius, which was discovered by Huxley in 1868 (Fig. 164). This wonderful Moneron lives in the deepest parts of the sea, especially in



FIG. 164.—*Bathybius Hæckelii* (Huxley). A small piece of the formless continually changing plasson-net of this Moneron from the Atlantic Ocean.

Atlantic Ocean, and in places covers the whole floor of the sea in such masses, that the fine mud on the latter consists, in great measure, of living slime. The protoplasm of these formless nets does not seem differentiated at all; each little piece is capable of forming an individual. The active amœboid movements of these formless pieces of Moneron, which were first observed by the English zoologists Huxley and Wyville Thomson, have recently been again observed by the German Arctic voyager, Emil Bessels, in *Bathybius* of the coast of Greenland.¹⁸⁷

The origin and importance of these huge masses of organic, formless plasson-bodies in the lowest depths of the

sea, raises many different inquiries and thoughts. Spontaneous generation, especially, is naturally suggested by the Bathybius. We have already found that, for the origin of the first Monera upon our globe, the assumption of spontaneous generation is a necessary hypothesis. We shall be all the more inclined to confirm it now that, in the Monera, we have recognized those simplest organisms, the origin of which by spontaneous generation, in the present condition of our science, no longer involves very great difficulties. For the Monera actually stand on the very boundary between organic and inorganic natural bodies.¹⁸⁸

Next to the simple cytod-bodies of the Monera, as the second ancestral stage in the human pedigree (as in that of all other animals), comes the simple cell, that most undifferentiated cell-form, which, at the present time, still leads an independent solitary life, as the Amœba. For the first and oldest process of organic differentiation, which affected the homogeneous and structureless plasmon-body of the Monera caused the separation of the latter into two different substances; an inner firmer substance, the kernel, or *nucleus* and an outer, softer substance, the cell-substance, or *protoplasma*. By this extremely important separative process, by the differentiation of the plasmon into nucleus and protoplasm, the organized cell originated from the structureless cytod, the nucleolated from the non-nucleolated plasmon. That the cells which first appeared upon the earth originated in this manner, by the differentiation of the Monera, a conception which in the present condition of histologic knowledge seems quite allowable; for we can even now directly observe this oldest histological process of differentiation in Ontogeny. It will be remembered that in the

egg-cell of animals, either before or after fertilization, the original kernel disappeared. We explained this phenomenon as a reversion or atavism, and assumed that the egg-cell, in accordance with the law of latent heredity, first falls back into the kernel-less, cytod stage (Fig. 165). It is only after fertilization is accomplished that a new cell-kernel arises in this cytod, which thus becomes the parent-cell (*Cytula*, Fig. 166). The transitory kernel-less cytod-condition, intermediate between the egg-cell and the parent-cell, is an interesting germ-form, because, in accordance with the fundamental law of Biogeny, it reproduces the original, oldest parent-form of the Moneron; we therefore call it the Monerula. (Cf. vol. i. pp. 178-183.)




FIG. 165.—Monerula of Mammal (Rabbit). The fertilized egg-cell after the loss of the nucleus is a simple ball of protoplasm (*d*). The outer covering of the latter is formed by the modified *zona pellucida* (*z*) together with a mucous layer (*h*) secreted on to the outside of the latter. In this a few sperm-cells are still visible (*s*).

FIG. 166.—Parent-cell (*Cytula*) of a Mammal (Rabbit): *k*, parent-kernel; *n*, nucleolus of the latter; *p*, protoplasm of the parent-cell; *z*, modified *zona pellucida*; *s*, sperm-cells; *h*, outer albuminous covering.

We have already explained the one-celled germ-form, which we see in the original egg-cell and the parent-cell which is originated by the fertilization of the egg-cell, as the reproduction of a one-celled parent-form, to which we ascribed the organization of an *Amœba* (cf. Chap. VI.). For the *Amœba*, as it yet lives widely distributed in the fresh and salt waters of the globe, must be regarded as the most undifferentiated and most original of the various one-celled Primitive Animals. As the immature primitive egg-cells (which as "primitive eggs" or *Protova* are found in the ovary of animals) are indistinguishable from ordinary *Amœbæ*, we are justified in pointing to the *Amœba* as the one-celled phylogenetic form, which, in accordance with the fundamental law of Biogeny, is at the present time yet reproduced in the ontogenetic primitive condition of the "Amœboid egg-cell." As evidence of the striking correspondence of the two cells, it was incidentally mentioned that in the case of some Sponges the real eggs of these animals were formerly described as parasitic *Amœbæ*. Large one-celled *Amœba*-like organisms were seen creeping about in the interior of the Sponge, and were mistaken for parasites. It was only afterwards that it was discovered that these "parasitic *Amœbæ*" (Fig. 168) are really the eggs of the Sponge, from which the young Sponges develop. These egg-cells of the Sponge are, however, so like the true common *Amœbæ* (Fig. 167) in size and structure, in the nature of their nuclei and in the characteristic form of movement of their continually changing false-feet (*pseudopodia*), that, unless their source is known, it is impossible to distinguish them.

This phylogenetic explanation of the egg-cell and its

reference to the primæval ancestral form of the Amœba, directly enables us to give a definite answer to the old humorous riddle: Which was first, the egg or the hen? We can




FIG. 167.—A crawling Amœba (much enlarged). The whole organism has the form-value of a simple naked cell and moves about by means of changeable processes, which are extended from the protoplasmic body and again drawn in. In the inside is the bright-coloured, roundish cell-kernel or nucleus.

FIG. 168.—Egg-cell of a Chalk-Sponge (*Olynthus*). The egg-cell creeps about in the body of the Sponge by extending variable processes, like those of the ordinary Amœba.

now very simply answer this Sphinx-question, with which our opponents try to shake or even to refute the Theory of Evolution. The egg existed much earlier than the hen. Of course it did not exist in the form of a bird's egg, but as an undifferentiated amœboid cell of the simplest form. The egg existed independently during thousands of years as a simplest one-celled organism, as the Amœba. It was only after the descendants of these one-celled Primitive Animals had developed into many-celled animal forms, and after these had sexually differentiated, that the egg, in the present physiological sense of the word, originated from the amœ-

boid cell. Even then, the egg was first a *Gastræa*-egg, then a Worm-egg, then an *Acrania*-egg, then a Fish-egg, an Amphibian-egg, a Reptile-egg, and lastly, a Bird-egg. The egg of the Bird, as it now is, is a most complex historical product, the result of countless processes of heredity, which have occurred in the course of many millions of years.¹²⁹

The fact that this primitive egg-form, as it first appears in the ovary of the most dissimilar animals, is always of one form, an undifferentiated cell, of the simplest amœboid character, has already been pointed out as an especially important phenomenon. In this earliest young condition immediately after the individual egg-cell has originated, the consequence of a separation of the cells of the maternal ovary, no essential difference is recognizable in the egg-cells of the most dissimilar animals. (Cf. Fig. 10, vol. i. p. 134.) It is not till later, when the primitive egg-cells, or the primitive eggs (*protova*), have absorbed different kinds of nutritive yelk, and have surrounded themselves with variously formed coverings, and in other ways differentiated—it is not till they have in this way changed into after-eggs (*metova*), that those of different classes of animals can usually be distinguished. These peculiarities of the developed after-egg, the mature egg, are naturally to be considered as only secondarily acquired, by adaptation to the different conditions of existence both of the egg itself and of the animal which forms the egg.

The two first and oldest ancestral forms of the human race, which we have now considered, the *Moneron* and the *Amœba*, are, considered from a morphological point of view, simple organisms and individuals of the first order, *Plastids*. All subsequent stages in the ancestral chain are, on the

other hand, compound organisms or individuals of higher order—social aggregations of a number of cells. The earliest of these, which, under the name of *Synamœbæ*, we must rank as the third stage of our pedigree, are quite simple societies of all homogeneous undifferentiated cells; amœboid communities. To be certain as to their nature and origin, we need only trace the ontogenetic product of the parent-cell step by step. After the cytula (Fig. 166) has originated, by the re-formation of a cell-kernel, from



FIG. 169.—Original or primordial egg-cleavage. The parent-cell, or cytula, which resulted from the fertilization of the egg-cell, first breaks up, by a continuous and regular process of division, into two cells (*A*), then into four (*B*), then into eight (*C*), and, lastly, into very numerous cleavage-cells (*D*).

the Morula (Fig. 165), the parent-cell breaks up, by repeated division, into numerous cells. We have already minutely examined this important process of egg-cleavage, and have found that all the various modes of the latter are modifications of a single mode, that of original or primordial cleavage. (Cf. Chap. VIII., p. 188.) In the Vertebrate line this palingenetic form of egg-cleavage has been accurately re-




FIG. 170.—Mulberry-germ, or morula.

tained to the present time only by the *Amphioxus*, while all other Vertebrates have assumed a modified kenogenetic form of cleavage. (Cf. Table III., vol. i. p. 241.) The latter certainly originated at a later period than the former, and the egg-cleavage of the *Amphioxus* is, therefore, extremely interesting (vol. i. p. 442). In this the parent-cell first parts into two similar cells, the two first cleavage-cells (Fig. 169, *A*). From these, by continuous division, arise 4, 8, 16, 32, 64 cells, etc., etc. (Fig. 169). The final result of this primordial cleavage was, we found, the formation of a globular mass of cells, which was entirely composed of homogeneous, undifferentiated cells of the simplest character (Figs. 170, and 171, *E*). On account of the resemblance which this globular mass of cells bears to a mulberry or blackberry, we called it the "mulberry-germ," or *morula*.

This "morula" evidently at the present day shows us the many-celled animal body in the same entirely simple primitive condition in which, in the earlier Laurentian primitive epoch, it first originated from the one-celled amœboid primitive animal form. The morula reproduces, in accordance with the fundamental law of Biogeny, the ancestral form of the *Synamœba*. For the first cell-communities, which then formed, and which laid the first foundation of the higher many-celled animal body, must have consisted entirely of homogeneous and quite simple amœboid cells. The earliest Amœbæ lived isolated hermit lives, and the amœboid cells, which originated from the division of these one-celled organisms, must also have long lived isolated and self-dependent lives. Gradually, however, by the side of these one-celled Primitive Animals, small amœboid communities arose, owing to the fact that the

GERMINATION OF A CORAL.

57



Various kinds of Primitive Animals, which yet both in the sea and in fresh water, are formed essentially like the blastula, and which, in a certain sense, may be considered as permanent or persistent blastula-forms, have been discovered. These Planæads, or Blastæads, as they may be called, are formed in the very mixed societies of the Flagellatæ, especially the Volvoces (for instance, *Synura*). I noticed in September, 1869, on the Island Gili, on the coast of Norway, another very interesting form, which I named *Magosphæra planula* (Figs. 172, 173). The developed body of this forms a globular vesicle, the wall of which is composed of from thirty to forty vibratory homogeneous cells, and which swims about freely in the sea.



FIG. 172.—The Norwegian Flimmer-ball (*Magosphæra planula*), swimming by means of its vibratile fringes; seen from the surface.

FIG. 173.—The same, in section. The pear-shaped cells are seen brought together in the centre of the gelatinous sphere by a thread-like process. Each cell contains both a kernel and a contractile vesicle.

g reached maturity the society dissolves. Each separate cell still lives a while independently, grows, and changes into a crawling Amœba. This afterwards assumes a globular form, and encases itself by exuding a structureless integument. The cell now has just the appearance of a common animal egg. After it has remained for a time in a quiescent state, the cell breaks up, by means of consecutive division, first into 2, then into 4, 8, 16, 32 cells. These again arrange themselves so as to form a globular mass, put forth cilia, and bursting the encasing integument, swim about in the same Magosphæra-form from which we started. This accomplishes the entire life-history of a remarkable Primitive Animal.¹⁴¹

If we compare these permanent blastula-forms with the free-swimming Flimmer-larvæ or planula-condition, of similar structure, of many other lower animals, we may with certainty infer therefrom the former existence of a general and long-extinct parent-form, the structure of which was essentially like that of the planula or blastula. We will call this the Planæa, or Blastæa. The whole body, in the fully developed condition, consisted of a simple hollow sphere filled with fluid or structureless jelly, the wall of which formed a single stratum of homogeneous cells, lined with cilia. Many different kinds and species of Planæa-like Primitive Animals must certainly have existed, forming a distinct class of Protozoa, which we may call Planæa-swimmers (*Planæada*). A remarkable proof of the philosophical genius with which Karl Ernst Baer penetrated into the deepest secrets of the history of animal creation, is that, as early as the year 1828 (ten years before cell-theory was established), he guessed the significance

of the blastosphæra, and, truly prophetically, insinuates it in his classical "Entwicklungsgeschichte der Thiere" (vol. i. p. 223). The passage in question says: "The further back we go in evolution, the more do we find correspondence in very different animals. This leads to the question: Are not all animals in the beginning of evolution essentially alike, and is there not a principle common to all? As the germ is the undeveloped form itself, it is not without reason that it is asserted that the simple vesicular form is the common primitive form from which all animals, not only ideally, but also historically develop." This latter sentence has not only ontogenetic but also phylogenetic significance, and is all the more worthy because the blastula of the most diverse animals and the constitution of its wall of a single cell-stratum were not then known. And yet Baer, in spite of the insufficiency of his empiric grounds, ventured the statement: "At their first appearance all animals are alike, and are merely hollow globes."

Next to the primæval ancestral form of the type, the fifth stage in the human pedigree, is the Gastrula, which arises from the Planæa. Of all ancestral forms, as we have already shown, is of pre-eminent phylogenetic significance. Its former existence is certainly proved by the very important gastrula, which is met with as a germ-stage in the ontogeny of the most varied animals (Fig. 171, I, K). We found that the gastrula, in its palæogenetic form, is a globular, oval or oblong-rod-like form with one axis which has a simple cavity with one opening (at one pole of the axis). This is the primitive gut-cavity with its mouth-opening. The intestinal wall

of two cell-strata, which are, in fact, the two primary germ-layers, the animal skin-layer, and the vegetative intestinal layer.

The ontogenetic origin of the gastrula from the blastula at the present day affords us trustworthy intelligence as to the phylogenetic origin of the Gastræa from the Planæa. We found that on one side of the globular germ-membrane vesicle a groove-like depression begins, and this inversion (*invagination*) becomes continually deeper (Fig. 171, *H*). At last it is so great, that the outer, inverted part of the germ-membrane, or blastoderm, attaches itself closely to the inner, uninverted portion (Fig. 171, *I*). Now, if guided by this ontogenetic process, we wish to conceive the phylogenetic origin of the Gastræa in accordance with the fundamental law of Biogeny, we must imagine that the one-layered cell-society of the globular Planæa began, especially at one point of its surface, to absorb nourishment. At the nutritive point on the surface of the ball a groove-like depression was gradually formed by natural selection. The groove, which was at first quite shallow, in course of time became continually deeper. The function of nourishing, of absorption of nutriment, and digestion, was soon limited to the cells which lined the groove, while the other cells undertook the function of movement and covering. Thus originated the first division of labour among the originally homogeneous cells of the Planæa.

The first result of this earliest histological differentiation was the distinction of two different kinds of cells; within the hollow the nutritive cells, without, on the surface, the motive or locomotive cells. The distinction of the two primary germ-layers was thus caused. The inner cells

of the hollow formed the inner or vegetative layer, accomplishing the functions of nutrition; the outer cells of the covering formed the outer or animal layer, exercising the functions of locomotion and covering the body. This first and oldest process of differentiation is of such fundamental significance that it deserves the deepest thought. When we consider that the body of the human being with all its different parts, and also the body of all other higher animals, originates from these two simple primary germ-layers, we cannot over-estimate the phylogenetic significance of the gastrula. For in the quite simple primitive intestine, or the primitive intestinal cavity of the gastrula and its simple mouth-opening, the first real organ of the animal body, in a morphological sense, is gained; the earliest genuine organ, from which all the other organs have differentiated at a later period. The whole body of the gastrula is really only a "primitive intestine."

We have already pointed out the remarkable agreement between the palingenetic gastrula-forms of animals of the most diverse classes; of Sponges (Fig. 174, *A*), Polyps, Corals (Fig. 171, *I*), Medusæ, Worms (Fig. 175, *B*) Star-animals (*Echinodermata*, *C*), Articulated Animals (*Arthropoda*, *D*), Soft-bodied Animals (*Mollusca*, *E*), and Vertebrates (*F*). All these various forms of the palingenetic gastrula are much alike, and are only distinguished by such unessential and subordinate peculiarities, that the systematic zoologist, in his "natural system," could only represent them as different species of a single genus. The various kenogenetic gastrula-forms which have been described were also referable to that original palingenetic form (vol. i. p. 231). The gastrula proved to be a germ-form common to all classes of

animals, with the exception of the Protozoa. This highly important fact justifies the inference in accordance with the fundamental law of Biogeny, that the various ancestral




FIG. 175.

FIG. 176.

FIG. 177.

FIG. 178.




FIG. 174.



FIG. 179

FIG. 174.—(A) Gastrula of a Zoophyte (*Gastrophysema*), Haeckel.

FIG. 175.—(B) Gastrula of a Worm (Arrow-worm, *Sagitta*). After Kowalevsky.

FIG. 176.—(C) Gastrula of an Echinoderm (Star-fish, *Uraster*). After Alexander Agassiz.

FIG. 177.—(D) Gastrula of an Arthropod (Primitive Crab, *Nauplius*).

FIG. 178.—(E) Gastrula of a Mollusc (Pond-snail, *Limnaeus*). After Karl Rabl.


FIG. 179.—(F) Gastrula of a Vertebrate (Lancelet, *Amphioxus*) After Kowalevsky.

lines of all these classes of animals have developed phylogenetically from the same parent-form. This most significant primæval parent-form is the *Gastræa*.

The *Gastrea* was at any rate already present in the sea during the Laurentian period, and by means of its vibratory fringe hurried about in the water, just like the yet extant free-moving ciliated gastrulæ of this age. Probably the primæval *Gastræa*, which has been extinct for many millions of years, differed from the living gastrula of the present day only in some unessential point. On grounds derived from Comparative Anatomy and Ontogeny, the explanation of which would lead us too far, we may assume that the *Gastræa* had already acquired sexual reproduction, and did not only propagate its species asexually (by division—bud-formation or spore-formation), as was probably the case with the four preceding ancestral stages. Presumably, single cells of the primary germ-layers assumed the character of egg-cells, others that of fertilizing seed-cells. (Cf. Chapter XXV.) This hypothesis is founded on the fact that sexual reproduction is yet met with in the same simple forms in the lowest Plant-Animals (*Zoophyta*), especially in the Sponges.

Two small animal forms are especially interesting in their bearing on this aspect of the *Gastræa* theory. They have as yet been little observed, but of all extant animals they are most nearly allied to the primæval *Gastræa*, and may therefore be called "the *Gastræads* of the present day."¹⁴² One of these animals, *Haliphysema* (Figs. 180 and 181), has been described by Bowerbank as a Sponge; the other, *Gastrophysema*, by Carter as a Rhizopod (as "Squamulina"). The entire mature body of the developed person

of *Haliphysema* forms a most simple, cylindrical or egg-shaped pouch, the wall of which consists of two cell-strata. The cavity of the pouch is the stomach-cavity, and the



FIGS. 180, 181.—*Haliphysema primordiale*, an extant Gastræa-form. Fig. 180. External view of the whole spindle-shaped animal (attached by its foot to seaweed). Fig. 181. Longitudinal section of the same. The primitive intestine (d) opens at its upper end in the primitive mouth (m). Between the whip-cells (g) lie amœboid eggs (e). The skin-layer (h) below is encrusted with grains of sand, above with sponge-spicules.

opening at the top is the mouth-opening (Fig. 181, m). The two cell-strata forming the wall of the pouch are the

two primary germ-layers. These most simple Plant-Animals differ from the gastrula principally in the fact that the former are attached by one end (that opposite to the mouth opening) to the bottom of the sea, while the latter are free. Moreover, the cells of the skin-layer are coalescent and have included many foreign bodies, such as sponge-spicules and sand-grains, etc., which serve to support the body-wall (Fig. 180). The intestinal layer, on the other hand, consists merely of a stratum of ciliated cells (Fig. 181). When the *Haliphysema* is sexually mature, individual cells of its entoderm assume the character of female egg-cells; on the other hand, individual cells of its exoderm become male seed-cells; the fertilization of the former by the latter



FIGS. 182, 183.—Ascula of a Sponge (*Olynthus*). Fig. 182, from the outside; Fig. 183, in longitudinal section: *g*, primitive intestine; *o*, primitive mouth; *i*, intestinal layer; *e*, skin-layer.

takes place directly through the stomach-cavity. A true palingenetic gastrula (Fig. 174) develops, just as in the *Monoxenia* (Fig. 171), from the fertilized egg. This swims about for a time in the sea, then attaches itself, and in this state resembles a simple young-form, which occurs in the course of the evolution of many other Plant-Animals, and which is called the ascula (Figs. 182, 183). In consequence of the absorption of foreign bodies by the exoderm, it becomes the *Haliphysema*.

When we consider that there is no other important difference between the free-swimming gastrula and this attached, simplest Plant-animal, we are fairly justified in stating that in the simplest form of *Gastræa* sexual reproduction must have taken place in the same way. In the *Gastræads*, just as in Plant-animals, both kinds of sexual cells—egg-cells and sperm-cells—must have formed in the same person; the oldest *Gastræads* must, therefore, have been hermaphrodite. For Comparative Anatomy shows that hermaphroditism, that is, the union of both kinds of sexual cells in one individual, is the oldest and original condition of sexual differentiation; the separation of the sexes (*Gonochorismus*) did not originate till a later period.

TABLE XVII.

Systematic Survey of the five earliest evolutionary stages of the Human Ancestral Line, compared with the five earliest stages of Individual and of Systematic Evolution.

Form-Value Of the five earliest stages of the animal body.	Phylogeny. The five earliest stages in the evolution of the tribe.	Ontogeny. The five earliest stages in the evolution of the germ.	The System. The five earliest stages in the animal system.
1. <i>First Stage.</i> A quite simple cytod (a non-nucleated plastid). 2. <i>Second Stage.</i> A simple cell (a nucleated plastid). 3. <i>Third Stage.</i> A quite simple aggregation of simple, similar cells. 4. <i>Fourth Stage.</i> A simple hollow globe, filled with liquid, the wall of which consists of a single stratum of homogeneous cells. 5. <i>Fifth Stage.</i> A hollow body, with a single axis, the wall of which consists of different cell-strata; with an opening at one pole of the axis.	1. Monera. The oldest animal Monera (originating by spontaneous generation). 2. Amœba. Oldest animal Amœba. 3. Synamœba. The oldest aggregation of animal Amœbæ. 4. Planœa. An animal hollow globe, the wall of which consists of a single stratum of ciliated cells. (<i>blastœa.</i>) 5. Gastrœa. Parent-form of intestinal animals, or <i>Metazœa</i>. Simple primitive intestine with primitive mouth. The body-wall is formed by the exoderm and the entoderm.	1. Monerula. A non-nucleated animal-egg (after fertilization and after loss of the germ-vesicle). 2. Cytula. A nucleated, fertilized animal-egg ("first cleavage globule"). 3. Morula. "Mulberry-germ." A globular mass of cleavage-cells. 4. Blastula. A hollow globe, the wall of which consists of a single stratum of homogeneous cells (the Planula of lower animals). (<i>blastophœra.</i>) 5. Gastrula. Intestinal larva. A simple intestinal cavity with a mouth-opening. The body-wall is formed by the two primary germ-layers.	1. Monera. Protamœba, Bathybius, and other extant Monera. 2. Amœba. Extant Amœba. 3. Labyrinthula. A mass of similar, one-celled primitive animals. 4. Magosphœra. A hollow globe, the wall of which consists of a single stratum of homogeneous ciliated cells. 5. Haliphysœma. A quite simple plant-animal. An unarticulated uniaxial person, the body-wall of which consists of the exoderm and the entoderm.

CHAPTER XVII

THE ANCESTRAL SERIES OF MAN.

II.—FROM THE PRIMITIVE WORM TO THE SKULLED ANIMAL.

The Four Higher Animal Tribes are descended from the Worm Tribe.—The Descendants of the *Gastræa*; in one direction the Parent Form of Plant-Animals (Sponges and Sea-Nettles), in the other the Parent Form of Worms.—Radiate form of the former, Bilateral form of the latter.—The Two Main Divisions of the Worms, *Acœlomi* and *Cœlomati*: the former without, the latter with, a Body Cavity and Blood-vessel System.—Sixth Ancestral Stage: *Archelminthes*, most nearly allied to *Turbellaria*.—Descent of the *Cœlomati* from the *Acœlomi*.—Mantled Animals (*Tunicata*) and Chorda-Animals (*Chordonia*).—Seventh Stage: Soft-Worms (*Scolecida*).—A Side Branch of the latter: the Acorn-Worm (*Balanoglossus*).—Differentiation of the Intestinal Tube into Gill-intestine and Stomach-intestine.—Eighth Stage: Chorda-Animals (*Chordonia*).—Ascidian Larva exhibits the Outline of a Chorda-Animal.—Construction of the Notochord.—Mantled Animals and Vertebrates as Diverging Branches of Chorda-Animals.—Separation of Vertebrates from the other Higher Animal Tribes (Articulated Animals, Star-Animals, Soft-bodied Animals).—Significance of the Metameric Formation.—Skull-less Animals (*Acrania*) and Skulled Animals (*Craniota*).—Ninth Ancestral Stage: Skull-less Animals.—*Amphioxus* and Primitive Vertebrate.—Development of Skulled Animals (Construction of the Head, Skull, and Brain).—Tenth Ancestral Stage: Skulled Animals, allied to the *Cyclostomi* (*Myxinoïdæ* and *Petromyzonidæ*).

“Not like the gods am I! Full well I know;
But like the worm which in the dust must go,
And, finding in the dust his life and weal,
Is crushed and buried by the traveller’s heel.—

Why dost thou grin at me, thou hollow skull?
As though of old thy brain, like mine, was vexed,
Had looked to find bright day, but in the twilight dull,
In search for truth, was sad and sore perplexed!"

GOETHE.

BOTH in prose and in poetry man is very often compared to a worm. "A miserable worm," "a poor worm," are common and almost compassionate phrases. If we cannot detect any deep phylogenetic reference in this zoological metaphor, we might at least safely assert that it contains an unconscious comparison with a low condition of animal development which is interesting in its bearing on the pedigree of the human race. For there is no doubt that the vertebrate tribe, in common with those of the other higher classes of animals, have developed phylogenetically from that multiform group of lower invertebrate animals which are now called Worms. However closely we limit the zoological significance of the word "Worm," it yet remains indubitable that a large number of extinct Worms must be reckoned among the direct ancestors of the human race.

The group of Worms (*Vermes*) is much more limited in the Zoology of the present day, than was the same class in the older Zoology, which followed the system of Linnæus. It, however, yet includes a great number of very diverse lower animals, which, phylogenetically, we may regard as the few last living twigs of an immense spreading tree, the trunk and main branches of which have for the most part long since died off. On the one side, among the widely divergent classes of Worms, are found the parent-forms of the four higher tribes of animals, the Molluscs, Star-animals, Articulates, and Vertebrates; on the other side,

eral comprehensive groups and also single isolated genera. Worms are to be regarded as root-suckers which have branched directly from the rest of the primæval family-tree of the Worms. Some of these suckers have evidently branched but little from the long-extinct parent-form, the Primitive Worm (*Prothelms*), which is immediately connected with the Gastræa.

Comparative Anatomy and Ontogeny clearly and significantly prove that the Gastræa must be regarded as the direct ancestor of this Primitive Worm. Even now, a Gastrula develops from the egg of all Worms after its hatching. The lowest and most imperfect Worms retain throughout life an organization so simple that they are but little raised above the lowest Plant-animals, which are also the immediate descendants of the Gastræa, and which also yet develop directly from the gastrula. If the genealogical relation of these two lower animal tribes, the Worms and the Plant-animals, is closely examined, it becomes evident that the most probable hypothesis of their descent is, that they have both originated, as independent branches, directly from the Gastræa. On the one side, the common parent-form of the Worms developed from the Gastræa; as, on the other side, did the common parent-form of the Plant-animals. (Tables XVIII. and XIX.)

The tribe of Plant-animals (*Zoophytes*, or *Cœlenterata*) now comprehends, on the one side, the main class of Sponges (*Spongiæ*); on the other, the main class of the Sea-nettles (*Calephæ*); to the former belong the Gastræads and Siphoniferæ, to the latter the Hydroid-polyps, the Medusæ, the Anthrophoræ, and Corals. From the Comparative Anatomy and the Ontogeny of these we may infer, with great pro-

bability, that all these Plant-animals descend from a common and very simple parent-form, the structure of which resembled that of the ascula in essential points (Figs. 182, 183, p. 68). The uniaxial outline of the ascula and the gastrula is usually retained by the Sponges, while in most Sea-nettles (*Acalephæ*) transverse axes have been differentiated in the course of further evolution, thus giving rise to a characteristic radiate structure with a pyramidal general outline.

In distinction from this predominant radiate outline of Plant-animals, a marked bilateral general outline is developed from the first in the second offshoot from the gastrula, in the Worms. As the radiate form is marked by adaptation to an adherent mode of life, so is the bilateral form by adaptation to certain definite acts of free locomotion. The constant direction and carriage of the body which would be maintained in this mode of free locomotion, conditioned the two-sided, or bilateral outline of the symmetrical Worms. Even the parent-form of the latter, the Primitive Worm (*Prothelms*) must have acquired this character, and thus have become distinguished from the uniaxial parent-form of the Plant-animals. In this simple mechanical impetus, in the defined free locomotion of the Worms, on the one hand, and in the stationary mode of life of the earliest Plant-animals on the other, we must look for the efficient cause which produced in the one the bilateral or two-sided, in the other the radiate outline of the body. The former, the bilateral outline, has been inherited by the human race from the Worms.

Except through the Gastræa, the common parent-form of Plant-animals and Worms, the human race is, therefore,

not related to the Plant-animals. It will be our next task to consider more closely the pedigree of Man in so far as it coincides with that of the Worms. Let us examine how far the Comparative Anatomy and Ontogeny of Worms justify us in looking among the latter for primæval ancestors of Vertebrates, and therefore of Man. For this end we must first consider the zoological system of Worms. In accordance with the most recent investigations of the Comparative Anatomy and Ontogeny of Worms, we divide (without reference to the many and various peculiarities of the numerous separate classes, which in this place do not interest us) the whole mass of forms within this tribe into two large main groups. The first main group, which we call Bloodless Worms (*Acoelomi*), comprehends the earlier division of the lower Worms, which have no true body-cavity, no system of blood-vessels, no heart, no blood, —in short, none of the parts connected with this organ-system. The second main group, on the contrary, called Blood-worms (*Cœlomati*), are distinguished from the former by the possession of a true body-cavity, and also by the presence of a blood-like fluid, which fills this cavity; most of them also develop special blood-vessels, which again cause further correlated advances in structure. The relation of these two main groups of Worms is very evidently phylogenetic. The *Acoelomi*, which are very nearly allied to the *Gastrea* and the Plant-animals, are to be regarded as an earlier and lower group, from which the more recent and higher division of the *Cœlomati* developed, perhaps towards the end of the Laurentian Period.

We will first carefully examine the lower group of Worms, the *Acoelomi*, among which we must look for the

sixth ancestral stage of the human race, the stage immediately following the gastrula. The name "Acoelom" signifies "Worms without a body-cavity, or coeloma," therefore without blood, or vascular system. The extant Acoelomi are generally included in a single class, which, account of their flattened bodies, are called Flat-worms (*Plathelminthes*). To this class belong the Gliding-worms (*Turbellaria*), which live independently in the water; the parasitic intestinal Sucking-worms (*Trematoda*), and the Tape-worms (*Cestoda*), which have become yet more degraded by parasitism. The phylogenetic relations of these three forms of Flat-worms are very evident; the Sucking-worms originated from the free Gliding-worms by adaptation to a parasitic mode of life; and, by a yet more complete parasitic life, the Tape-worms originated from the Sucking-worms. These are striking examples of the gradual increasing degeneration of the most important organs.

In addition to these well-known extant Flat-worms, great numbers of other Acoelomi must have lived during the Archilithic Epoch, which in general form were very much like those of the present day, but were, in several respects, yet more simply organized, and were, in the lowest stages of development, immediately connected with the Gastræans. The whole of these lowest Acoelomi, among which the common parent-form of the whole Worm tribe (the *Prothelminis*) must have been, may be classed as "Primitive Worms" (*Archelminthes*).

The two classes of the Acoelomi, the Primitive Worms and the Flat-worms, represent in their external form the simplest bilateral condition of the animal body. The body is a simple oval, usually somewhat flattened, with

out any appendage (Figs. 184, 185). The dorsal side of the leaf-like body differs from the ventral side, on which the Worm creeps. Accordingly, even in these most simple Worms there are the three definite axes which mark the bilateral type-form, and which re-occur in the human body and in that of all higher animals: (1) a longitudinal axis (main axis), which passes from front to rear; (2) a lateral axis, passing from right to left; and (3) a sagittal axis, passing from the dorsal to the ventral surface. (Cf. vol. i. p. 257.) This so-called symmetrical or "bilateral" arrangement of the outline of the body is simply the mechanical result of adaptation to a creeping form of locomotion, during which one end of the body is always directed forwards. The geometric outline of the gastrula, as of the ascula, has but one axis with unequal poles (*Monaxonia diplopola*). The typical outline of Worms, as of Vertebrates, is, on the contrary, bilateral, with tranverse axes (*Stauraxonia dippleura*).¹⁴⁴

The whole outer surface of the Gliding-worms (*Turbellaria*) is covered, as in the gastrula, with a thick, fine ciliated coat; that is, with a fur-like covering of extremely fine and close microscopic hairs, which are direct processes of the uppermost cells of the epidermis, and maintain an uninterrupted whirling or vibratory motion (Fig. 184, *f*). The constant vibrations of these cilia cause a continued current of water over the surface of the body. Fresh water is constantly conveyed to the surface of the skin by this current, thus permitting respiration in its simplest form (skin-respiration). A similar ciliated covering, just as is seen in the extant Gliding-worms of our fresh-water seas, presumably covered our extinct ancestors of the Primitive

Worm group, the *Archelminthes*. They inherited the ciliated dress directly from the *Gastræa*.

If we now make various vertical sections (longitudinal and transverse) through the simple body of the Gliding-worm (and that of the *Archelminthes* which are certainly very closely allied to the former), we soon discover that the internal structure is considerably higher than that of the *Gastræads*. We first observe that the two primary germ-layers (inherited from the *Gastræa*) have differentiated into several cell-strata. The skin-layer and the intestinal layer have each split into two strata. The four secondary germ-layers, which are thus produced, are the same that we found resulted from the first differentiation of the two primary germ-layers in the embryo of the Vertebrate also. (Cf. the transverse sections through the larval *Amphioxus* and Earth-worm, Figs. 50 and 51, p. 236, and Plate IV. Fig. 2 and Plate V. Fig. 10.)

The highly important histological differentiation of these four secondary germ-layers led directly to further organological processes of differentiation, by which the organism of the Primitive Worms was soon considerably raised above that of the *Gastræads*. In the latter there was really, in a morphological sense, but a single organ, the primitive intestine, with its mouth-opening. The whole body was nothing but an intestinal canal; the intestinal wall was at the same time the wall of the body. Of the two cell-layers forming this intestinal wall, the inner accomplished the functions of nutrition, the outer those of motion and covering. As some of the cells of the primary germ-layers developed into egg-cells, and others into sperm-cells, these layers also performed the function of reproduction. In the

Primitive Worms, however, simultaneously with the formation of the secondary germ-layers, these various functions also began to be distributed to various organs, which detached themselves from the original main organ, the primitive intestine. Special organs originated for reproduction (sexual glands), for secretion (kidneys), for motion (muscles), and for sensation (nerves and sense-organs).

In order to obtain an approximate picture of the simplest form in which all these various organs first appeared in the Primitive Worms, it is only necessary to examine the most imperfect forms of Gliding-worms (*Turbellaria*), as they exist at the present time in salt and fresh water. They are mostly very small and insignificant Worms of the simplest form, many being scarcely a millimetre or a few millimetres in length. In the simplest species of Gliding-worms the greater part of the oval body is occupied by the intestinal canal. This is a very regularly shaped pouch with an opening, representing both mouth and anus (Fig. 184, *m*). At the anterior section of the intestinal tube, which is separated as a throat (*pharynx*, *sd*), the fibrous layer is very thick, a thick muscular layer. Immediately outside the intestinal-fibrous layer lies the skin-fibrous layer, which in most worms appears as a large skin-muscle sac. Above the throat in Gliding-worms a nerve system of the simplest form is already visible in front, a pair of small nerve-knots, or ganglia, which from their position are called the "upper throat ganglia," or "brain" (Fig. 185, *g*). Delicate nerve-threads (*n*) pass from this to the muscles and to the ciliated skin-sensory layer. A pair of quite simple eyes (*au*) and nose-pits (*na*) are to be found in a few Gliding-worms. The Flat-worms are also universally provided with

a pair of simple kidney-canals ("excretory organs"), the form of two long, thin, glandular tubes, which traverse the right and left sides of the intestine and open at the hinder end of the body (Fig. 184, *nm*). We found that





FIG. 184.—A simple Gliding-worm (*Rhabdocalum*): *m*, mouth; *sd*, the epithelium; *sm*, throat-muscles; *d*, stomach-intestine; *nc*, kidney due *nm*, opening of the kidneys; *au*, eye; *na*, nose-pit.

FIG. 185.—The same Gliding-worm, showing the remaining organs brain; *au*, eye; *na*, nose-pit; *n*, nerves; *h*, testes; ♂, male opening; ♀, female opening; *e*, ovary; *f*, ciliated outer-skin.

primitive kidney canals in the vertebrate embryo are formed at a very early period, shortly after the first formation of the middle germ-layer (*mesoderma*). The presence of these at so early a period shows that they are very important primordial organs. It also shows their universal existence in all Flat-worms; for even Flat-worms, which, in consequence of the adoption of a parasitic mode of life, have lost the intestine, yet have the rudimentary primitive kidneys, or "excretory ducts." The existence of these, therefore, is to be older and of greater physiological importance than the blood-vessel system, which is wholly wanting in the Flat-worms. The sexual organs appear in the Gliding-worms in a very complex form; in others their form is very simple. Most of them are hermaphrodites; that is, each individual worm has both male and female sexual organs. In the simplest we find a testis in the anterior part (Fig. 185, *h*), and a single or double ovary behind (*a*). One of these simplest forms, the *Acœlomi*, such as we find among the lowest Rhabdionchæ, may give us an approximate idea of the structure of the Primitive Worm, which forms the sixth stage in the human pedigree.

These ancestors of the human race, which, on account of their general organization, must be placed among the lowest Worms (*Acœlomi*), must have been represented in the Archilithic Epoch by a large number of various forms. The lowest of these must have been directly connected with the Gastræans (the fifth ancestral stage); the highly developed must, on the other hand, have been connected with the Cœlomati (the seventh stage). However, our present knowledge of the Comparative

Anatomy and Ontogeny of the Accelomi is very fragmentary, and much too imperfect to enable us to point with certainty to the series of the various stages, we will attempt a detailed arrangement of them. We will take instead to the seventh stage in the human pedigree, which belonged to the multiform group of the Blood-bearing Worms (*Cœlomati*).

The great organic advance in structure by which the Blood-bearing Worms, or Cœlomati, developed from the older Bloodless Worms, or Accelomi, consisted in the formation of a body-cavity (*cœloma*), and of a nutritive juice filling the latter, the first blood. All the lower animals with which we have yet occupied ourselves in our Phylogeny, all the Primitive Animals and Plant-animals, as like the Accelomi, bloodless and without a body-cavity. The formation of a special vascular system, the earliest Cœlomati made a very great advance. Much of the complexity in the organic structure in the four higher tribes of animals is based on the differentiation of the vascular system, which they have inherited from the Blood-bearing Worms.

The first development of a true body-cavity (*cœloma*) is referable to the separation of the two fibrous layers; the formation of a spacious cavity between the outer skin-fibrous layer and the inner intestinal-fibrous layer. In the fissure-like gaps, which formed between the two germ-layers a juice collected, which penetrated through the intestinal wall. This juice was the first blood, and the gaps between the two germ-layers formed the first rudiment of the body-cavity. The union of these gaps formed the simple cœloma, the large cavity, containing blood or lymph, which plays

important a part in all the higher animals as the receptacle of the very extensive intestines. The formation of this coelom, and of the blood-vessels developed in connection with it, exercised a very great influence on the further evolution of the animal organization. The most important result was, that it allowed the conveyance of rich nutritive juices to those parts of the body lying near the circumference, and developing at a considerable distance from the intestinal canal. The intimate correlation, or reciprocity of the parts, necessarily occasioned, in direct connection with the progressive development of the blood-vessel system, many other important advances in the structure of the body of the Blood-bearing Worms.

Just as among the Acoelomi, so also among the Coelomati, the pedigree of our race must have passed through a large number of diverse ancestral stages. But among extant Coelomati (which form but a very small fraction of this once multifarious group), there are but very few Worms which can with certainty be regarded as nearly allied to the long-extinct ancestors of Man. In this respect, but a single class of Coelomati is really of prominent importance; these are the Mantled Animals (*Tunicata*), to which belong the Ascidia already known to us. Our careful examination of the structure and germ-history of the Ascidian and the Amphioxus have shown the extreme importance of these very interesting animal forms. (Cf. Chapters XIII. and XIV.) That examination fully justifies us in asserting that among the ancestors of the Vertebrates (and therefore of Man) there was an unknown extinct coelomate species, to which the nearest allied form among extant animals is the Appendicularia (Fig. 187), of which we have already

spoken, and the tailed Ascidian larva. We will for the present call this kind of Worm, which was primarily distinguished by the possession of a notochord, the Chorda-animal (*Chordonium*). The Ascidians on the one hand, and the Vertebrates on the other, developed, as two diverging branches, from these Chorda-animals. The common parent form of the Chorda-animals themselves was a coelomate form which finally must have descended from the Accelomi, and from the Archelminthes.

Many connecting intermediate forms must, of course, have existed between these two groups of Worms, between the Primitive Worms and the Chorda-animals. Unfortunately however, zoological knowledge is at present especially imperfect with regard to these important intermediate forms of the multiform Worm tribe. For very evident reasons none of these Worms could leave fossil remains. For, like the great majority of other Worms, they had no hard parts in their bodies. Most even of the known fossil Worms are worthless, for they tell us little or nothing of the most important structural features of the soft body. Fortunately however, we can in great measure satisfactorily fill the considerable palæontological gap in this part of our pedigree with the help of the Comparative Anatomy and Ontogeny of Worms. If, on the one hand, we examine the structure and mode of development of the lower Worms from the Gliding Worms (*Turbellaria*), and, on the other hand, the Anatomy and Ontogeny of the Ascidians, it is not difficult, step by step, to re-construct in imagination the connecting intermediate forms, and to insert a series of extinct ancestor forms between the Accelomi and the Chordonia. This series of forms under the name of Soft-worms (*Scolec*

we will consider as the seventh stage in the human pedigree.

An examination of the Comparative Anatomy of the various Scolecoid forms, which we might perhaps distinguish here, would lead us much too far into the difficult details of the Comparative Anatomy and Ontogeny of the Worms. For our purpose it seems more important to call attention to those phylogenetic advances, by means of which the organization of the earliest Blood-bearing Worms was in the end elevated to that of the Chorda-animals. The Comparative Anatomy and Ontogeny of the Gliding-worms and of the Ascidians justify us in giving special weight to the significant differentiation of the intestinal canal into two distinct divisions; into an anterior division (the gill-intestine), which accomplishes respiration, and a posterior division (the stomach-intestine), which accomplishes digestion. As in *Gastræads* and Primitive Worms, so also in the Ascidian larva, the intestinal canal is at first a simple pouch-like body, provided merely with a mouth-opening. A second opening, the anus, does not develop till a later period. Gill-openings afterwards appear in the anterior section of the intestinal canal, by which the whole anterior intestine is transformed into a gill-body. This remarkable arrangement is, as we found, quite peculiar to Vertebrates, and, except in the Ascidians, occurs nowhere else. Among extant Worms there is, however, a single isolated and very remarkable Worm form, which in this respect may be regarded as distantly allied to the Ascidia and to Vertebrates, and perhaps as an off-shoot from the Soft-worms (*Scolecida*). This is the so-called "Acorn-worm" (*Balanoglossus*, Fig. 186), which lives in the sand of the sea-shore. The in-

interesting points connecting this with Ascidians and

Skull-less Animals (*Acra* were first accurately observed and explained by Gegenbaur. though this singular *Balanoglossus* is in many other respects peculiar in its organization, so that Gegenbaur rightly ranked it as the representative of a special class (*Enteropneusta*), yet the structure of the anterior section of the testinal tube is exactly similar to that of Ascidians and Skull-less Animals (*k*), a gill body, the wall of which are pierced on either side by gill-openings and are supported by gill-arches. Now, although the *Acorn-worm* in other points of structure may differ very considerably from those extinct *Acorn-worms* (*Scolecidae*), which we regard as direct ancestors of the *Acorn-worm* race, and as intermediate links between the Primitive *Worms*



FIG. 186.—A young *Acorn-worm* (*Balanoglossus anoglossus*). (After Alexander Agassiz.) *r*, acorn-like proboscis; *h*, collar; *k*, gill openings and gill-arches of the anterior testine, in a long row one behind another on each side; *d*, digestive posterior intestine, filling the greater part of the body cavity; *v*, intestinal vessel, lying between two parallel folds of skin; *a*, anus.

and the Chorda-animals, yet, in virtue of this characteristic structure of the gill-intestine, it may be considered a remotely allied collateral line of the Soft-worms. The development of an anus (Fig. 186, *a*) at the end opposite to the mouth, is also a considerable advance in the structure of the intestine. The further development of the blood-vessel system in the Acorn-worm also indicates a marked advance. In the ciliary surface of the skin, on the contrary, it recalls the Gliding-worms. The sexes are separated, while our scolecid ancestors were probably hermaphrodite.¹⁴⁵

From a branch of the Soft-worms, the group of Chorda-animals (*Chordonia*), the common parent-group of the Mantle-animals and Vertebrates also developed. The process which primarily led to the development of this important group of the *cœlomati*, was the formation of the inner axial skeleton (the notochord, or *chorda dorsalis*), which at the present day we find permanently retained in its simplest form in the lowest Vertebrate, the *Amphioxus*. We saw that this notochord is already found in the tailed and free-swimming larva of the Ascidian (Plate X. Fig. 5). The chorda does, indeed, serve specially as a support for the rudder-like tail of the larval Ascidian, but its anterior extremity passes in between the intestinal and medullary tubes within the actual body of the larva. A transverse section of this larva therefore shows that arrangement of the most important organs which is characteristic of the vertebrate type: in the centre is the firm notochord, which supports the other organs and serves especially as a base and point of attachment for the motive trunk muscles; above this notochord, on the dorsal side, is the central

nervous system in the form of a medullary tube; below, on the ventral side, is the intestinal tube, the anterior half of which is a respiratory gill-intestine, its posterior half a digestive stomach-intestine. It is true that the free-swimming larva of the extant Ascidian possesses this typical vertebrate character only for a short time; it soon relinquishes its free roving mode of life, puts off its oar-like tail with the notochord, adheres to the bottom of the sea, and then undergoes that very great retrogression, the surprising final result of which we have already observed (Chapters XIII. and XIV.). Nevertheless, the Ascidian larva, in its very transitory evolution (for a brief space), affords us a picture of the long extinct Chordona-form, which must be regarded as the common parent-form of Mantle-animals and Vertebrates. There is even yet extant a small and insignificant form of Mantle-animal which throughout life retains the structure of the Ascidian larva with its oar-like tail and its free-swimming mode of life, and which reproduces itself in this form. This is the remarkable Appendicularia (Fig. 187), which we have already examined.

If we ask ourselves what conditions of adaptation could possibly have had so remarkable a result as the development of the notochord, and the modification of a branch of the Soft-worms into the parent-form of the Chorda-animals, we may with great probability answer, that this result was effected by the habituation of the creeping Soft-worm to a swimming mode of life. By energetic and continued swimming movements, the muscles of the trunk would be greatly developed, and a strong internal point of attachment would be very favourable to this muscular

activity. A support of this kind might arise by enlargement and concrescence of the germ-layers along the longitudinal axis of the body; and the differentiation of an independent bony cord from this axial cord gave rise to the notochord. (Cf. Fig. 88, 89, vol. i. pp. 300, 301.) In correlation to the formation of this central notochord, the simple nerve-ganglia, lying over the throat in the Soft-worms, lengthened into a long nerve-cord, reaching from front to rear, above the notochord; in this way, the medullary tube originated from the "upper throat ganglia."

As we have already minutely considered the great significance of the Ascidians (Fig. 188) in this respect, as well as their close relations to the *Amphioxus* (Fig. 189), we will not tarry longer over this point now. I will repeat, that we must by no means regard the Ascidian as the direct parent-form of the *Amphioxus* and of the other Vertebrates. On the contrary, we assert that, on the one hand the Ascidians, and on the other the Vertebrates, have both descended from one unknown Worm form, which has long been extinct; the nearest relatives of this among existing animals are the Ascidian larvæ and the Appendicularia (Fig. 187). This unknown common parent-form must have belonged to the group of Chorda-animals, which we pointed out as the eighth ancestral stage in the human pedigree.¹⁴⁶ Although we cannot form an entirely satisfactory idea as to all points of external and internal structure of this Chorda-animal, there is no doubt that, like its near relatives, the Mantle-animals, and like the preceding ancestral stage represented by the Soft-worms and Primitive Worms, it must be classified in the natural system of the animal

kingdom as a genuine Worm. The difference between and other genuine Worms cannot have been greater than





FIG. 187.—Appendicularia, seen from the left side: *m*, mouth; *k*, *g* intestine; *o*, oesophagus; *v*, stomach; *a*, anus; *n*, nerve-ganglia (up throat-knots); *g*, ear-vesicle; *f*, ciliated groove under the gill; *h*, heart; *t*, testes; *e*, ovary; *c*, notochord; *s*, tail.

FIG. 188.—Structure of an Ascidian (seen from the left, as in Fig. 1 and Fig. 14, Plate XI.): *sb*, gill-sac; *v*, stomach; *t*, large intestine; heart; *t*, testes; *vd*, seed-duct; *o*, ovary; *o'*, matured eggs in the body cavity. (After Milne Edwards.)

the difference between the extant Tape-worms and Ringed Worms (*Annelida*). Moreover, in a certain sense we may regard the extant Appendicularia as a last remnant of the Chordonia class.

We have now studied the most important animal forms which occur in the pedigree of the human race, and which, in the zoological system, must be classed among the Worms. In leaving this lower class, and tracing our ancestry henceforth exclusively within the vertebrate tribe, we at once leave behind the great majority of animal forms, which branched off from the worm tribe in entirely different directions. When, in a previous chapter (IX.), the vertebrate nature of man was proved, it was incidentally mentioned that the very great majority of animals are in no way directly allied to our tribe. The parent-forms of the three other higher Animal tribes, the Articulated Animals (*Arthropoda*), Star-animals (*Echinoderma*), and Soft-bodied Animals (*Mollusca*), like the vertebrate tribe, originated from the




FIG. 189.—Lancelet (*Amphioxus lanceolatus*), twice the actual size, seen from the left (the longitudinal axis is represented vertically, the mouth turned upward, the tail downward, as in Plate XI. Fig. 15): a, mouth-opening, surrounded by cilia; b, anal opening; c, ventral opening (*Forus abdominalis*); d, gill-lesds; e, stomach; f, liver-cæcum; g, large intestine; h, coelom; i, notochord (under it the aorta); k, arches of the aorta; l, main gill-artery; m, swellings on its branches; n, hollow vein; o, intestinal vein.


TABLE XVIII.

Systematic Survey of the Phylogenetic System of the Animal Kingdom
founded on the Gastræa Theory and the Homology of the Germ-layers.

<i>Tribes or Phyla of the Animal Kingdom.</i>	<i>Main Classes or Branches of the Animal Kingdom.</i>	<i>Classes of the Animal Kingdom.</i>	<i>Systematic Names of the Classes.</i>
FIRST SUB-KINGDOM: PRIMITIVE ANIMALS (<i>Protozoa</i>).			
Animals without germ-layers, intestine, or true tissues.			
A. Primitive Animals Protozoa	I. Egg-animals <i>Ovularia</i> II. Infusorial animals <i>Infusoria</i>	1. Monera 2. Amœbas 3. Gregarines 4. Sucking Infusoria 5. Ciliated Infusoria	1. Monera 2. Lobosa 3. Gregarines 4. Acinetes 5. Ciliata
SECOND SUB-KINGDOM: INTESTINAL ANIMALS (<i>Metazoa</i>).			
Animals with two primary germ-layers, intestines and tissues.			
B. Plant- Animals Zoophytes	III. Sponges <i>Spongia</i> IV. Sea-nettles <i>Acalephas</i>	6. Primitive intestinal animals 7. Sponges 8. Corals 9. Hood-jellies 10. Comb-jellies	6. <i>Gastræa</i> 7. Porifera 8. Coralla 9. Hydromedusæ 10. Ctenophora
C. Worms Vermes	V. Bloodless worms <i>Acaloma</i> VI. Blood-worms <i>Oelomati</i>	11. Primitive worms 12. Flat-worms 13. Round-worms 14. Arrow-worms 15. Wheel-animalcules 16. Moss-polyps 17. Mantle-animals 18. Acorn-worms 19. Star-worms 20. Ringed-worms	11. Archelminthes 12. Plathelminthes 13. Nematelminthes 14. Chaetognathid 15. Rotatoria 16. Bryozoa 17. Tunicata 18. Enteropneusta 19. Gephyrea 20. Annelida
D. Soft-bodied Animals Mollusca	VII. Headless shell-fish <i>Acephala</i> VIII. Head-bearing shell-fish <i>Eucephala</i>	21. Lamp-shells 22. Mussels 23. Snails 24. Cuttles	21. Spirobranchia 22. Lamellibranchia 23. Cochilides 24. Cephalopoda
E. Star-Animals Echinoderms	IX. Ring-d-arms <i>Cyathruchia</i> X. Armless <i>Lipobruchia</i>	25. Sea-stars 26. Sea-lilies 27. Sea-urchins 28. Sea-cucumbers	25. Asterida 26. Crinoida 27. Echinida 28. Holothurina
F. Articulated Animals Arthropoda	XI. Gill-breathers <i>Carides</i> XII. Tube-breathers <i>Tracheata</i>	29. Crabs 30. Spiders 31. Centipedes 32. Flies	29. Crustacea 30. Arachnida 31. Myriopoda 32. Insecta
G. Vertebrate Animals Vertebrata	XIII. Skull-less <i>Acrania</i> XIV. Single-nostrilled <i>Monorrhina</i> XV. Amnion-less <i>Anamnia</i> XVI. Amnion-animals <i>Amniota</i>	33. Tube-hearts (Lancelets) 34. Round-mouths (Lampreys) 35. Fishes 36. Mud-fish 37. Amphibians 38. Reptiles 39. Birds 40. Mammals	33. Leptocardia 34. Cyclostoma 35. Pisces 36. Dipneusta 37. Amphibia 38. Reptilia 39. Aves 40. Mammalia

TABLE XIX.

Phyletic Pedigree of the Animal Kingdom, founded on the Gastræa Theory and the Homology of the Germ-layers.²⁴



worm tribe; but the parent-forms of the three forms belong to worm-groups quite distinct from that of Chordonia. It is only far down at the common root of the group of Coelomati, that we assume a common source for these various tribal forms. (Cf. Tables XVIII and XIX.) It is especially necessary to remember that there is no direct blood-relationship between Vertebrates and Articulated Animals.

The Articulated Animals (*Arthropoda*), to which the most comprehensive of all classes of animals, that of Insects, and also the Spiders, Centipedes, as well as the Crabs, and Crustaceans, belong, are descendants of articulated Worms, the nearest allies of which are the extant Ringed Worms (*Annelida*). The tribe of Star-animals (*Echinodermata*), which includes the Star-fishes, Sea-lilies, Sea-urchins, and Sea-cucumbers, must also have descended from similar articulated Worms.¹¹⁸ The parent-form of the Soft-bodied Animals (*Mollusca*), which include the Cuttles, Snails, Mussels, and Lamp-shells, must also be sought among the Worms. But the Coelomati, from which these three higher animal tribes originated, differed entirely in character from the Chordate animals. Unlike the latter, they never developed a notochord. In them, the anterior section of the intestinal tube was never modified into a gill-body with gill-openings; nor were the upper throat-ganglia developed into a medullary tube. In a word, in Articulated Animals, Star-animals, and Soft-bodied Animals, as well as in their ancestors among the Blood-bearing Worms, the typical structural peculiarities which are exclusively characteristic of the vertebrate tribe and of their immediate invertebrate progenitors, were never present. Thus the great majority of all animals are

no way the subject of our further investigations, which is only concerned with the Vertebrates.

The development of the Vertebrates from the Invertebrates most nearly related to them, the Chorda-Animals, occurred millions of years ago, during the Archilithic Epoch. (See Table XII, p. 11.) This is unmistakably shown by the fact that the most recent sedimentary rock-strata which were deposited during that immense period of time, the higher layers of the Upper Silurian formation, contain remains of fossil Fishes (Primitive Fishes, *Selachii*). As these Fishes, although they belong to the lowest stage of the Skulled Animals (*Craniota*), yet possess a comparatively high organization, and as they must necessarily have been preceded by a long progressive series of lower Skullless Vertebrates, we must attribute the origin of the oldest skull-less Animals (*Acrania*) from the Chorda-animals to a much earlier part of the Archilithic Epoch. Therefore, not only all the invertebrate ancestors of our race, but also the earliest form of our vertebrate progenitors must have developed in that primordial time, which includes the Laurentian, Cambrian, and Silurian Periods. (Cf. Tables III, XIV., and XVI., pp. 12, 19, 44.)

Unfortunately, Palæontology can give us absolutely no information with regard either to the structure of our oldest vertebrate ancestors, or to the time of their appearance; for their bodies were as soft and as destitute of hard parts capable of fossilization, as were the bodies of all the preceding invertebrate ancestors. It is, therefore, not surprising, but quite natural, that we find no fossil remains of the former in the Archilithic formations. The fishes in which the soft cartilaginous skeleton was partly

modified into hard bone, are the earliest Vertebrates capable of leaving petrified records of their existence and structure.

Fortunately, this want is more than counterbalanced by the much more important testimony of Comparative Anatomy and Ontogeny, which henceforth form the safest guides within the Vertebrate pedigree. Thanks to the classic researches of Cuvier, Johannes Müller, Huxley, and especially of Gegenbaur, we are in possession of an extensive and instructive record of creation in this important branch of tribal history, that we can prove at least the more significant features in the development of Vertebrate ancestors, with the most gratifying certainty.

The characteristic peculiarities by which Vertebrates in general are distinguished from all Invertebrates, engaged our attention some time ago, when we examined the structure of the ideal Primitive Vertebrate (Figs. 52-56, p. 256). The most prominent characters were as follows: (1) the formation of the notochord between the medullary and intestinal tubes; (2) the differentiation of the intestinal tube into an anterior gill-intestine and a posterior stomach-intestine; (3) the inner articulation, or formation of metamers. The Vertebrates share the first two qualities with the larval Ascidians and with the Chorda-animals; the third quality is entirely peculiar to them. Accordingly, the most important structural advance, by which the earliest vertebrate forms originated from the most nearly allied Chorda-Animals, consisted in an internal metameric structure. This showed itself first most distinctly in the articulation of the muscular system, which broke up on the right and left into a series of consecutive muscular plates. At a later period the articulation declared itself prominently in the skeleton,

and blood-vessel systems. As we have already seen, this process of articulation, or metameric formation, is essentially to be regarded as terminal germination. Each distinct trunk-segment, or metameron, represents an individual. Thus the Vertebrates with their internal articulation stand in a similar relation to their inarticulate invertebrate ancestors, the Chorda Animals, as do the out-segmented Ringed Worms (*Annelida*) and Articulate Animals (*Arthropoda*) to the simple inarticulate animals from which they originated.

The tribal history of Vertebrates is rendered much more intelligible by the natural classification of the tribe which was proposed first in my *Generelle Morphologie* (1866), and afterwards improved in many ways in "The Natural History of Vertebrates" (Chap. XX., p. 192, etc.). In accordance with the existing Vertebrates must be divided into at least eight classes, as follows:—

SYSTEMATIC SURVEY OF THE EIGHT CLASSES OF VERTEBRATES.

Skull-less (<i>Acrania</i>)		1. Tube-hearted	1. Leptocardia ✓
Skulled (<i>Crania</i>)	a. Single-nostrilled (<i>Monorhina</i>)	2. Round-mouths	2. Cyclostoma
	b. Double-nostrilled (<i>Amphirhina</i>)	I.	
		Amnion-less	3. Fishes
		<i>Anamnia</i>	4. Mud-fishes
			5. Amphibians
		II.	
		With Amnion	6. Reptiles
		<i>Amniota</i>	7. Birds
			8. Mammals

The whole Vertebrate tribe may primarily be divided into the two main sections of the Skull-less and the Skulled Vertebrates. Of the earlier and lower section, that of the Skull-less (*Acrania*), the *Amphioxus* is alone extant. Of the more recent and higher section, the Skulled (*Crania*), belong all other existing Vertebrates up to Man. The

Craniota branched off from the Acrania, as these did the Chorda Animals. Our exhaustive study of the Comparative Anatomy and Ontogeny of the Ascidian and Amphioxus have already afforded proof of this relation. Chapters XIII. and XIV., and Plates X. and XI. with explanations.) I will only repeat, as the most important fact, that the Amphioxus develops from the egg in exactly the same way as the Ascidian. In both, the original gastrula (Figs. 4 and 10) originates in an exactly similar manner, by primordial cleavage from the simple parent (Figs. 1 and 7). From this originates that remarkable larva which develops a medullary tube on the dorsal side of intestinal tube, and between the two a notochord. At a later period, both in the Ascidian and in the Amphioxus intestinal tube differentiates into an anterior gill-intestine and a posterior stomach-intestine. In accordance with the fundamental principle of Biogeny, from these very important facts we may deduce the following statement of great phylogenetic importance: the Amphioxus, the lowest Vertebrate form, and the Ascidian, the most nearly allied Invertebrate form, have both descended from one single extinct Vertebrate form, which must have possessed the essential structure of the Chorda Animals.

- The Amphioxus, as has already been often shown, is of extreme importance; not only because it thus fills a great gap between the Invertebrates and the Vertebrates but also because it represents, at the present time, a typical Vertebrate in its simplest form; and because it directly affords the best standpoint from which to examine the gradual historic evolution of the whole tribe. If the structure and germ-history of the Amphioxus were

own to us, the whole subject of the development of the Vertebrate tribe, and thus of our own race, would be developed in an impenetrable veil. The accurate anatomical and ontogenetic knowledge of the Amphioxus, attained during the last few years, has alone pierced that heavy veil, formerly supposed to be impenetrable. If the Amphioxus is compared with the developed Man or any other of the higher Vertebrates, a great number of striking dissimilarities will be seen. The Amphioxus has no specialized head, no chin, no skull, no jaws, no limbs; it is without a central heart, a developed liver and kidneys, a jointed vertebral column; every organ appears in a much simpler and more primitive form than in the higher Vertebrates and in Man. (Table X., vol. i. p. 466.) And yet, in spite of all these obvious deviations from the structure of other Vertebrates,

Amphioxus is a genuine, unmistakable Vertebrate; and if, instead of the developed Man, the human embryo at an early period of its Ontogeny is compared with

Amphioxus, we shall find perfect parallelism between the two in all essential points. (Cf. Table IX., vol. i. p. 465.) This highly important parallelism justifies the conclusion that all the Skulled Animals (*Craniota*) have descended from a common primæval parent-form, the structure of which was essentially that of the Amphioxus. This parent-form, the earliest Primitive Vertebrate, possessed the peculiar characters of the Vertebrates, and yet was without those important peculiarities that distinguish the Skulled Animals from the Skull-less. Although the Amphioxus appears peculiarly organized in many respects, and although it may not be regarded as an unmodified descendant of the primitive Vertebrate, yet it must have inherited from the

latter the distinguishing characteristic features already mentioned. We cannot therefore say that the *Amphioxus* is the progenitor of the Vertebrates; but we may certainly say that the *Amphioxus* of all known animals is nearest all to this progenitor; both belong to the same limited family group, to the lowest Vertebrate class, that of the Skull-less Animals (*Acrania*). In the human pedigree, this group forms the ninth stage of the ancestral chain, the first among the Vertebrate ancestors. From this Skull-less group have developed the *Amphioxus* on the one side, and on the other the parent-form of the Skulled Animals (*Craniota*).

The comprehensive group of the Skulled Animals includes all known Vertebrates, with the single exception of the *Amphioxus*. All these Skulled Animals possess a distinct head, inwardly specialized from the trunk, and this contains a skull, enclosing a brain. The head also carries three of the higher sense-organs, which are partially wanting in the Skull-less Animals (nose, ears, and eyes). At first, the brain appears in a very simple form as an anterior bladder-like extension of the medullary tube (Plate XI. Fig. 16, *m*₁). This, however, is soon distributed into several transverse grooves—first into three, and afterwards into a series of five consecutive brain-bladders. In the formation of the head, skull, and brain, together with the higher sense-organs, lies the most essential advance made by the skulled parent-form beyond its skull-less ancestor. Other organs, however, also soon rose to a higher grade of development; a compact centralized heart appeared, a more perfect liver and kidneys; and in other directions also important advance was made.

The Skull-less Animals may be primarily subdivi-

differing main sections, that of the Single-nostrils (*Amphirhina*), and that of the Double-nostrils (*Amphirhina*). Former there are but very few extant forms, which are the Round-mouths (*Cyclostoma*). These are, however, of great interest, because in their whole structure they are intermediate between the Skull-less Animals and the Double-nostrils (*Amphirhina*). Their organization is much higher than that of the Skull-less Animals, much lower than that of the Double-nostrils; they thus form a very welcome genetic link between those two divisions. We may represent them as a special, tenth stage in the ancestral series.

Few existing species of the class of Round-mouths are divided into two different orders, which are distinguished as the Hagfishes and the Lampreys. The Hagfish (*Myxinoidei*) are long, cylindrical, worm-like bodies. Linnæus classed them among Worms, but later zoologists have placed them, as among the Fishes, sometimes Amphibians, and with Molluscs. The Hagfish live in the sea and are parasitic on Fishes, into the skin of which they enter by means of their round sucking mouths and their fleshy tongues. They are occasionally found in the vicinity of Fishes—for example, of the Cod and Sturgeon, having penetrated to the interior in their passage through the skin. The second order, that of the Lampreys (*Petromyzontes*), includes those well-known "Nine eyes," which live at the seaside; the little river Lamprey (*Petromyzon fluviatilis*) and the large sea Lamprey (*Petromyzon marinus*, Fig. 190).

The animals included in the two groups of the *Myxinoidei* and the *Petromyzontes*, are called Round-mouths

(*Cyclostoma*), from the fact that their mouth forms a circular or semi-circular opening. The upper and under jaws which appear in all the higher Vertebrates, are completely wanting in the Round-mouths, as in the *Amphioxus*. All other Vertebrates are therefore distinguishable from them as "Jaw-mouthed" (*Gnathostomi*). The Round-mouths may also be called "Single-nostrils" (*Monorhina*), because they have but a single nasal tube, while the *Gnathostomi* are all furnished with a pair of nasal cavities, a right and a left nose-cavity ("Double-nostrilled," *Amphirrhina*). But in addition to these peculiarities, the Jaw-mouths are also distinguished by many other remarkable structural arrangements, and are further removed from the Fishes than the latter are from Man. They must, therefore, evidently be regarded as the last remnant of a very old and very low class of Vertebrates, which are far below the structural stage of a genuine Fish. To mention here briefly only the most important, the Round-mouths are entirely without any trace of limbs. Their slimy skin is quite naked and smooth, without scales. They are wholly destitute of a bony skeleton. The inner skeleton axis is a very simple inarticulate notochord, like that of the *Amphioxus*. In the Lampreys alone a rudimentary articulation is indicated by the fact that upper arches appear in the vertebral tube proceeding from the notochord sheath. At the anterior end of the chorda a skull is developed in its very simplest form. From the notochord sheath proceeds a small soft-membraneous skull capsule, which becomes partly cartilaginous: this capsule encloses the brain. The important apparatus of the gill-arches, the tongue-bone, etc., which is inherited by all Vertebrate

shes to Man, is wholly wanting in round-mouths. They have, indeed, a small, cartilaginous gill-skeleton, but this is of different morphological significance. On the other hand, in them we meet, for the first time, with a brain, that important organ, which has been transmitted unchanged up to Man. It is true that in the Round-mouths the brain appears as a very small and comparatively insignificant swelling of the spinal chord; at the anterior end a large bladder (Plate XI. Fig. 16, m_1), which afterwards separates into five consecutive bladder-like swellings, as in the brains of all higher Vertebrates. These five simple primitive brain-blisters, which reappear in a similar form in the embryos of all higher Vertebrates, exist up to Man, and which undergo a very complex modification, remain in the Round-mouths, in a very low and undifferentiated stage of development. The histological structure of the nervous system is much more imperfect than in other Vertebrates.

While in the latter the organ of hearing always has three semi-circular canals, in lampreys it has but two, and in the Round-mouths but one. In most other points also, the organization of the Round-mouths is

Q.—The large Sea-lamprey (*Petromyzon marinus*), reduced in size. A series of seven gill-openings visible below the eye.



much simpler and more imperfect, as, for instance, in the structure of the heart, the circulatory system, and the kidneys. In them, as in the *Amphioxus*, the anterior portion of the intestinal canal does, indeed, form respiratory gills; but these respiratory organs are developed in a very peculiar way: in the form of six or seven little pouches, or sacs, which lie on both sides of the anterior intestine and communicate with the throat (*pharynx*) by inner openings and by outer ones with the external skin. This is a very peculiar formation of the respiratory organs, quite characteristic of this class of animals. They have therefore been called the "Pouch-gills" (*Marsupobranchii*). The absence of one very important organ found in the Fishes, the swimming-bladder, from which the lungs of the higher Vertebrates have developed, should be especially noticed.

In their germ-history, as in their whole anatomical structure, the Round-mouths present many peculiarities. They are even peculiar in the unequal cleavage of the egg, which most nearly approaches that of the Amphibians (Fig. 31, vol. i. p. 203). This results in the formation of a Hood-gastrula, like that of Amphibians (Plate II. Fig. 11). From this develops a very simple organized larval form, which is closely allied to the *Amphioxus*, and which, for that reason we examined and compared with the latter (vol. i. p. 428 and Plate VIII. Fig. 16). The gradual germ-evolution of these larvæ of the Round-mouths explains very clearly and unmistakably the gradual evolution of the Skulled from the Skull-less class of Vertebrates. At a later period, from this simple Lamprey larva is developed a blind and toothless larval form, which is so very different from the mature Lamprey that, until twenty years ago, it was generally

cribed as a peculiar form of fish under the name of *ammocetes*. By a further metamorphosis this blind and toothless *Ammocetes* is transformed into the Lamprey with eyes and teeth (*Petromyzon*).¹⁴⁷

Summing up all these peculiarities in the structure and embryology of the Round-mouths, we may assert that the first Skulled Animals, or *Craniota*, diverged in two lines; one of these lines has continued up to the present time almost little modified; it is represented by the *Cyclostoma*, or *Monorhina*, forming a collateral line which has made almost little progress, but has remained at a very low stage of development. The other line, the direct line in the pedigree of the Vertebrates, advanced in a straight line to the Fishes, and by new adaptations attained many important improvements.

In order rightly to appreciate the phylogenetic significance of interesting remnants of primæval groups of animals, such as the Round-mouths, it is necessary to study minutely their various peculiar characters philosophically and with the aid of Comparative Anatomy. A careful distinction must be drawn between the hereditary characters which have been accurately transmitted to the present day by heredity from common, primæval ancestors, now extinct, on the one hand; and, on the other, those special adaptive peculiarities which the existing remnant of that primæval group have, in the course of time, gained secondarily by adaptation. To the latter class belong, for example, in the Round-mouths, the peculiar formation of the single nostril and the round sucking mouth; as well as special structural arrangements of the epidermis and the pouch-shaped gills. But, on the other hand, to the

former class of characteristics, which alone have any phylogenetic significance, belong to the primitive formation of the vertebral column and the brain, the absence of the swim-bladder, of jaws, limbs, etc.

In the animal system, the Round-mouths are usually classed among Fishes; but that this is quite incorrect is apparent from the simple fact that, in all important prominent structural peculiarities, they are further removed from the Fishes than the Fishes are from the Mammals; from Man.

CHAPTER XVIII.

THE PEDIGREE OF MAN.

III. FROM THE PRIMITIVE FISH TO THE AMNIOTIC ANIMAL.

Comparative Anatomy of the Vertebrates.—The Characteristic Qualities of the Double-nostrilled and Jaw-mouthed: the Double-Nostrils, the Gill-arch Apparatus, with the Jaw-arches, the Swimming-bladder, the Two Pairs of Limbs.—Relationship of the Three Groups of Fishes: the Primitive Fishes (*Selachii*), the Ganoids (*Ganoides*), the Osseous Fishes (*Teleostei*).—Dawn of Terrestrial Life on the Earth.—Modification of the Swimming-bladder into the Lungs.—Intermediate Position of the Dipneusta between the Primitive Fishes and Amphibia.—The Three Extant Dipneusta (*Protopterus*, *Lepidosiren*, *Ceratodus*).—Modification of the Many-toed Fin of the Fish into the Five-toed Foot.—Causes and Effects of the latter.—Descent of all Higher Vertebrates from a Five-toed Amphibian.—Intermediate Position of the Amphibians between the Lower and Higher Vertebrates.—Modification or Metamorphosis of Frogs.—Different Stages in Amphibian Metamorphosis.—The Gilled Batrachians (*Proteus* and *Axolotl*).—The Tailed Batrachians (Salamanders and Mud-fish).—Frog Batrachians (Frogs and Toads).—Chief Group of the Amnion Animals, or *Amniota* (Reptiles, Birds, and Mammals).—Descent of all the *Amniota* from a Common Lizard-like Parent-form (*Protamnion*).—First Formation of the Allantois and of the Amnion.—Branching of the Amnion Animals in Two Lines: on the one side, Reptiles (and Birds), on the other side, Mammals.

"The imagination is an indispensable faculty; for it is that which, by forming new combinations, occasions important discoveries. The naturalist needs both the discriminating power of abstract reason, and the generalizing power of the imagination, and that the two should be harmoniously inter-

related. If the proper balance of these faculties is destroyed, the man is hurried into chimerical fancies by his imagination; while the man leads the gifted naturalist of sufficient strength of reason to the important discoveries."—JOHANNES MÜLLER (1834).

THE further we proceed in human tribal history, the lower does that part of the animal kingdom become to which we must look for extinct ancestors of the human race. At the same time, the evidence as to the history of the evolution of our race given by what we have called the records of creation, the evidence of Ontogeny, of Comparative Anatomy, and of Palæontology, grows constantly extensive, complete, and trustworthy. It is then natural that Phylogeny should assume a more definite form the nearer we approach the higher and the highest stages of the animal kingdom.

Comparative Anatomy especially has done far more to extend our knowledge of these higher stages of evolution in the animal kingdom than for the lower. This important science, which aims at a true philosophy of organic life, has made greater progress in the Vertebrate tribe than in any section of the Invertebrate. Cuvier, Meckel, and Johannes Müller had already laid a deep and extensive foundation, and now the Comparative Anatomy of Vertebrates has recently been powerfully advanced by the admirable investigations of Owen and Huxley, and, especially, has been perfected to such a degree by the unsurpassed labors of Gegenbaur, that it now forms one of the strongest supports of the Theory of Descent. Relying on the evidence furnished, we can now, with a great degree of certainty, recognize the most important outlines of the series of stages and the ramifications of the Vertebrate pedigree.

at part of the animal kingdom with which we are concerned has become so narrow, even before we have the Archilithic Epoch, that but a single one of the tribes of the animal kingdom forms the object of our study. Even within this tribe we have passed the lowest stage, and have risen above the Skull-less (*Acrania*) and Single-nostrilled Vertebrates (*Monorhina*), to the class of Double-nostrilled Vertebrates distinguished by mouths with jaws and by Double-nostrils (*Amphirhina*, or *Gnathostoma*). From Fishes we start again, as from that class of Vertebrates which are indisputably shown by Comparative Anatomy and Ontogeny to be the ancestral class of all higher Vertebrates, all Amphirhina. Of course no existing Fish can be regarded as the direct parent-form of the higher Vertebrates. But it is fully certain that from a common extinct Fish-like parent-form we may trace all those Vertebrates from Fishes to Man, which are included under the name of Amphirhina. If this primæval parent-form were extant, we could undoubtedly describe it as a genuine Fish and class it among Fishes. Fortunately, the Comparative Anatomy and Classification of the Fishes has been so far advanced thanks to the labours of Johannes Müller and Gegenbaur, that we can very clearly distinguish these most important and interesting genealogical relations.

In order correctly to understand the human pedigree within the Vertebrate tribe, it is very important to bear in mind the distinguishing characteristics, separating Fishes from all the other Double-nostrilled (*Amphirhina*) from Single-nostrilled and Skull-less Animals (*Monorhina* and *Acrania*). These very distinguishing characteristic marks

Fishes have in common with all other Double-nostrils to Man, and it is on this parallelism that we found claim of relationship to Fishes. (Cf. Table X, vol. p. 466.) The following characters of the Double-nostrils must be especially indicated as the systematic anatomical features of the highest importance: (1) the double structure of the nose; (2) the internal gill-arch apparatus, together with the jaw-arches; (3) the swimming-bladder, or lung; and (4) the two pairs of limbs.

As to the nasal structure, on which is based the distinction of the Single-nostrils (*Monorhina*) from the Double-nostrils (*Amphirrhina*), it is certainly significant that even in Fishes the earliest rudiment of the nose consists of two entirely distinct lateral grooves or pits in the outer surface of the head, just as is the case in the embryo of Man and of all higher Vertebrates. On the other hand, in Single-nostrilled and Skull-less Vertebrates the first rudiment of the nose is from the first, a single pit in the centre of the forehead region. No less important is the higher development of the skeleton of the gill-arch and of the jaw apparatus connected with it, as it occurs in all Double-nostrils from Fishes to Man. It is true that the primitive modification of the anterior intestine into the gill-intestine, which occurs even in Ascidians, is developed in all Vertebrates from one simple rudiment; and in this respect the gill-openings, which in all Vertebrates and also in Ascidians pierce the wall of the gill-intestine, are quite characteristic. But the external framework of the gills, which in all Skull-less and Single-nostrilled Animals (*Acraniota* and *Monorhina*) supports the gill-body, is displaced in all Double-nostrils (*Amphirrhina*) by an internal gill-skeleton which replaces the former

This internal gill-support consists of a consecutive series of cartilaginous arches, which are situated between the gill-openings within the wall of the throat (pharynx), and extend round the throat. The foremost of these pairs of gill-arches changes into the jaw-arch (maxillary arch), which gives rise to the upper and lower jaws.

A third essential character by which all Double-nostrils are well distinguished from all those lower Vertebrates which we have already considered, is the formation of a blind sac which protrudes from the anterior portion of the intestinal canal, and which in the Fishes becomes the air-filled swimming-bladder (Plate V. Fig. 13, *lu*). As this organ, in proportion as it contains a greater or less quantity of air, or in proportion as this air is more or less compressed, imparts a higher or lower specific gravity to the Fish, it acts as a hydrostatic apparatus. By this means the Fish can rise or sink in the water. This swimming-bladder is the organ from which the lung of higher Vertebrates has developed. The fourth and last main character of Double-nostrils is the presence of two pairs of extremities or members in the primitive arrangement of the embryo; a pair of fore limbs, which in Fishes are called pectoral fins (Fig. 191, *v*), and a pair of hind limbs, which in Fishes are called ventral fins (Fig. 191, *h*). The Comparative Anatomy of these fins is of supreme interest, because they contain the rudiments of all those parts of the skeleton which, in all the higher Vertebrates up to Man, form the skeleton or support of the extremities of the fore and hind limbs. In Skull-less and Single-nostrilled Animals there is, on the contrary, no trace of these extremities. In addition to these four most important main characters of the Amphi-

rhina, we might further mention the presence of a sympathetic nerve-system, a spleen, a ventral salivary gland, and other organs which are not represented in the lower Vertebrates already considered. All these important parts have transmitted themselves from Fishes up to Man, and from this circumstance alone it is evident how wide a chasm separates the Fishes from the Skull-less and Single-nosed Animals (*Acraniota* and *Monorhina*). Fishes and Man possess all these characters in common (Table X.).

Turning now to consider the Fish class in greater detail, we may divide it primarily into three main groups, or sub-classes, the genealogies of which are evident. The first and most ancient group is that of the Primitive Fishes (*Selachii*), the best-known extant representatives of which are the members of the much-varied orders of Sharks and Rays (Figs. 191, 192). These are followed by a series of further developed Fish forms, by the sub-class of Mucous Fishes (*Ganoides*). The greater number of these have long been extinct, and only very few living representatives are known; these are the Sturgeon and Huso of European seas, the Polypterus of African, and the Lepidosteus and Amia of American rivers. The earlier abundance of forms belonging to this interesting group is, however, proved by the abundance of their fossil remains. From these Mucous Fishes originated the third sub-class, that of the Osseous Fishes (*Teleostei*), to which belong most extant Fishes, especially nearly all our river fish. Comparative Anatomy and Ontogeny very clearly show that the Ganoids sprang from the Selachii, just as the Teleostei sprang from the Ganoids. But, on the other hand, a second side-line, or rather the main ascending line of the Vertebrate tribe,




FIG. 191.



FIG. 192.



FIG. 191.—Embryo of a Shark (*Scymnus lichia*), seen from ventral side; *v*, pectoral fins (in front of these five pairs of gill-openings); *h*, ventral fin; *a*, anal opening; *s*, tail fin; *k*, external gill-tufts; *d*, yolk-sac (the greater part of this has been removed); *g*, eye; *n*, nose; *m*, mouth fissure.

FIG. 192.—Developed Man-shark (*Carcharias melanopterus*), seen from the left side: *r*₁, first, *r*₂, second dorsal fin; *s*, tail fin; *a*, anal fin; *v*, pectoral fins; *h*, ventral fins.

developed in another direction from the Primitive Fishes this line leads upward through the Dipneusta group to the important class of Amphibia.

This significant relationship between the three groups of Fishes has been placed beyond all doubt by the researches of Gegenbaur on the subject. The lucid discussion on the "systematic position of the Selachii" which the author inserted in the introduction to his classic study of the "head skeleton of the Selachii," must be regarded as definitely proving this important relation.¹⁴⁸ In Primitive Fishes (*Selachii*), however, the scales (skin appendage) and the teeth (jaw appendages) are identical in formation and structure, while in the other two groups of Fishes (Mucous and Osseous Fishes) these organs have already become distinct and differentiated. Moreover, in Primitive Fishes, the cartilaginous skeleton (the vertebral column and the skull, as well as the members) is of the simple and most primitive nature, of which the bony skeleton of Mucous and Osseous Fishes must be regarded as a modification. It is true that in certain respects (in the structure of the heart and of the intestinal canal) Mucous Fishes fully coincide with Primitive Fishes, and differ from Osseous Fishes. But a comparative review of all the anatomical relations plainly shows that the Mucous Fishes constitute a connecting group between Primitive and

Osseous Fishes. The Primitive Fishes (*Selachii*) form the most ancient and original group of Fishes. From these, in one direction, all other Fishes have developed; the Mucous Fishes first, which, at a much later period (in the Jurassic, or the Chalk Period), gave rise to the Osseous Fishes. In another direction, the Primitive Fishes gave rise to the parent-forms of the higher Vertebrates, directly to the Dipneusta, and thus to Amphibians. Regarding the *Selachii* as forming the eleventh stage in our pedigree, these would be followed by the Dipneusta group as the twelfth stage, and by the Amphibian group as the thirteenth stage.

The advance effected in the development of the Mud-fishes (*Dipneusta*) from the Primitive Fishes is of great moment, and is connected with a very noticeable change, which took place in the beginning of the Palæozoic, or Primary Period in organic life as a whole. For the very numerous fossil remains of plants and animals which are now known to belong to the first three epochs of the history of the earth—to the Laurentian, the Cambrian, and the Silurian Periods, are exclusively those of aquatic plants and animals. From this palæontological fact, taken in connection with certain weighty geological and biological considerations, we may infer, with tolerable certainty, that at that time no land animals yet existed. During the whole of the enormous Archizoic Period—during many millions of years—the living population of our globe were all water-dwellers: a very remarkable fact, when it is remembered that this period embraces the larger half of the entire organic history of the earth. The lower animal tribes are even now exclusively, or with very few exceptions, aquatic. But during the Archizoic, or Primordial Epoch, the higher animal tribes

continued exclusively adapted to aquatic habits of life. It was not till later that they adopted a land life. The earliest fossils of terrestrial animals occur in the Devonian strata which were deposited in the beginning of the second great division of the earth's history (the Palæozoic Epoch). They increase greatly in number in the deposits of the Coal and Permian Periods. Even in these early formations many terrestrial and air-breathing species, both of the Arthropod and of the Vertebrate tribe, occur; while their aquatic ancestors of the Silurian Period breathed nothing but water. This physiologically significant modification of the mode of respiration is the most influential change that affected the animal organism in the transition from water to dry land. In the first place it caused the development of an air-breathing organ, the lung, the water-breathing gills having previously acted as respiratory organs. Simultaneously, however, it effected a remarkable change in the circulation of the blood and in the organs connected with this; for these are always most closely correlated with the respiratory organs. In addition to these, other organs also either in consequence of more remote correlation with the respiratory organs, or in consequence of new adaptation, were more or less modified.

Within the Vertebrate tribe it was undoubtedly a branch of the Primitive Fishes (*Selachii*) which, during the Devonian Period, made the first successful effort to accustom itself to terrestrial life and to breathe atmospheric air. In this the swimming-bladder was especially of service, for it succeeded in adapting itself to respiration of air, and so became a lung. The immediate consequence of this was the modification of the heart and nose. While true Fishes

have only two blind nose-pits on the surface of the head, these now became connected with the mouth-cavity by an open passage. A canal formed on each side, leading directly from the nose-pit into the mouth-cavity, and thus even while the mouth-opening was closed the necessary atmospheric air could be introduced into the lungs. While, moreover, in all true Fishes the heart consists simply of two compartments, an auricle, which receives the venous blood from the veins of the body, and a ventricle, which forces this blood through an arterial expansion into the gills, the auricle, owing to the formation of an incomplete partition wall, is now divided into a right and a left half. The right auricle alone now received the venous blood of the body, while the left auricle received the pulmonic venous blood passing from the lungs and the gills to the heart. The simple blood-circulation of the true Fishes thus became the so-called double circulation of the higher Vertebrates; and this development resulted, in accordance with the laws of correlation, in further progress in the structure of other organs.

The vertebrate class, which thus first adapted itself to the habit of breathing air, and which originated from a branch of the Selachii, are called Mud-fishes (*Dipneusta*), or Double-breathers, because, like the lowest Amphibia, they retain the earlier mode of breathing through the gills, in addition to the newly acquired lung-respiration. This class must have been represented by numerous and diverse genera during the Palæolithic Epoch (during the Devonian, Carboniferous, and Permian Periods). As, however, the skeleton is soft and cartilaginous, like that of the Selachii, they naturally left no fossil remains. The hard teeth of

single genera (*Ceratodus*) could alone endure; these, for instance, in the Trias. At the present time there are only three extant genera of this whole class: *Protopterus annectens*, in the rivers of tropical Africa (White Niger, Quillimane, etc.); *Lepidosiren paradoxa*, in the rivers of South America (in the tributaries of the Amazon); *Ceratodus Fosteri*, in the swamps of Southern Australia (Plate XII).¹⁴⁰ This wide distribution of the three immediate descendants of the class is alone sufficient to prove that they are the last remnants of a group which was formerly very widely developed. The whole structure of their bodies shows that the group to which they belong forms a transition between Fishes and Amphibia. The direct transitional structure between the two classes is so clearly expressed in the whole organization of these curious animals that zoologists yet dispute whether the Dipneusts are Fishes or Amphibia. Some well-known zoologists still place them among Amphibia, while they are usually placed among Fishes. In fact, the characters of both the classes are so united in the Dipneusta that the answer to the question as to their nature depends entirely upon the meaning being attached to the terms "Fish" and "Amphibian." In their mode of life they are true Amphibia. During the tropical winter, in the rainy season, they swim in the water like Fishes and inhale water through the gills. During the dry season they burrow in the mud as it dries up, and during that period breathe air through lungs, like Amphibians and higher Vertebrates. In this two-fold respiration they do, it is true, coincide with the lower Amphibia and stand far above Fishes. Yet, in most other characters they more nearly resemble the latter, and stand below






Fig. 2.

Ceratodus
Forsteri



former. Their external appearance is entirely like that of Fishes.

The head of the Dipneusta is not distinct from the trunk. The skin is covered with large fish-scales. The skeleton is soft, cartilaginous; its development has been arrested at a very low stage, just as in the lower Primitive Fishes. The notochord is retained entire. The two pairs of limbs are very simple fins of primitive structure, like those of the lowest Primitive Fishes. The structure of the brain, of the intestinal tube, and the sexual organs, is also as in Primitive Fishes. The Dipneusta, or Mud-fishes, have, therefore, by heredity, accurately retained many features of a lower organization derived from our primæval Fish ancestors, while their adoption of the habit of breathing air through lungs introduced a great advance in the vertebrate organization.

Moreover, the three extant Mud-fishes differ a good deal from one another in important points of structure. The Australian Mud-fish (*Ceratodus*), which was first described at Sidney in 1870 by Gerard Krefft, and which attains a length of six feet, appears in an especial degree to represent a primæval and very conservative animal form (Plate XII). This is especially true of the structure of its simple lung, and of its fins, which contain a pinnate skeleton. In the African Mud-fish (*Protopterus*), on the contrary, and in the American form (*Lepidosiren*) the double lung is present, as in all higher Vertebrates; nor is the fin-skeleton pinnate. In addition to the internal gills, *Protopterus* has also external gills, which are wanting in *Lepidosiren*. Those unknown Dipneusta, which were among our direct ancestors, and which formed the connecting link between the Selachii

TABLE XX.


Systematic Survey of the Phylogenetic Classification of Vertebrates.

I. Skull-less (<i>Acrania</i>), or Tube-hearted (<i>Leptocardia</i>). Vertebrates without a specialized head, skull, brain, or centralized heart.			
1. Skull-less <i>Acrania</i>	I. Tube-hearted <i>Leptocardia</i>	{ 1. Lancelets	1. Amphioxids

II. Animals with skulls (<i>Craniota</i>) and with centralized hearts (<i>Pachyo</i>). Vertebrates with specialized head, with skull and brain, and with centralized heart.			
Main-classes of the Skulled Animals.	Classes of the Skulled Animals.	Sub-classes of the Skulled Animals.	Systematic of the Sub-classes
2. Single- Rostrilled Monorhina	{ II. Round mouths <i>Cyclostoma</i>	2. Hags, or Mucous Fish	2. Hyperot (Myxinoi)
		3. Lampreys	3. Hyperos (Petromy)
3. Non- amniote Anamnia	{ III. Fishes <i>Pisces</i>	4. Primitive Fish	4. Selachii
		5. Ganoid Fish	5. Ganoide
		6. Osseous Fish	6. Teleoste
		IV. Mud-fishes <i>Dipneusta</i>	7. Single-lunged
8. Double-lunged	8. Dipneun		
	{ V. Batrachians <i>Amphibia</i>	9. Mailed Batra- chians	9. Phractan
		10. Naked Batra- chians	10. Lissamp
4. Amnion- Animals Amniota	{ VI. Reptiles <i>Reptilia</i>	11. Lizards	11. Lacertili
		12. Snakes	12. Ophidia
		13. Crocodiles	13. Crocodil
		14. Tortoises	14. Chelonias
		15. Sea-dragons	15. Haliosaur
		16. Dragons	16. Dinosaur
		17. Flying Reptiles	17. Pterosaur
		18. Beaked Animals	18. Anomoda
	{ VII. Birds <i>Aves</i>	19. Long-tailed	19. Saururæ
		20. Fan-tailed	20. Carinata
		21. Bush-tailed	21. Ratitæ
	{ VIII. Mammals <i>Mammalia</i>	22. Cloacal Animals	22. Monotres
		23. Pouched Ani- mals	23. Marsupia
		24. Placental Ani- mals	24. Placenta

TABLE XXI.

Pedigree of Vertebrates. (Cf. Plate XV.)



the most important of the Vertebrates, and group
the most important of the Fishes, is as yet entire-
ly unknown. It is only some future time it will ac-
quire an important importance as to the whole history
of the Vertebrate and of our ancestors.

Very important indication of this kind is
suggested by the main Vertebrate class, that of the
class *Amphibia* which is directly connected with
the *Lepidoptera* from which it originated. To this class
the *Amphibia* *Salamanders* *Plate XIII* *Toads* and
Barren after the example of *Linnæus* all *Reptiles*,
snakes *Crocodiles* and *Turtles* were also classed
Amphibia. But these animals are of a far higher or-
der, and in the most important characters of the
vertebral structure are more nearly allied to *Birds*
Amphibians. The true *Amphibia* on the other hand
more nearly allied to the *Double-breathers* and to *P*
Fishes: they are also much older than *Reptiles*. As
early as the *Carboniferous Period* numerous very
developed *Amphibia* (some of large size) were extant.

the yet extant Amphibians have a smooth and slippery skin. The latter, also, are on an average smoother than the former, and must be regarded as their stunted posterity.

Among the Amphibia of the present time we are, therefore, unable to find any forms that are directly referable to the pedigree of the human race, or that are to be regarded as ancestors of the three higher Vertebrate classes; yet, in important points of their internal anatomical structure, and especially in their germ-development, they correspond so closely with us, that we are justified in affirming that between the Double-breathers (*Dipneusta*) on the one hand, and the three higher Vertebrate classes (grouped together as Amniota) on the other, there existed a series of extinct intermediate forms which, if we had them before us, we should class among Amphibia. The whole organization of the extant Amphibia represents a transitional group of this kind. In the important matters of respiration and circulation of the blood, they are still closely allied to the Double-breathers, although in other respects they rise above the latter. This is especially true with respect to the advanced structure of their limbs or extremities. The latter here for the first time appear as feet with five digits. The thorough researches of Gegenbaur have shown that the fins of Fishes, concerning which very erroneous views were previously held, are feet with numerous digits; that is to say, the several cartilaginous or osseous rays, many of which occur in every Fish-fin, correspond to the fingers or digits on the limbs of higher Vertebrates. The several joints of each ray correspond to the several joints of each digit. In the Double-breathers the fin yet retains the same structure as in Fishes, and it was only gradually that the five-toed form of foot,

which occurs for the first time in Amphibians, was developed from this multi-digitate form. This reduction in the number of the digits from ten to five occurred in those Dipnoids which must be regarded as the parent-forms of the Amphibia, probably as early as the latter half of the Devonian Period, or, at latest, in the immediately subsequent Carboniferous Period. Several fossil Amphibia with five digits have already been found in the strata of the latter period. Fossil prints of the same animals are very numerous in the Trias (*Cherotherium*).

The great significance of the five digits depends on the fact that this number has been transmitted from the Amphibia to all higher Vertebrates. It would be impossible to discover any reason why in the lowest Amphibia, as well as in Reptiles and in higher Vertebrates up to the present time there should always originally be five digits on each anterior and posterior limbs, if we denied that here, as in all other cases, the cause of this phenomenon is the inheritance from a common five-fingered parent-form. In many Amphibia, certainly, as well as in many higher Vertebrates, we find less than five digits. But in all cases it can be shown that separate digits have retrograded and have finally been completely lost.

The causes which effected the development of the five-fingered foot of the higher Vertebrates in this Amphibian parent-form from the many-fingered foot, must certainly be found in the adaptation to the totally altered function which the limbs had to discharge during the transition from an exclusively aquatic life to one which was partly terrestrial. While the many-fingered fins of the Fish previously served almost exclusively to propel the

the water, they had now also to afford support to the body while creeping upon land. This effected a change both of the skeleton and of the muscles of the body.

The number of fin rays was gradually lessened, and finally reduced to five. These five remaining rays, however, developed more vigorously. The soft cartilaginous rays became hard bones. The rest of the skeleton also became considerably more firm. The movement of the body became not only more vigorous, but more varied. The separate portions of the skeleton and consequently those of the muscular system also, became more and more differentiated. Owing to the intimate connection of the muscular to the nervous system, the latter naturally made marked progress in point of function and structure. We therefore find that the brain is very more developed in the higher Amphibia than in the lower Mud-fishes, and in the lower Amphibia.

The organs which are most modified in consequence of the amphibious mode of life are, as we have already seen in the Lungs of the Lungs-breathers (*Dipneusta*), those of respiration and circulation of the blood. The first advance in evolution necessitated by the transition from aquatic to land habits of life was, of course, the formation of an air-breathing organ, a lung. This developed directly from the swimming-bladder which these animals had inherited from their Fishes. At first the function of this organ would be subordinate to the more ancient organ, used for the respiration of water, the gills. Hence we find that the Amphibia, the Gilled Amphibia, like the *Dipneusta*, spend the greater part of their lives in the water, and that they breathe water through gills. It is only

for brief intervals that they rise to the surface of the water or creep out of the water on to the land ; and at these times they breathe air through lungs. Some, however, of the Tailed Amphibians, the Axolotl and the Salamander, live exclusively in the water only when young, and afterwards usually remain on land. In the adult state they breathe only air through lungs. This is also the case with the most highly developed Amphibians, the Frog-amphibia (Frogs and Toads); some of the latter have even entirely lost the gilled larval form.¹⁵⁸ The same is true of a few small snake-like Amphibia, the Cæciliæ, which, like earth-worms, live in the ground.

The high degree of interest attached to the natural history of the Amphibian class is especially due to the fact that they hold a position exactly intermediate between the higher and the lower Vertebrates. While the lower Amphibia are in their whole organization directly allied to the Dipneusta and the Fishes, living mostly in the water and respiring water through gills, the higher Amphibia are no less directly related to the Amnion Animals, for, like the latter, they live mostly on land, and breathe air through lungs. But when young the higher forms resemble the lower, and only attain their own higher degree of development after undergoing complete modification. The individual germ-history of most higher Amphibians still accurately reproduces the tribal history of the whole class ; and the various stages of modification which were necessitated in certain low Vertebrates by the transition from aquatic to terrestrial habits during the Devonian or Carboniferous Period, are still to be seen every spring in each Frog as it develops from the egg in our ditches and pools.

Like the Tailed Salamanders (Fig. 193), each common Frog emerges from the egg in a larval form, totally different from that of the full-grown Frog (Fig. 194). The short




FIG. 193.—Larva of Spotted Land-Newt (*Salamandra maculata*), from the ventral side. In the centre a yolk-sac yet protrudes from the intestine. The external gills are prettily branched and tree-like. The two pairs of limbs are yet very small.

FIG. 194.—Larva of the Common Grass-Frog (*Rana temporaria*), a so-called tadpole: *m*, mouth; *n*, a pair of suction cups used in clinging to stones; *d*, skin-fold, which gives rise to the gill-roof; behind are the gill-openings, from which the gill branches protrude; *s*, tail-muscles; *f*, skin-fold of the tail, forming a float.

trunk is produced into a long tail, which in form and struc-

ture resembles the tail of a Fish (*s*). At first it has no limbs. Respiration is accomplished solely by gills, which are at first external (*k*) and afterwards internal. Correspondingly, the heart is also of the same form as in the Fishes, and consists of only two compartments—an auricle, which receives the venous blood of the body, and a ventricle, which drives it through the arterial bulb into the gills.

Numbers of these fish-like Frog larvæ, or "tadpoles," as they are called, swim about every spring in all ponds and pools, using their muscular tails for propulsion, just as is done by Fishes and larval Ascidiæ. The remarkable transformation of the fish-like form into that of the Frog does not take place till after the tadpole has grown to a certain size. From the throat grows a closed sac which develops into a pair of large sacs; these are the lungs. The simple chamber of the heart is divided into two auricles, owing to the formation of a partition wall, and simultaneously considerable changes of structure occur in the main arterial trunks. Previously all the blood passed from the heart-chamber through the aorta arches into the gills; but only part of it now passes to the gills, while another part passes through the newly formed lung arteries into the lungs. From the lungs arterial blood returns into the left auricle of the heart, while the venous blood of the body collects in the right auricle. As both of the auricles open into the simple ventricle, the latter contains mixed blood. The fish-like form has now passed into the Dipneusta form. During the further course of modification the gills, with the gill-vessels, are entirely lost, and respiration is now performed by the lungs alone. Yet later, the long tail is also

Fig. 1.



Fig. 2.



Fig 1. *Siredon pisciformis.*

Fig 2. *Salamandra maculata.*

The following is a list of the names of the persons who have been appointed to the various positions of the Board of Directors of the City of New York, for the year 1890. The names are given in alphabetical order, and the positions are given in parentheses.



ected, and the Frog now leaps about on the land on legs which have sprouted in the mean time.¹⁵⁰

This remarkable metamorphosis of the higher Amphibia is very instructive in its bearing on Man's ancestral history, and is especially interesting owing to the fact that the various groups of extant Amphibia have remained stationary at various stages of their tribal history, which, in accordance with the fundamental law of Biogeny, are reproduced in this germ-history. First, there is a very low order of Amphibia, the Gilled Batrachians (*Sozobranchia*), which, like fishes, retain their gills throughout life. To this order belong, among others, the well-known blind "Olm" of the Adelsberg Cave (*Proteus anguineus*), the Mud-eel of South Carolina (*Siren lacertina*), and the Axolotl of Mexico (*Sirenon pisciformis*; Plate XIII. Fig. 1). All these Gilled Batrachians are fish-like animals with long tails, and in point of respiratory organs and of circulation of the blood they remain throughout life stationary at the Dipneusta stage. They possess both gills and lungs, and can either respire water through the gills or air through the lungs, as occasion requires. In another order, the Salamanders, the gills are lost during metamorphosis, and in the adult state air only is breathed through lungs. This order bears the name of Tailed Batrachians (*Sozura*) because they retain the tail throughout life. To this order belong the common Water-Newts (*Triton*) which swarm in all ponds during the summer, and the black, yellow-speckled Land-Salamanders (*Salamandra*) found in damp woods (Plate XIII. Fig. 2). The latter are among the most remarkable of our indigenous animals, sundry anatomical characters proving them to be very ancient and highly conservative Vertebrates.¹⁵¹ A

few Tailed Batrachians retain the gill-opening in the skin of the neck, though the gills themselves are lost (*Mentopoma*). If the larvæ of the Salamanders (Fig. 193) and Tritons are compelled to remain in water, and not allowed to get on land, they may, under favourable conditions, be made to retain their gills. In this fish-like condition they become sexually mature, and will throughout life remain compulsorily in the lower stage of development of the Gilled Batrachians. The opposite experiment was made some years ago in the case of the Mexican Gilled Batrachian, the fish-like Axolotl (*Siredon pisciformis*; Plate XIII. Fig. 1). This animal had previously been regarded as a permanent Gilled Batrachian, remaining throughout life in this fish-like condition. But of the hundreds of these animals kept in the Jardin des Plantes at Paris, a few individuals, for some unknown reason, crept to land, lost their gills, and changed into a form closely allied to that of the Salamander (*Amblystoma*, Fig. 2). In this state they became sexually mature.¹⁵³ This phenomenon, which first excited a lively interest, has since been repeatedly observed with care. Zoologists regarded the fact as something peculiarly wonderful, though each spring even a common Frog and Salamander passes through the same modification. In these animals we can in the same way follow each step in the significant metamorphosis of the aquatic and gill-respiring animal into the terrestrial and lung-respiring animal. That which thus takes place in the individual during germ-evolution, took place in the same way in the whole class during the course of its tribal history.

The metamorphosis which takes place in the third order

Amphibia, the Frog Batrachians (*Batrachia*, or *Anura*), more complete than in the Salamanders. To these belong all the various kinds of Toads, Water-frogs, Tree-frogs, etc. In the course of transformation these lose not only the gills, but also the tail, which drops off in some cases earlier, in others later. In this respect the various species differ somewhat from one another. In most Frog Batrachians the larvæ drop the tail very early, and the frog-like form subsequently grows considerably.

Other species, on the contrary, as, for instance, the *Pelodytes paradoxus* of Brazil, as also an European Toad (*Pelodytes*) remain for a very long time in the fish form, retaining a lengthy tail till they have almost attained full size; hence, after their metamorphosis is complete, they appear much smaller than before. The opposite case is seen in some Frogs but recently brought under notice, which have lost the whole of their historic metamorphosis, and in which no tailed and gilled larva emerges from the egg, but the perfect Frog, without tail or gills.

Frogs inhabit isolated oceanic islands, the climate of which is very dry, and which are often for a considerable length of time without fresh water. As fresh water is indispensable for gill-respiring tadpoles, these Frogs have adapted themselves to this local deficiency and have completely relinquished their original metamorphosis, e.g., *Pelodytes martinicensis*.¹⁵³

The ontogenetic loss of gills and tail in Frogs and Toads can only be phylogenetically explained as owing to the fact that these animals have descended from long-tailed salamander-like Amphibians. This is also proved without doubt by the Comparative Anatomy of the two

development of the human embryo; all the many peculiarities in the development of the organs which we shall presently notice in detail; and, finally, the chief special arrangements of the internal structure of the body in the fully developed Amnion Animals; all these so clearly demonstrate the common origin of all Amnion Animals from a single extinct parent-form, that it is impossible to conceive their origin as polyphyletic, and that they originated from several independent parent-forms. This unknown common parent-form is the Primitive Amnion Animal (*Protamnion*). In external appearance the Protamnion was most probably an intermediate form between the Salamanders and the Lizards.

It was probably during the Permian Period that the Protamnion originated; perhaps at the beginning, perhaps at the close of that period. This we know from the fact that the Amphibia did not attain their full development till the Carboniferous Period, and that toward the close of the Permian the first fossil Reptiles make their appearance—or, at least, fossils (*Proterosaurus*, *Rhopalodon*) which must in all probability be referred to lizard-like Reptiles. Among the great and pregnant modifications of the vertebrate organization determined during this period by the development of the first Amnion Animals from salamander-like Amphibians, the three following are especially important: the total loss of water-breathing gills and modification of the gill-arches into other organs; the formation of the allantois, or primitive urinary sac; and, finally, the development of the amnion.

The total loss of the respiratory gills must be regarded as one of the most prominent characters of all Amnion

Animals. All these, even such as live in the water, *e.g.*, whales, respire only air through lungs, never water through gills. While all Amphibians, with very few exceptions, in the young state retain their gills for a longer or shorter period, and breathe through gills for some time (if not always), from this point gill-respiration entirely ceases. Even the Protamnion must have entirely ceased to breathe water. The gill-arches, however, remain, and develop into very different organs (partly rudimentary); into the various parts of the tongue-bone, into certain portions of the jaw apparatus, the organ of hearing, etc. But no trace of gill-leaves, of real respiratory organs on the gill-arches, are ever found in the embryo of Amnion Animals.

With this total loss of the gills is probably connected the formation of another organ, which we have already described as occurring in human Ontogeny; this is the allantois, or primitive urinary sac. (See vol. i. p. 379.) In all probability the urinary bladder of the Dipneusta is to be regarded as the first beginning of the allantois. Even in the American Mud-fish (*Lepidosiren*) we find an urinary bladder, which grows from the lower wall of the posterior extremity of the intestine, and serves as a receptacle for the renal secretions. This organ has been inherited by the Amphibia, as may be seen in any Frog. But it is only in the three higher Vertebrate classes that the allantois attains a special development; in these it protrudes at an early period from the body of the embryo, forming a large sac filled with liquid, and traversed by a considerable number of large blood-vessels. This sac also discharges a portion of the nutritive functions. In the higher Mammals and in Man the allantois afterwards forms the placenta.

The formation of the amnion and the allantois, together with the total loss of the gills and the exclusive adoption of lung-respiration, are the most important characters by which all Amnion Animals are distinguished from the lower Vertebrates which we have been considering. In addition to these there are a few subordinate characters which are constantly inherited by Amnion Animals, and are altogether wanting in animals without an amnion. One striking embryonic character of the Amnion Animals is the great curvature of the head and neck of the embryo. In the Anamnia the embryo is from the first either nearly straight, or else the whole body is bent in a sickle-shaped curve corresponding to the curvature of the yolk sac, to which the embryo is attached by its ventral surface; but there are no marked angles in the longitudinal axis (Plate VI. Fig. *F*). In all Amnion Animals, on the contrary, the body is very noticeably bent at an early age, so that the back of the embryo is much arched outwards, the head pressed almost at right angles against the breast, and the tail inclined on to the abdomen. The tail extremity, as it bends inwards, approaches so near to the frontal side of the head, that the two often nearly touch (Plates VI. and VII.). This striking triple curvature of the embryonic body, which has already been considered when we studied the Ontogeny of Man, and in which we distinguished the skull-curve, neck-curve, and tail-curve (vol. i. p. 371), is a characteristic peculiarity common to the embryos of all Reptiles, Birds, and Mammals. But in the formation of many internal organs also, an advance is observable in all the Amnion Animals which ranks them above the highest of the non-amnionate forms. Above all, a partition wall forms

in the simple ventricle of the heart, dividing it into a right and a left ventricle. In connection with the complete metamorphosis of the gill-arches, a further development of the organ of hearing takes place. A considerable advance is noticeable in the development of the brain, the skeleton, the muscular system, and other parts. Finally, the construction of the kidneys must be regarded as a most important modification. In all the lower Vertebrates as yet considered, we have found the primitive kidneys, which appear very early in the embryos of all higher Vertebrates. In Man, acting as a secretory or urinary apparatus. In lower Animals, however, these early primitive kidneys perform their function at an early period of embryonic life, and are superseded by the permanent "secondary kidneys," which are situated at the terminal portion of the primitive kidney.

Looking back at the whole of these characters of Amnion animals, it is impossible to doubt that all animals of this class, all Reptiles, Birds, and Mammals, had a common ancestor, and constitute a single main division of kindred animals. To this division belongs our own race. In his organization and germ-history Man is a true Amnion animal, and, in common with all other Amniota, has descended from the Protamnion. Although this whole class originated at the end, or perhaps even in the middle, of the Palæozoic Epoch, it did not attain its full development and its full perfection till the Mesozoic Epoch. In the two classes of Birds and Mammals then first appeared. It was in the Mesozoic Epoch, which is, therefore, called the "Age of Reptiles." The unknown and extinct Protamnion, the

parent-form of the entire group, must have been very nearly allied to the Reptiles in its whole organization, even though it cannot be regarded as a true Reptile in the proper meaning of the term. Of all known Reptiles, certain Lizards are most nearly allied to the Protamnion; and in the outward form of its body we may imagine the latter as an intermediate form between the Salamander and the Lizard.¹⁵⁶

The Comparative Anatomy and Ontogeny of the Protamnionate group clearly explains its genealogy. The group which directly descended from Protamnion gave rise to two divergent branches. The first of these, which will in future receive our whole attention, forms the Mammalian group. The other branch, which assumed an entirely different course of progressive development, and which is connected with the mammalian branch only as the root, is the comprehensive group constituted by Reptiles and Birds. The latter forms may be classed together as *Monocondylar Sauropsides*. The common parent-form of these is an extinct lizard-like Reptile. From this, the Serpents, Crocodiles, Tortoises, Dragons, etc.—in short, all the various members of the Reptilian group—developed in different directions. The remarkable group formed by the Birds also developed directly from an offshoot of the Reptilian group, as is definitely proved. Down to a late time the embryonic Reptiles and of Birds are yet identical, and even later are in some respects surprisingly similar. (See Plate VI, *T* and *C*.) In their entire organization the resemblance between the two is so great that no anatomist now doubts that the Birds originated from Reptiles. The Mammalian line is connected at its roots with the Reptilian line

nds diverged entirely from the latter, and developed
ntirely peculiar direction. The highest result of the
ment of the Mammalian line is Man, the so-called
of Creation."

CHAPTER XIX.

THE PEDIGREE OF MAN.

IV. FROM THE PRIMITIVE MAMMAL TO THE APE.

The Mammalian Character of Man.—Common Descent of all Mammals from a Single Parent-form (Promammalian).—Bifurcation of the Amnion Animals into Two Main Lines: on the one side, Reptiles and Birds, on the other, Mammals.—Date of the Origin of Mammals: the Trias Period.—The Three Main Groups or Sub-classes of Mammals: their Genealogical Relations.—Sixteenth Ancestral Stage: Cloacal Animals (*Monotremata*, or *Ornithodelphia*).—The Extinct Primitive Mammals (*Promammalia*) and the Extant Beaked Animals (*Ornithostoma*).—Seventeenth Ancestral Stage: Pouched Animals (*Marsupialia*, or *Didelphia*).—Extinct and Extant Pouched Animals.—Their Intermediate Position between Monotremes and Placental Animals.—Origin and Structure of Placental Animals (*Placentalia*, or *Monodelphia*).—Formation of the Placenta.—The Deciduous Embryonic Membrane (*Decidua*).—Group of the *Indecidua* and of the *Deciduata*.—The Formation of the *Decidua (vera, serotina, reflexa)* in Man and in Apes.—Eighteenth Stage: Semi-apes (*Prosimia*).—Nineteenth Stage: Tailed Apes (*Momocerca*).—Twentieth Stage: Man-like Apes (*Anthropoides*).—Speechless and Speaking Men (*Mali. Homines*).

"A century of anatomical research brings us back to the conclusion of Linnaeus, the great lawgiver of systematic zoology, that man is a member of the same order as the apes and lemurs. Perhaps no order of mammals presents us with so extraordinary a series of gradations as this, leading us insensibly from the crown and summit of the animal creation down to creatures from which there is but a step, as it seems, to the lowest, smallest, and least intelligent of the placental mammalia. It is as if nature herself

"MAN'S PLACE IN NATURE."

had foreseen the arrogance of man, and with Roman severity had that his intellect, by its very triumphs, should call into prominence, admonishing the conqueror that he is but dust."—THOMAS (1863).

AMONG those zoological facts which afford us points of support in researches into the pedigree of the human race, the position of Man in the Mammalian class is one of the most important and fundamental. Much as zoologists have long disagreed in their opinions as to Man's particular place in this class, and especially in their ideas of his relation to the most nearly related group, that of the Apes, yet no naturalist has ever doubted that Man is a genuine Mammal in the whole structure and development of his body. Every anatomical museum, every manual of Comparative Anatomy, affords proof that the structure of the human body shares all those peculiarities which are common to all Mammals, and by which the latter are definitely distinguished from all other animals.

Now, if we examine this established anatomical fact phylogenetically, and in the light of the Theory of Descent, we arrive immediately at the conclusion that Man is of a common stock with all the other Mammals, and springs from a root common to them. The various characteristics in which all Mammals coincide, and in which they differ from all other animals, are, moreover, of such a kind, that a polyphyletic hypothesis appears in a special degree inadmissible in their case. It is inconceivable that all existing and extinct Mammals have sprung from several different and originally separate root-forms. We are compelled, if we in any way acknowledge the Theory of Evolution, to assume the monophyletic hypothesis, that all Mammals,

including Man, must be traced from a single common mammalian parent-form. This long extinct primæval root-form and its immediate descendants—which differ from each other hardly more than do several species of one genus—we will call Primitive Mammals (*Promammalia*). As we have already seen, this root-form developed from the ancient parent-form of the Primitive Amnion Animals in a direction wholly different from that followed by the Reptile group, which afterwards gave rise to the more highly developed class of Birds. The differences which distinguish Mammals on the one side, from Reptiles and Birds on the other, are so important and characteristic, that we may quite safely assume a bifurcation of this kind in the vertebrate family tree. Reptiles and Birds—which we classed together as *Monocodylea*, or *Sauropsida*—coincide entirely, for instance, in the characteristic structure of the skull and brain, which is strikingly dissimilar from that of the same parts in Mammals. In Reptiles and Birds, the skull is connected with the first cervical vertebra (the atlas) by a single joint-process (condyle) of the occipital bone; in Mammals, on the contrary (as in Amphibians), the condyle is double. In the former, the under jaw is composed of many parts, and is connected with the skull by a peculiar bone of the jaw (the square bone) so as to be movable; in the latter, on the contrary, the lower jaw consists of but two bone-pieces, which are directly attached to the temporal bone. Again, the skin of the *Sauropsida* (Reptiles and Birds) is covered with scales or feathers, that of the Mammals with hair. The red blood-cells of the former are nucleated, those of the latter non-nucleated. The eggs of the former are very large, are provided with a large nutritive yelk, and undergo

discoidal cleavage resulting in a Disc-gastrula; the eggs of the latter are very small, and their unequal cleavage results in the formation of a Hood-gastrula. Finally, two characters entirely peculiar to Mammals, and by which these are distinguished both from Birds and Reptiles and from all other animals, are the presence of a complete diaphragm, and of the milk-glands (*mammæ*), by means of which the new-born young are nourished by the milk of the mother. It is only in Mammals that the diaphragm forms a transverse partition-wall across the body-cavity (*cæloma*), completely separating the chest from the ventral cavity. (Cf. Plate V. Fig. 16 z.) It is only among Mammals that the mother nourishes the young with her milk; and the whole class are well named from this.

These important facts in Comparative Anatomy and Ontogeny clearly show that the tribe of Amnion Animals (*Amniota*) bifurcated from the very first into two main diverging lines; on the one side, the Reptilian line, from which the Birds afterwards developed; on the other side, the Mammalian line. The same facts also prove as indubitably that Man originated from the latter line. For Man, in common with Mammals, shares all the characteristics we have mentioned, and is distinguished by them from all other animals. And, finally, these facts indicate as certainly those advances in vertebrate structure by which one branch of the Primitive Amnion Animals developed into the parent-form of Mammals. The most prominent of these advances were (1) the characteristic modification of the skull and brain; (2) the formation of a covering of hair; (3) the complete development of the diaphragm; and (4) the formation of the milk-glands and the adaptation to the suckling of

the young. Intimately connected with these, other important structural modifications gradually occurred.

The period at which these important advances, which laid the first foundation of the Mammalian class, took place may most probably be placed in the first part of Mesolithic, or Secondary Epoch, in the Triassic Period. For the oldest known fossil remains of Mammals, occur in the sedimentary rock-strata of the most recent deposits of the Triassic Period, in the upper Keuper. It is possible indeed, that the parent-forms of Mammals may have appeared earlier (perhaps even at the close of the Palæolithic Epoch, in the Permian Period). But no fossil remains of Mammals belonging to that period are as yet known. Throughout the Mesolithic Epoch, throughout the Triassic, Jurassic, and Calcareous Periods, fossil remains of Mammals are very scarce, and indicate a very limited development of the whole class. During this Mesolithic Epoch, Reptiles play the chief part, and Mammals are of quite secondary importance. It is, however, especially significant and interesting, that all mammalian fossil remains of the Mesozoic Epoch belong to the older and inferior divisions of Pouched Animals (*Marsupialia*), a few probably extend to the yet older division of the Cloacal Animals (*Metremata*). Among them, no traces of the third and most highly developed division of the Mammals, the Placental Animals, have as yet been found. The last, to which I belong, are much more recent, and their fossil remains do not occur till much later—in the succeeding Cænozoic Epoch; in the Tertiary Period. This palæontological fact is very significant, because it harmonizes perfectly with that order of the development of Mammals which is

stakably indicated by Comparative Anatomy and Ontogeny.

These show that the whole Mammalian class is divisible into three main groups, or sub-classes, corresponding to three successive stages of phylogenetic evolution. These three stages, which consequently represent three important ancestral stages in the human pedigree, were first distinguished in the year 1816 by the celebrated French zoologist, Blainville, who named them, according to the different structure of the female organs of reproduction, *Ornithodelphia*, *Didelphia*, and *Monodelphia* (ὄελφύς, which, being interpreted, is *uterus*). It is not, however, only in the varied structure of the sexual organs that these three classes differ from one another, but in many other respects also, so that we can safely maintain the important phylogenetic statement: The *Monodelphia*, or Placental Animals, have descended from the *Didelphia*, or Pouched Animals; and the latter, again, have descended from the Cloacal Animals, or *Ornithodelphia*.

Accordingly we have now to consider, as the sixteenth ancestral stage in the human pedigree, the oldest and lowest in group of Mammals; the sub-class of the Cloacal Animals (*Monotremata*, or *Ornithodelphia*). They are so named in consequence of the cloaca, which they have in common with the other lower Vertebrates. This so-called cloaca is the common excretory channel for the excrement, urine, and the sexual products (Fig. 327). For, in these Cloacal Animals, the urinary duct and the sexual ducts yet open into the posterior parts of the intestine, while in all other Mammals they are wholly separated from the rectum and anus, and open by a special orifice (*porus*

urogenitalis). The urinary bladder in the Monotremes also opens into the cloaca, and is separate from the two urinary ducts (Fig. 327, *ro*); in all other Mammals the latter open directly into the urinary bladder. The structure of the milk-glands, by means of which all Mammals suckle their new-born young for a time, is also quite peculiar in the Cloacal Animals. In them the milk-gland has no nipple which the young animal can suck; there is only a peculiar sieve-like place in the skin, perforated with holes through which the milk passes out, and from which the young animal has to lick it. For this reason they have also been called Nipple-less Mammals (*Amast*). Again, the brain of the Cloacal Animals has remained at a much lower stage of development than that of any other Mammal. The fore-brain, or cerebrum, is so small that it does not overhang the hind-brain, or cerebellum. In the skeleton (Fig. 196), the structure of the shoulder girdle, as well as of other parts, is remarkable, differing entirely from the same part in other Mammals, and resembling rather those of the lower Vertebrates, especially Reptiles and Amphibians. Like the latter, the Cloacal Animals have a well-developed coracoid bone (*coracoideum*), a strong bone uniting the shoulder-blade with the breast-bone. In all other Mammals the coracoid bone (as in Man) has degenerated, has coalesced with the shoulder-blade, and appears only as an insignificant process of the latter. These and many other less striking peculiarities prove beyond doubt that the Cloacal Animals occupy the lowest rank among Mammals, and represent a direct intermediate form between the Protamnia and other Mammals. All these marked Amphibian characters must have been present in the paren

a of the whole vertebrate class, in the Primitive animal, by which they must have been inherited from Primitive Amnion Animals.

During the Triassic and Jurassic Periods, the sub-class the Cloacal Animals seems to have been represented by Primitive Mammals of very varied form. At present is represented only by two isolated members, which grouped together as the Beaked Animal family (*Ornithomata*). Both of these are confined to Australia and the neighbouring island of Van Diemen's Land, or Tasmania; they are becoming less numerous year by year, and will be classed, with all their blood relations, among the extinct animals of our globe. One of these forms passes life swimming about in rivers, and builds subterranean dwellings on the banks: this is the well-known Duck-billed Platypus (*Ornithorhynchus paradoxus*): it is web-footed, has a thick, soft skin, and broad, flat jaws, which very much resemble a duck's bill (Figs. 195, 196). The other form, the Porcupine Ant-eater (*Echidna hystrix*), much resembles the Ant-eaters, in its mode of life, in the characteristic form of its slender snout, and in the great length of its tongue; it is covered with prickles, and can roll itself into a ball like a hedgehog. Neither of these extant Beaked Animals possesses true bony teeth, and, in this respect, they resemble the Toothless Mammals (*Edentata*). The absence of teeth, together with other peculiarities of the Ornithostomata, is probably the result of comparatively recent adaptation. Those extinct Cloacal Animals which embraced the parent-forms of the whole Mammalian class, the Promammalia, must certainly have been provided with a developed set of teeth, inherited from Fishes.¹⁵⁷ Some



FIG. 195.—The Duck-billed Platypus (*Ornithorhynchus paradoxus*).

FIG. 196.—Skeleton of Platypus.

all single molars, found in the uppermost strata of the Keuper formation in England and Würtemberg, and which are the oldest known vertebrate remains, probably belonging to these primæval Promammalia. These teeth, by their form, indicate species that lived on insects; the species has been called *Microlestes antiquus*. Teeth belonging to a more closely allied Primitive Mammal (*Dromatherium strepsirrhin*) have recently been discovered in the North African Trias.

On the one hand, the still extant Beaked Animals, and, on the other, the parent-forms of the Pouched Animals (*Marsupialia*, or *Didelphia*), must be regarded as representing distinct and divergent lines of descent from the Promammalia. This second Mammalian sub-class is very interesting as a perfect link between the two other sub-classes. While the Pouched Animals, on the one side, retain many of the characters of the Cloacal Animals, they also, on the other, possess many placental characters. A few characters are quite peculiar to Pouched Animals alone; such, for instance, is the structure of the male and female genital organs, and the form of the lower jaw. The distinctive feature of the latter in these Pouched Animals is a peculiar hook-shaped bony process, passing inward horizontally from the angle of the lower jaw. As neither Cloacal Animals nor Placental Animals have this process, this structure is alone sufficient to distinguish the Pouched Animals (*Marsupialia*). Nearly all the known mammalian fossils from the Jurassic and Cretaceous formation are lower classes. Our whole knowledge of numerous mesolithic mammals, the former existence of which would otherwise never have been known, is solely derived from their fossilized

lower jaws, no fragment of the rest of their bodies having been reserved. According to the logic usually applied to palæontology by the "exact" opponents of the theory of evolution, the inference drawn from this fact would be that these Mammals had no bones except lower jaws. The remarkable circumstance is, after all, very easily accounted for. The lower jaw of Mammals being a solid and exceptionally hard bone, but very loosely attached to the skull, it is easily detached from the carcase as the latter is carried down by some river, and, falling to the bottom, is retained in the mud. The rest of the carcase is carried on further, and is gradually destroyed. As all the mammalian lower jaws found, in England, in the Jurassic strata of Stonesfield and Purbeck, exhibit this peculiar process characteristic of the Pouched Animals (*Marsupialia*), we may infer, from this palæontological fact, that they belonged to Marsupials. No Placental Animals appear to have existed during the Mesolithic Epoch. At least no fossil remains, undoubtedly belonging to these and dating from that epoch, are known.

The extant Pouched Animals, the most generally known of which are the gramnivorous Kangaroos and the carnivorous Pouched Rats, display very considerable difference in their organization, in the form of their bodies and in size, and in many respects correspond to the several orders of Placental Animals. The great majority of them live in Australia, in New Holland, and in a few of the Australian and South Asiatic islands; some few species occur in America. On the other hand, there is no longer a single indigenous Pouched Animal on the continents of Asia, of Africa, or of Europe. The case was very different during the Mesolithic, and also during the earlier Cærolithic Epochs

The Neptunian deposits of these epochs in all quarters of the globe, and even in Europe, contain abundant marsupial remains in great variety, some of them being of very large size. From this we may infer that the extant Pouched Animals are but the last remnant of a group which was once much more widely developed, and which was distributed over the whole surface of the globe. During the Tertiary Period, these succumbed in the struggle for life with the stronger Placental Animals, and the survivors were gradually driven back by the latter into their present restricted area.

From the Comparative Anatomy of the extant Pouched Animals, very important conclusions may be drawn as to their phylogenetic intermediate position between Cloacal Animals and Placental Animals. The incomplete development of the brain, especially of the fore-brain (*cerebrum*), the possession of marsupial bones (*ossa marsupialia*), the simple structure of the allantois (which does not as yet develop a placenta), with many other characters, have been inherited by the Pouched Animals from Cloacal Animals. On the other hand, they have lost the independent coracoid bone (*os coracoideum*) attached to the shoulder girdle. A more important step consists in the fact that a cloaca is no longer formed; the cavity of the rectum, together with the anal opening, is separated by a partition wall from the urinary and sexual opening (*sinus urogenitalis*). Moreover, all Pouched Animals develop special nipples on the milk-glands, which are sucked by the young after birth. These nipples project into the cavity of a pouch, or marsupium, in the ventral side of the mother. This pouch is supported by a couple of marsupial bones. In it the young, which are

born in a very imperfect condition, are carried by the mother for a long time; until, in fact, they are completely developed (Fig. 197). In the large Giant Kangaroo, which



FIG. 197.—The Crab-eating Pouched Rat (*Phitander cancrivorus*). A female with two young in its pouch. (After Brehm.)

attains the height of a man, the embryo develops in the uterus but for a month; it is then born in a very incomplete condition, and attains all its further development in the mother's pouch, where, for about nine months, it remains attached to the milk-glands.

All these and other characters (especially the peculiar structure of the internal and external sexual organs of the male and female) clearly show that the whole sub-class of the Pouched Animals (*Marsupialia*) are a single group, which originated from the promammalian branch. From a branch of these Pouched Animals (perhaps from several branches) the parent-forms of the higher Mammals, the Placental Animals, afterwards sprang. Hence we must reckon a whole series of Pouched Animals among the ancestors of the human race; and these constitute the seventeenth stage in the human pedigree.¹⁵⁸

The remaining stages of our ancestral line, from the eighteenth to the twenty-second, all belong to the group of Placental Animals (*Placentalia*). This very highly developed group of Mammals, the third and last, came into the world at a considerably later period. No single known fossil, belonging to any portion of the Secondary or Mesolithic Epoch, can be referred with certainty to a Placental Animal, while we have plenty of placental fossils dating from every part of the Tertiary or Cærolithic Epoch. From this palæontological fact we may provisionally infer that the third and last main division of Mammals did not develop from the Pouched Animals until the beginning of the Cærolithic Epoch, or, at the earliest, till the close of the Mesolithic Epoch (during the Chalk Period). In our survey of geological formations and periods (pp. 12, 19) we found

how comparatively short this whole Tertiary or Cænozoic Epoch was. Judging from the relative thicknesses of various strata-formations we were able to say that whole period, during which Placental Animals first appeared and assumed their respective forms, amounted at most about three per cent. of the entire duration of the organic history of the earth. (Cf. p. 18.)

All Placental Animals are distinguished from the lower Mammalian groups already considered, from Cloacal Animals and Pouched Animals, by many prominent peculiarities. All these characters are present in Man, the most significant fact. For on the most accurate comparative anatomical and ontogenetical researches, we may base an irrefutable proposition that Man is in every respect a Placental Animal; in him are present all those peculiarities in the structure and in the development of the body which distinguish Placental Animals from the lower Mammalian groups, and at the same time from all other animals. Among these characteristic peculiarities the higher development of the brain, the organ of the mind, is especially prominent. The fore-brain, or large brain (*cerebrum*), much more highly developed in these than in the lower animals. The body (*corpus callosum*), which, like a bridge connects the two hemispheres of the fore-brain, attains full development only in Placental Animals; in the Pouched Animals and Cloacal Animals it exists merely as an insignificant rudiment. It is true that in their brain structure the lowest of the Placental Animals yet resemble Pouched Animals very nearly; but within the Placental group we can trace a continuous series of progressive stages in the development of the brain, ascending quite gradually

the lowest stage to the very highly developed mind-organ of the Monkey and of Man. (Cf. Chapter XX.) The human mind is but a more highly developed ape-mind.

The milk-glands of Placental Animals, as of Marsupials, are provided with developed nipples; but the pouch in which the immature young of the latter are carried about and suckled is never present in the former. Nor are the marsupial bones (*ossa marsupialia*) present in Placental Animals; these bones, which are embedded in the abdominal wall, and rest on the anterior edge of the pelvis, are common to Pouched Animals and Cloacal Animals, originating from a partial ossification of the tendons of the inner oblique muscle of the abdomen. It is only in a few beasts of prey that insignificant rudiments of these bones are found. The hook-shaped process of the lower jaw, which characterizes Pouched Animals, is also entirely wanting in Placental Animals.

The character, however, which especially distinguishes Placental Animals, and which has justly given its name to the entire sub-class, is the development of the placenta, or vascular cake. We have already spoken of this organ, in describing the development of the allantois in the human embryo (vol. i. p. 382). The urinary sac or allantois, that peculiar bladder which grows out of the posterior portion of the intestinal canal, is, we found, formed at an early stage in the human embryo just as in the germs of all other Amnion animals. (Cf. Figs. 132-135, vol. i. p. 377-380.) The thin wall of this sac consists of the same two layers, or skins, as the wall of the intestine itself; internally of the intestinal-glandular layer, and externally of the intestinal-fibrous layer. The cavity of the allantois is filled with fluid; this primi-

tive urine must be chiefly the product of the primitive kidneys. The intestinal fibrous layer of the allantois is traversed by large blood-vessels which accomplish the nutriment and, especially, the respiration of the embryo; these are the navel-vessels, or umbilical vessels (vol. i. p. 400). In all Reptiles and Birds the allantois becomes an immense sac, which encloses the embryo with the amnion, and which does not coalesce with the outer covering of the egg (*chorion*). In Cloacal Animals (*Monotremata*) and Pouched Animals (*Marsupialia*) the allantois is also of this nature. It is only in Placental Animals that the allantois develops into that very peculiar and remarkable formation, called the *placenta*, or "vascular cake." The nature of the placenta is this: the branches of the blood-vessels which traverse the wall of the allantois, penetrate into the hollow tufts of the chorion, which are inserted into corresponding depression in the mucous membrane of the maternal uterus. As this mucous membrane is also abundantly supplied with blood vessels, which conduct the mother's blood into the uterus and as the partition between these maternal blood-vessels and the embryonic vessels in the chorion-tufts soon become extremely thin, a direct exchange of substance is soon developed between the two sets of blood-vessels, which is of the utmost importance for the nutrition of the young Mammal. The maternal blood-vessels do not, however, pass directly (anastomosis) into the blood-vessels of the embryonic chorion-tufts, so that the two kinds of blood simply mix, but the partition between the two sets of vessels becomes so thin, that it permits the passage of the most important food-materials, freed from unnecessary matter (transudation, or diosmosis). The larger the embryo

rows in Placental Animals, and the longer it remains in the maternal uterus, the more necessary does it become that special structural arrangements should meet the increased consumption of food. In this point there is a very striking difference between the lower and the higher Mammals. In Locustal Animals and Pouched Animals, in which the embryo remains for a comparatively brief time in the uterus, and is born in a very immature condition, the circulation as it exists in the yolk-sac and in the allantois suffices for nutrition, as in birds and reptiles. In Placental Animals, on the contrary, in which gestation is very protracted, and the embryo remains much longer in the uterus, there attaining its full development within its investing membranes, a new apparatus is required to convey a direct supply of richer nutritive matter; and this is admirably effected by the development of the placenta.

In order rightly to understand and appreciate the formation of this placenta and its important modifications in different Placental Animals, we must once more glance at the external coverings of the mammalian egg. The outermost of these was originally, and during the cleavage of the egg and the first formation of the axial portion of the germ, formed by the so-called *zona pellucida*, and by the thick albuminous covering deposited externally on the latter (Fig. 19, Fig. 21, *z, h*, vol. i. p. 178).

We called these two outer coverings, which afterwards amalgamate, the *prochorion*. This prochorion very soon disappears (in man perhaps in the second week of development), and is replaced by the permanent outer egg-membrane, the chorion. The latter, however, is simply the serous membrane, which, as we have already seen, is the

product of the outer germ-layer of the germ-membrane vesicle. (See vol. i. p. 401, and Fig. 139, 4, 5, *sh*, p. 385.) It is at first a very smooth, thin membrane, surrounding the entire egg, as a closed spherical vesicle, and consisting of a single layer of exoderm cells. The chorion, however, very soon becomes studded with a number of little processes or tufts (Fig. 139, 5, *chz*). These fit themselves into the indentations in the mucous membrane of the uterus, thus securing the egg to the wall of the latter. They are, however, not solid, but hollow, like the finger of a glove. Like the whole chorion, these hollow tufts consist of a thin layer of cells belonging to the horn-plate. They very soon attain an extraordinary development, grow and branching rapidly. In the spaces between them




FIG. 198.—Egg-covers of the human embryo (mammalian): *m*, the thick fold of the uterus; *plu*, the inner stratum; *chf*, tufted, *chl*, smooth, the chorion; *a*, amnion; *as*, amniotic cavity; *as*, amnion; *dg*, yolk-duct; *ds*, decidua (decidua vera); *dv*, *dr*, decidua (decidua falsa). The uterus opens below into the vagina; above, on the right hand, into an oviduct (*t*). (Kölliker.)

tufts arise in all directions from the serous membrane, thus before long (in the human embryo in the third

the whole outer surface of the egg is covered with a dense forest of tufts (Fig. 134).

These hollow tufts are now penetrated from within by the branching blood-vessels, which originate from the intestinal fibrous layer of the allantois, and which contain the blood of the embryo, introduced through the navel vessels (Fig. 198, *chz*). On the other hand, dense networks of blood-vessels develop in the mucous membrane, which lines the inner surface of the uterus, particularly in the neighbourhood of the depressions into which the chorion-tufts penetrate (*plu*). These vascular networks receive the blood of the mother introduced through the uterus vessels. The whole mass of these two sets of vessels, which are here most intimately connected, together with the connecting and enveloping tissues, is called the placenta, or "vascular cake." Properly speaking, the placenta consists of two quite different, though closely connected, parts; internally, of the embryonic placenta (*placenta fœtalis*, Fig. 198, *chz*), and externally of the maternal placenta (*placenta uterina*, Fig. 198, *plu*). The latter is formed by the uterine mucous membrane and its blood vessels: the former by the secondary chorion and the navel vessels of the embryo.

The mode in which these two "vascular cakes" combine to form the placenta, as well as the structure, form, and size of the latter, differs much in different Placental animals, and affords valuable data for natural classification, and hence also for the tribal history of the whole sub-class. The latter is primarily divisible into two main divisions, based on these differences: the lower Placental Animals, which are called *Indecidua*, and the higher Placental animals, or *Deciduata*.

To the Indecidua, or lower Placental Animals, belong two very comprehensive and important vertebrate groups (1) the Hoofed Animals (*Ungulata*)—the Tapirs, Horses, Swine, Ruminants, and others; (2) the Whale-like animals (*Cetomorpha*)—the Sea-cows, Porpoises, Dolphins, Whales, and others. In all these Indecidua the chorion tufts are distributed, singly or in bunches, over the entire surface of the chorion, or over the greater part of it. They are but loosely attached to the mucous membrane of the uterus, so that the entire outer egg-membrane with its tufts may be easily and without using force be drawn out of the depression in the uterine mucous membrane, just as the hand is drawn from a glove. The two "vascular cakes" do not really coalesce at any point of their contact. Hence at birth the "embryonic cake" (*placenta fœtalis*) is removed; the "maternal cake" (*placenta uterina*) is displaced. The entire mucous membrane of the uterus is but little altered, and, at parturition, suffers no direct loss of substance.

The structure of the placenta in the second and higher division of Placental Animals, the Deciduata, is very different. To this comprehensive and very highly developed mammalian group belong all Beasts of Prey and all Insect-eaters, Gnawers (*Rodentia*), Elephants, Bats, Semi-apes, and lastly, Apes and Man. In all these Deciduata the entire surface of the chorion is also at first thickly covered with tufts. These, however, afterwards disappear from part of the surface, while they develop all the more vigorously on the remainder. The smooth chorion (*chorion læve*, Fig. 197, *chl*) thus becomes distinct from the tufted chorion (*chorion frondosum*, Fig. 198, *chf*). On the former there are

minute and scattered tufts, or none at all; while the latter is thickly overgrown with highly developed and large tufts. In the Deciduata the tufted chorion alone forms the placenta.

Yet more characteristic of the Deciduata is the very peculiar and intimate connection which is developed in these between the tufted chorion and the contiguous portion of the uterine mucous membrane, and which must be regarded as a true coalescence. The vascular tufts of the chorion push their branches into the sanguineous tissue of this mucous membrane in such a way, and the two sets of vessels are in such close contact and are so interlaced, that the embryonic placenta is no longer distinguishable from the maternal placenta; the two form one whole—a compact and apparently simple placenta. Owing to this intimate coalescence, a portion of the uterine mucous membrane of the mother comes away, at birth, with the firmly adherent egg-membrane. The portion of the mother's body which is thus removed in parturition is called, on account of its separable nature, the deciduous membrane (*decidua*). All Placental Animals which possess this deciduous membrane are classed together as Deciduata. The removal of this membrane at parturition, of course, causes a greater or less loss of blood by the mother, which does not occur in the Indecidua. In the Deciduata, moreover, the lost portion of the uterine mucous membrane must be replaced, after parturition, by a renewal of the tissue.

The structure of the placenta and deciduous membrane, however, by no means identical throughout the comprehensive group of Deciduata. On the contrary, there are many important differences in this respect, which are in

some degree connected with other important structural characters (*e.g.*, the structure of the brain, of the teeth, of the feet), and which may justly, therefore, be turned to account in the phylogenetic classification of Placentals. In the first place, two great groups of Deciduata may be distinguished according to the form of the placenta: in the one group it is ring-shaped or girdle-shaped; in the other it is discoid or cake-shaped. In Deciduata with girdle-shaped placenta (*Zonoplacentalia*) the poles of the oval egg take no part in the formation of the placenta. The "vascular cake" resembles a broad ring-like girdle, embracing the central zone of the egg. It is so in Beasts of Prey (*Carnassia*), both in the terrestrial forms (*Carnivora*) and in the marine forms (*Pinnipedia*). A similar girdle-shaped placenta is found in the False-hoofed Animals (*Chelophora*): the elephants, and Klip Das (*Hyrax*) with its allies, which were formerly classed as Hoofed Animals. All these *Zonoplacentalia* belong to one or more side-branches of the Deciduata which are not nearly allied to Man.

The second and most highly developed group is formed by the Deciduata with discoidal placenta (*Discoplacentalia*). The formation of the placenta is here most localized and its structure most fully developed. The placenta forms a thick, spongy cake, usually in the form of a circular or oval disc, and attached only to one side of the uterine wall. The greater part of the embryonic egg-membrane is, therefore, smooth, without developed tufts. To the *Discoplacentalia* belong the Semi-apes and Insect-eaters, the Diggers (*Effodienta*) and the Sloths, Rodents and Bats, Apes and Man. Comparative Anatomy enables us to infer that of these various orders the Semi-apes are the parent

group from which all other Discoplacentals, and perhaps even all Deciduous Animals, have developed as divergent branches. (Cf. Tables XXIII. and XXIV.)

The Semi-apes (*Prosimiæ*) are now represented only by very few forms. These, however, are very interesting, and must be regarded as the last remnants of a group once rich in forms. This group is certainly very ancient, and was probably very prominent during the Eocene Epoch. Their present degraded descendants are scattered widely over the southern portion of the Old World. Most of the species inhabit Madagascar; a few the Sunda Islands; a few others the continents of Asia and Africa. No living or fossil Semi-apes have, as yet, been found in Europe, America, or Australia.¹⁵⁰ The widely scattered posterity of the Semi-apes is considerably diversified. Some forms seem nearly allied to the Marsupials, especially to the Pouched-rats. Others (*Macrotarsi*) are very near akin to the Insect-eaters, and yet others (*Cheiromys*) to the Gnawers (*Rodentia*). One genus (*Galeopithecus*) forms a direct transition to the Bats. Finally, some of the Semi-apes (*Brachytarsi*) approach very near to true Apes. Among the latter are some tail-less forms (e.g., the Lori, *Stenops*, Fig. 199). From these highly interesting and important relations of the Semi-apes to the various Discoplacental orders, we may fairly infer that of the extant representatives of this group, they are the nearest to the common primitive parent-form. Among the direct common ancestors of Apes and Men, there must have been some Deciduata which we should class among the Semi-apes, were we to see them alive. We may therefore consider this order as a special stage, following the Pouched Animals, as the eighteenth stage in the human pedigree.

Probably our ancestors among the Semi-apes closely resembled the extant Brachytarsi or Lemurs (*Lemur*, *Lichas*).



FIG. 199.—The Slender Loris of Ceylon (*Stenops gracilis*).

otus, *Stenops*), and, like these, led a quiet life, climbing trees. The extant Semi-apes are mostly nocturnal animals of gentle and melancholy disposition, subsisting on fruits

Semi-apes are immediately followed by the true (*Simiæ*), as the nineteenth stage in the human pedigree. It has long been beyond doubt that of all animals these are in all respects the most nearly allied to Man. On the one side, the lowest Apes approach very near semi-apes, so, on the other side, do the highest Apes closely resemble Man. By carefully studying the Comparative Anatomy of Apes and Man, it is possible to trace a uninterrupted advance in the Ape-organization up to the fully human structure; and on impartially testing this question," which has lately been agitated with such intense interest, we shall infallibly have to acknowledge an important fact, which was first explicitly laid down by Huxley, that "whatever system of organs be studied, the sequence of their modifications in the ape series leads to the same result—that the structural differences which separate Man from the Gorilla and Chimpanzee are as great as those which separate the Gorilla from the lower Apes." In phylogenetic language this pregnant law is expressed in so masterly a manner by Huxley, is equivalent to the popular phrase: Man is descended from the Ape. In order to become convinced of the truth of this law, let us now once more consider the placenta and deciduous membrane, on the varied structure of which we have justly laid stress. Men and Apes, in the structure of their discoplacenta and in their decidua, do, indeed, coincide wholly with all other Discoplacental Animals. But in the more delicate structure of these parts Man is distinguished by peculiarities which he shares only with Apes, which are absent in other Deciduata. Thus in Man and the Apes three distinct parts are recognized in the

deciduous membrane; these parts may be called the outer, the inner, and the placental deciduous membrane. The outer or true membrane (*d. externa* or *vera*, Fig. 198, *e*, Fig. 200, *g*), is that portion of the uterine mucous membrane which coats the internal surface of the uterus wherever



FIG. 200.—Human embryo, twelve weeks old, with its coverings; natural size. The navel cord passes from the navel to the placenta: *b*, and *c*, chorion; *d*, placenta; *d'*, remains of tufts on the smooth chorion; *e*, *decidua reflexa* (inner); *g*, *decidua vera* (outer). (After Bernhard Schu

latter is not attached to the placenta. The placental spongy deciduous membrane (*d. placentalis* or *sera*, Fig. 198, *plu*, Fig. 200, *d*) is simply the maternal pla

self, or the maternal part of the "vascular cake" (*placenta uterina*), i.e., that part of the uterine mucous membrane which coalesces intimately with the chorion-tufts of



o. 201.—Mature human embryo (at the end of pregnancy), in its natural position, taken out of the uterus. On the inner surface of the latter (on the left) is the placenta, which is attached to the navel of the child by the umbilical cord. (After Bernhard Schultze.)

embryonic placenta (*placenta fœtal*). Lastly, the true or false deciduous membrane (*d. interna* or *reflexa*, 198, *dr*, Fig. 200, *f*) is that portion of the uterine mucous membrane which, as a peculiar thin envelope, covers all the surface of the egg-surface, lying immediately over the tuftless chorion (*chorion læve*). The origin of these three distinct deciduous membranes, concerning which erroneous notions have been entertained (still retained in the nomenclature), is plain enough; the external or true deciduous

membrane is a peculiar modification, afterwards lost, of the superficial layer of the original mucous membrane of the uterus. The placental membrane is that portion of the chorion preceding which is completely modified by the intrusion of the chorion-tufts and is employed in forming the placenta. Lastly, the inner deciduous membrane is formed by a ring-shaped fold of the mucous membrane (at the point of union of the *d. vera* and the *d. serotina*) which rings round the egg, and closes in the same way as the amnion.¹⁰⁰

The peculiar anatomical characters which mark the human egg-membrane re-occur, in the same form, only in Apes. In other Discoplacental Animals present greater or less differences in these points, the conditions being generally more simple. This is the case, for instance, in the structure of the placenta itself, in the coalescence of the chorion-tufts with the *decidua serotina*. The matured human placenta is a circular (rarely oval) disc of a soft, spongy character, 6 to 8 inches in diameter, about 1 inch thick, and weighs from 1 to 1½ lb. Its convex, external surface (that which coalesces with the uterus) is very uneven, and tufted. Its internal, concave surface (that which is turned towards the cavity of the egg) is quite smooth, and clothed by the amnion (Fig. 198, a). From near the centre of the placenta springs the navel cord (*funiculus umbilicalis*), the development of which we have already observed (vol. i. p. 383). It is coated by the amnion as with a sheath, which at the navel end passes directly into the abdominal skin (Fig. 200, 2). The mature navel cord is a cylindrical cord, coiled spirally around its axis, and usually about 20 inches long and ½ inch thick. It consists of gelatinous connective tissues ("W

's jelly"), in which are contained the remnants of the k-vessels and of the great navel vessels; the two navel arteries which convey the blood of the embryo to the placenta, and the great navel vein which brings back the blood in the latter to the heart. The numerous fine branches of these embryonic navel vessels pass into the branched arbor-tufts of the foetal placenta, and with these, finally, flow, in a very peculiar way, into large blood-filled cavities, which spread themselves in the uterine placenta and convey blood from the mother. The anatomical relations, very complex and difficult to comprehend, which are developed between the embryonic and the maternal placenta, exist in this form only in Man and in the higher Apes, while in all other Deciduous Animals their form is more or less different. The navel cord, also, is proportionately longer in Man and Apes than in other Mammals.

As in these important characters, so also in every other morphological respect, Man appears as a member of the order of Apes, and cannot be separated from the latter. The great originator of systematic description of nature, Karl Linnæus, with prophetic penetration, united Men, Apes, Mi-apes, and Bats in a single natural division, under the name of Primates, that is, the first, the lords of the animal kingdom. Later naturalists dissolved this order of Primates. The Göttingen anatomist, Blumenbach, first placed Man in a special order, which he called that of Two-handed Animals (*Bimana*); in a second order, he united Apes and Semi-apes under the name of Four-handed Animals (*Quadrumana*), while a third order included the distantly related Bats (*Chiroptera*). The separation of the *Bimana* and *Quadrumana* was retained by Cuvier and most succeeding

zoologists. It seems very important, but is really unjustifiable. This was first shown in the year 1863 by Huxley. Supported by very accurate Comparative Anatomical researches, he proved that Apes are as "two-handed" as Men, or, conversely, that Men are as "four-handed" Apes. Huxley showed, with convincing clearness, that the ideas previously held of the hand and the foot were and were incorrectly founded on physiological instead of morphological distinctions. The circumstance that in the hand, the thumb may be opposed to the other four fingers thus permitting the act of grasping, appeared especially to distinguish the hand from the foot, in which the corresponding great toe cannot be thus opposed to the four remaining toes. Apes, on the contrary, can grasp in this way with the hind-foot as well as with the fore-foot, and were therefore regarded as four-handed. Many tribes, however, among the lower races of men, especially many negro tribes, use the foot in the same way as the hand. In consequence of early and continued practice, they are able to grasp as well with the foot as with the hand (for example, in climbing or to grasp the branches of trees). Even new-born children of our own race have a very strong grasping power in the great toe with which they can hold a spoon as fast as with the thumb. The physiological distinction between hand and foot is therefore, neither be strictly carried out, nor scientifically established. Morphological characters must be used for the purpose.

A sharp morphological distinction of this kind—one founded on anatomical structure—between hand and foot, between the anterior and the posterior limbs, is as possible. There are essential and permanent differences

in the structure of the bony skeleton and in that of muscles which are attached to the hand and the foot; these are exactly the same in Man and in the Ape. There is, for instance, an essential difference in the arrangement and number of the wrist-bones of the hand (*carpus*) and ankle-bones of the foot (*tarsus*). The muscle-masses show equally constant differences. The posterior extremity, the foot, has always three muscles (a short flexor, a short extensor muscle, and a long muscle attached to the muscles of the tibia) which are never present in the anterior extremity, the hand. The disposition of the bones is also very different in the two sets of limbs. The characteristic differences between the anterior and posterior extremities occur in Man just as in Apes. It can, therefore, be no doubt, that the foot of the Ape serves the name as truly as that of the Man; and that true Apes are as genuinely two-handed animals as Man. Thus the usual distinction of the Apes from *Homo sapiens* is wholly unjustifiable.

It might now be asked whether, quite apart from these, there are not other marks by which Man is more widely separated from the Apes than are the different species of Apes from each other. Huxley has given a final negative answer to this question so convincingly, that the opposition now raised against him in many quarters must be regarded as wholly unfounded and ineffective. Based on an accurate knowledge of the Comparative Anatomy of all parts of the body, Huxley has brought forward very significant proof that, in anatomical respect, the differences between the highest and lowest Apes are greater than the corresponding differences between the highest Apes and Man. He there-

fore restored Linnæus's order of *Primates* (excluding the Bats), and divided it into three different sub-orders, the first of which is formed by the Semi-apes (*Lemurida*), the second by the true Apes (*Simiadae*), and the third by Man (*Anthropidae*).¹⁶¹

Yet, if we proceed logically and without prejudice, accordance with the principles of scientific reasoning, find, on the basis of Huxley's own law, this division adequate, and must go considerably further. As I showed in 1866, in treating this question in my *General Morphologie*, we are fully justified in taking at least one important step further, in assigning to Man his natural place in one of the divisions of the Ape-order. All characters distinctive of this one division of the Apes present in Man, while they are absent in other Apes. We are, therefore, not justified in forming a distinct order for Man apart from the true Apes.

The order of the true Apes (*Simiæ*), the Semi-apes being excluded, has long been divided into two natural groups, which, among other points, are distinguished by their geographical distribution. Those of one division (*Hesperopithecæ*, or Western Apes) live in the New World in America. The other division, to which Man belongs, that of the *Heopithecæ*, or Eastern Apes; these live in the Old World, in Asia, Africa, and, formerly, in Europe. In the Apes of the Old World, all *Heopithecæ*, share, in common with Man, all those characteristics to which special prominence is justly given, in distinguishing these two groups of Apes, in zoological classification; among these characteristics the structure of the teeth is most prominent. The objection is at once evident that the teeth are, in a physiolog-

much too subordinate a part of the body to justify so great a weight being attached to their structure in so important a question. There are, however, good reasons for prominent consideration of the structure of the teeth; it is with perfect correctness and propriety that systematic zoologists have, for more than a century, given great weight to this character in systematically describing and arranging the mammalian orders. The number, form, and disposition of the teeth are transmitted more accurately within the respective orders of the mammals than are most other zoological characteristics. The structure of the human teeth is well known. In maturity there are 32 teeth in our jaws, and of these 32 teeth, 8 front-teeth, 4 canine-teeth, and 20 molar-teeth. The 8 front-teeth or incisors (*dentes incisivi*), which are situated in the centre of the jaws, exhibit characteristic differences in the upper and lower jaw. In the upper the inner incisors are larger than the outer; in the lower jaw, the contrary, the inner incisors are smaller than the outer. Next to these, on each side, both in the upper and lower jaw, is a corner-tooth, which is larger than the inner, the so-called eye-tooth, or canine (*dens caninus*). Sometimes this tooth becomes very prominent in Men, as in Apes and many other Mammals, and forms a sort of fang. Finally, next to this, on each side, and in each jaw, are situated five back-teeth, or molar-teeth (*dentes molares*), of which the two foremost (the bicuspid teeth) are small, but a single fang, and are subject to the change of deciduous teeth, while the three hinder molars are much larger, have no fangs, and do not appear till after the temporary teeth have been shed (so-called "grinders"). The Apes of the

Old World have exactly this human structure of the teeth,—all Apes which have as yet been found, either living or as fossils, in Africa, Asia, and Europe. All Apes of the New World, on the contrary, all American Apes, have an extra tooth on both sides of each jaw; this is a bicuspid tooth. Thus they have six back-teeth on both sides of each jaw,—in all, thirty-six teeth. This characteristic difference between the Eastern and Western Apes has been so constantly transmitted within the two groups, that it is of the greatest value to us. A small family of South American Apes does, indeed, appear to form an exception in this respect. The pretty little Silk Apes, or Marmosets (*Hapalida*), namely, to which the Brush-monkey (*Midas*) and the tufted Marmoset (*Jacchus*) belong, have but five back-teeth in each half of the jaw, instead of six, and, accordingly, seem to approach nearer the Eastern Apes. But on closer observation it is found that, like all the Western Apes, they have the three bicuspid, and that the hindmost grinder has been lost. Thus this apparent exception confirms the value of the distinction.

Among the other marks by which the two main groups of the Apes are distinguished, the structure of the nose is specially important and prominent. In all Old World Apes the structure of the nose is the same as in Man; namely, a comparatively narrow partition of the two halves, so that the nostrils are directed downwards. In a few Eastern Apes, the nose projects as prominently and is as characteristically formed as in Man. We have already called attention, in this respect, to the remarkable Nose-ape (*Semnopithecus nasicus*), which has a well-curved and long nose (Fig. 202). Most of the Eastern Apes have, it is true, a

somewhat flatter nose, as, for instance, has the white-nosed Sea-cat (*Cercopithecus petaurista*, Fig. 203); yet in all the partition of the nose is narrow and thin. On the contrary, all American Apes have a different nasal structure. In them, the partition is peculiarly broadened and thickened below, and the wings of the nose are not developed, in consequence of which the nostrils are not below, but are turned outwards. This characteristic difference in the structure of the nose has also been so accurately trans-



FIG. 202.—Head of Nose-ape (*Semnopithecus nasicus*).

FIG. 203.—The white-nosed Sea-cat (*Cercopithecus petaurista*).

mitted in both groups, that, on account of it, the Apes of the New World have been called Flat-nosed (*Platyrrhinæ*), and those of the Old World Narrow-nosed (*Catarhinæ*). The former are, on the average, inferior in organization.

The division of the order of Apes into two sub-orders, the *Platyrrhinæ* and the *Catarrhinæ*, is, on account of the constant hereditary characters, now generally accepted by zoologists, and receives much support from the geographical distribution of the two groups between the New and Old Worlds. From this follows the direct inference, very important in its bearing on the Phylogeny of Apes, that, from the primæval common parent-form of the Ape-order, two diverging lines branched out at a very early period, one of which spread over the New World, the other over the Old. It is certain that all the Flat-nosed Apes, on the one hand, are descendants of a common parent-form, and, on the other hand, all the Narrow-nosed Apes from another.

An inference concerning our own pedigree may be drawn from this. Man has exactly the same characters, the same peculiar formation of the teeth and nose, as all the *Catarrhinæ*, and is as thoroughly distinguished by these characteristics from the *Platyrrhinæ*. We are therefore compelled, in classifying the Primates, to assign to Man a place in the Narrow-nosed group. The bearing of this on our tribal history is, that Man is immediately related in blood to the apes of the Old World, and may be traced from a parent-form common to all other *Catarrhinæ* also. Man is a genuine Narrow-nosed Ape in his whole structure and in origin, and has descended from some unknown, extinct *Catarrhine* form in the Old World. On the other hand, the Apes of the New World, the Flat-nosed group, constitute a diverging branch of our family tree, and stand in no near genealogical relation to the human race.

We have now reduced the circle of our nearest allies to the small group, containing comparatively few forms,

h is represented by the sub-order of the Narrow-nosed, Eastern Apes. Finally, the question which now remains to be answered is—what position in this sub-order is to be assigned to Man, and whether other inferences as to the structure of our immediate ancestors may be drawn from this position. The comprehensive and acute researches of the Comparative Anatomy of Man and the various Primates, which Huxley has recorded in his work on the "Evidence as to Man's Place in Nature," are of the greatest value in furnishing the answer to these important questions. The inevitable conclusion is, that the difference between Man and the highest Narrow-nosed Apes (the Gorilla, Chimpanzee, Orang) is slighter in every respect than the corresponding differences between the highest and the lowest Primates (the Sea-cat, Macaque, Baboon). Even within the small group of the Tail-less man-like Apes (*Anthropoidea*) the several genera do not differ less from each other than they do from Men. This is seen by a glance at the skeletons represented here, as arranged by Huxley (Figs. 208). If the skull, or the vertebral column, together with the rib-system, or the anterior or posterior members, be compared; or if the comparison is extended to the muscular system, the circulatory system, the brain, etc., an impartial and unprejudiced examination always results in the same conclusion, that Man does not differ more from the higher Catarrhines than the extreme forms of the latter (for example, the Gorilla and Baboon) differ from each other. We can, therefore, complete the important proposition already quoted from Huxley: We may take whatever system of organs we will,—the comparison of their variations within the ranks of the Catarrhines leads us



FIG. 204.
(*Cf.* p. 181.)
Gibbon.

FIG. 205.
(Plate XIV Fig. 3.)
Orang-outang.

FIG. 206.
(Plate XIV Fig. 1.)
Chimpanzee.

FIG. 207.
(Plate XIV Fig. 2.)
Gorilla.

FIG. 208.
(Plate XIV Fig. 4.)
Man.

to one and the same conclusion: that the anatomical differences that distinguish Man from the most highly developed Catarhinæ (the Orang, Gorilla, Chimpanzee), are not so great as those which separate the latter from the lowest Catarhinæ (Sea-cat, Macaque, Baboon).

We must, therefore, consider the proof complete, that Man is descended from other Narrow-nosed Apes (*Catarhinæ*). Although future researches into the Comparative Anatomy and Ontogeny of the existing Catarhines, as well as of their fossil relatives, promise us various new details, yet no future discovery can ever overthrow that important proposition. Our Catarhine ancestors must, of course, have passed through a long series of varied forms, before Man finally developed as the most perfect form. The following must be considered as the most important advances by which this "Creation of Man," his differentiation from the most nearly allied Catarhine Apes, was effected: Habituation to upright carriage and, in connection with this, the greater differentiation of the anterior and posterior limbs; also, the development of articulate speech and its organ, the larynx; and lastly, and especially, the more perfect development of the brain and its function, the soul; sexual selection must have exerted an extraordinarily important influence, as Darwin has conclusively proved in his celebrated work on sexual selection.¹⁶³

With reference to these advances, we may, among our Catarhine ancestors, distinguish at least four important ancestral stages, marking prominent epochs in the great historical process of the origin of Man. As the nineteenth stage in the human pedigree, next to the Semi-apes, we may place the oldest and lowest Catarhine Apes, which developed

from the former by the formation of the characteristic catarhine head, and by the peculiar modification of the teeth, the nose, and the brain. This oldest parent-form of the whole Catarhine group must, certainly, have been thickly covered with hair, and must have had a long tail; was, in fact, a Tailed Ape (*Menocerca*, Fig. 203). They were already in existence during the earlier part of the Tertiary Epoch (during the Eocene Period), as is shown by fossil remains of Eocene Catarhines. Among extant Tailed Apes, the Slender Apes (*Semnopithec*i) are perhaps most nearly related to this parent-form.¹⁶⁸

As the twentieth stage in the human pedigree, next to these Tailed Apes, we must rank the Tail-less man-like Apes (*Anthropoides*), under which name the most highly developed Catarhines, those most nearly related to Man, have been grouped. They originated from the Tailed Catarhines, by the loss of the tail, the partial loss of their hairy covering, and the further development of the brain, the latter being indicated in the preponderating development of the brain-skull over the facial skull. At the present time but few forms of this remarkable family are in existence; they are distributed into two different groups, an African and an Asiatic group. The African Man-like Apes are limited to the western part of tropical Africa, but are probably distributed over Central Africa in several species. Only two species are well known: the Gorilla (*Pongo gorilla*, or *Gorilla engina*), the largest of all Apes (Fig. 207); and the smaller Chimpanzee (*Pongo troglodytes*, or *Engeco troglodytes*), which may be seen in several zoological gardens (Figs. 206, Plate XIV. Figs. 1, 2). Both the African Man-like Apes are black in colour, and like their countrymen,

11

1. Chimpanzee.

2. Gorilla.



3. Orang.

4. Negro.

the Negroes, have the head long from back to front (dolichocephalic). The Asiatic Man-like Apes are, on the contrary, mostly of a brown, or yellowish brown colour, and have the head short from back to front (brachycephalic), like their countrymen, the Malays and Mongols. The largest Asiatic Man-like Ape is the well-known Orang, or Orang-outang (Fig. 128), which is indigenous in the Sunda Islands (Borneo, Sumatra), and is brown in colour. Two species have recently been distinguished: the great Orang (*Satyrus Orang*; Fig. 205, Plate XIV. Fig. 3), and the small Orang (*Satyrus morio*). A genus of smaller Anthropoids (Fig. 204), the Gibbons (*Hylobates*), live on the main-land of Southern Asia and on the Sunda Islands; from four to eight different species of these have been distinguished. Neither of these living Anthropoids can be indicated as the Ape absolutely most like Man. The Gorilla approaches nearest to Man in the structure of the hand and foot, the Chimpanzee in important structural details in the skull, the Orang in the development of the brain, and the Gibbon in that of the thorax. It is evident that no single one of these existing Man-like Apes is among the direct ancestors of the human race; they are all the last scattered remnants of an old, catarrhine branch, once numerous, from which the human race has developed as a special branch and in a special direction.

Although Man (*Homo*) ranks immediately next to this anthropoid family, from which he doubtless directly originated, yet the Ape-men (*Pithecanthropi*) may be inserted here, as an important intermediate form between the two, and as the twenty-first stage in our ancestral series. In the "Natural History of Creation" (vol. ii. p. 293), I have

applied this name to the speechless Primitive Men (*Alali*) who made their appearance in what is usually called the human form, that is, having the general structure of Man (especially in the differentiation of the limbs)—but being destitute of one of the most important qualities of Man, namely, articulate speech, as well as of the high mental development connected with speech. The high differentiation of the larynx and of the brain occasioned the latter, first gave rise to the true “Man.”

Comparative Philology has recently shown that the present human language is polyphyletic in origin, that several, and probably many, different original languages must be recognized, as having developed independently from each other. The history of the development of language also teaches us (its Ontogeny in every child, as well as its Phylogeny in every race), that the actual rational language of men developed gradually, only after the body had developed into the specific human form. It is even probable that the formation of language did not begin until after the differentiation of the various species, or races of men, and this presumably occurred in the beginning of the Quaternary Epoch, or the Diluvial Period. The Ape-men or *Alali*, were therefore probably already in existence toward the close of the Tertiary Epoch, during the Pliocene Period, perhaps even as early as the Miocene Period.¹⁶⁴

Lastly, the genuine or speaking human being (*Homo*) must be considered as the twenty-second and final stage in our animal pedigree. Man originated from the preceding stage in consequence of the gradual improvement of inarticulate animal sounds into true human articulate speech. Only very uncertain conjectures can be formed

the time and place of this true "Creation of Man." is probable that Primæval Man originated during the Tertiary Epoch, in the torrid zone of the Old World, either the continent of tropical Asia or Africa, or on an earlier continent which has now sunk below the surface of the Indian Ocean, and which extended from Eastern Africa (Madagascar and Abyssinia) to Eastern Asia (the Sunda Islands and Eastern India). In my "Natural History of Creation" Chapter XXIII. and Table XV.), I have already fully discussed the important evidence as to the former existence of this large continent, called Lemuria, and how the distribution of the various species and races of men probably took place from this "Paradise" over the surface of the earth. In the same place, I have also fully discussed the interrelations of the various races and species of the human race.¹⁶³



TABLE XXII.

SYSTEMATIC SURVEY OF THE PERIODS IN THE TRIBAL HISTORY OF HUMAN RACE.

(Compare Table VIII., vol. i. p. 402.)

FIRST MAIN PERIOD IN TRIBAL HISTORY.

The Plastid Ancestors of Man.

The form of the ancestors of man is equal to the simple individual of first order, a single plastid.

First Stage : **Moneron Series** (Fig. 163, p. 46).

The ancestors of man are single, living, simple cytoda.

Second Stage : **Amœba Series** (Fig. 167 p. 58).

The ancestors of man are single, living, simple cells.

SECOND MAIN PERIOD IN TRIBAL HISTORY.

The many-celled Primitive Animal Ancestors of Man.

The ancestors of man consist of a closely-united society of many homogeneous cells; hence their form-value is that of individuals of the second order, of *Idorgana*.

Third Stage : **Synamœba Series** (Fig. 170, p. 55).


The ancestors of man are many-celled primitive animals of the simplest kind: solid masses of simple, homogeneous cells.

Fourth Stage : **Planœa Series** (Figs. 172, 173, p. 60).

The ancestors of man are many-celled primitive animals of a character like that of the *Magosphœra* and certain planula-larvæ, of equal rank to the ontogenetic *Blastula* or *Blastosphœra*; hollow spheres, the wall of which consists of a single stratum of ciliated cells.



PEDIGREE OF MAN.





THIRD MAIN PERIOD IN TRIBAL HISTORY.

The Invertebrate Intestinal Animal Ancestors of Man.

The ancestors of man have the form-value of individuals of the third order, of inarticulate individuals. The body encloses an intestinal cavity with a mouth, and consists at first of two primary germ-layers, afterwards of four secondary germ-layers.

Fifth Stage : *Gastræa Series* (Figs. 174-179, p. 65).

The ancestors of man have the form-value and structure of a *Gastrula*. The body consists merely of a simple primitive intestine, the wall of which is formed of the two primary germ-layers.

Sixth Stage : *Chordonium Series* (Figs. 184-188, p. 80-90).

The ancestors of man are worms : at first, primitive worms, allied to the *Turbellaria*; afterwards worms of higher rank, *Scolecida*; finally, notochord-animals with the organization of the ascidian larvæ. The body is composed of four secondary germ-layers.

FOURTH MAIN PERIOD IN TRIBAL HISTORY.

The Vertebrate Ancestors of Man.

The ancestors of man are vertebrates, and their form-value is, therefore, that of an articulated individual, or a chain of metamera. The skin-sensory layer is specialized into the horn-plate, medullary tube, and primitive kidneys. The skin-fibrous layer has divided into the leather-plate, primitive vertebræ (muscular plate and skeleton-plate), and the notochord. From the intestinal-fibrous layer originates the heart with the main blood-vessels and the fleshy intestinal wall. From the intestinal-glandular layer, the epithelium of the intestinal tube is formed. The formation of metamera is constant.

Seventh Stage : *Acrania Series* (Fig. 189; Pl. XI. Fig. 15).

The ancestors of man are skull-less vertebrates, like the extant *Amphioxus*. The body already forms a chain of metamera, several primitive vertebræ having separated off. The head is not yet entirely distinct from the trunk. The medullary tube has not separated into brain-bladders. The heart is very simple, without chambers. The skull is still wanting; as are also the jaws and limbs.

Eighth Stage: Monorhina Series (Fig. 190; Pl. XI. Fig. 16).

The ancestors of man are jaw-less skulled animals (resembling developed *Myxinoidea* and *Petromysontes*). The number of the metameres increasing. The head is becoming more distinctly differentiated from trunk. The anterior end of the medullary tube swells into a bladder-structure and forms the brain, which is soon differentiated into five or six bladders. At the sides of these appear the three higher organs of sense. The heart is divided into auricle and ventricle. The jaws, limbs, swimming-bladder are still wanting.

Ninth Stage: Ichthyoda Series (Figs. 191, 192; Pl. XII. and XIII.).

The ancestors of man are fish-like skulled animals: first, *Primitive Fishes (Selachii)*, then *mud-fishes (Dipneusta)*, then gilled *Batrachians (Serranidae)*. The ancestors belonging to this Ichthyoda stage develop two pairs of limbs: a pair of anterior limbs (pectoral fins) and a pair of posterior limbs (ventral fins). The gill-arches are formed between the gill-openings, and from them are formed the first pair of jaw-arches (upper and lower jaws). The swimming-bladder (lungs), liver, and pancreas grow from the intestinal canal.

Tenth Stage: Amniota Series (Figs. 195-208; Pl. XIV.).


The ancestors of man are amnion-animals or gill-less vertebrates: first, *Primitive amniota (Protamnion)*, then *Primitive mammals (Monotremata)*; then *Pouched animals (Marsupialia)*; then *Semi-apes (Prosimia)*, and, last, *Apes (Simia)*. The ape-ancestors of man are first tailed *Catarrhini*, then tail-less *Catarrhini (Anthropoidea)*, then speechless Ape-men (*Alali*), and last genuine, speaking men. The ancestors belonging to this amniota series develop an amnion and allantois, and gradually acquire the mammalian structure, and at last the specific human form.

TABLE XXIII.

Systematic Survey of the Phylogenetic Classification of Mammals.

I. First -class of animals	Cloacal Animals (<i>Monotrema</i> , or <i>Ornithodelphia</i>)	1. Primitive Mammals	<i>Promammalia</i>
		2. Beaked Animals	<i>Ornithostoma</i>
II. Second -class of animals	Pouched Animals (<i>Marsupialia</i> , or <i>Didelphia</i>)	3. Herbivorous Pouched Animals	<i>Notanophaga</i>
		4. Carnivorous Pouched Animals	<i>Zoophaga</i>
III. Third class of animals: Placental animals (<i>Eutentalia</i> , <i>Monotremalia</i> , <i>Monotremalia</i>)	III. (a) Placental Mammals with- out Decidua, with Tufted Placenta <i>Indecidua</i> <i>Villioplacentalia</i>	5. Hoofed Animals { <i>Ungulata</i> { Single-hoofed Double-hoofed	<i>Perissodactyla</i> <i>Artiodactyla</i>
		6. Whale-like Animals { <i>Cetomorpha</i> { Sea-cows Whales	<i>Sirenia</i> <i>Cetacea</i>
	III. (b) Placental Mammals with Decidua, with Girdle Placenta <i>Deciduata</i> <i>Zonoplacentalia</i>	7. Pseudo-hoofed Animals { <i>Chelophora</i> { Rock Conies Elephants	<i>Lamnungia</i> <i>Proboscidea</i>
		8. Beasts of Prey { <i>Carnassia</i> { Land Beasts of prey Marine Beasts of prey	<i>Carnivora</i> <i>Pinnipedia</i>
	III. (c) Placental Mammals with Decidua, with Discoid Placenta <i>Deciduata</i> <i>Discoplacentalia</i>	9. Semi-apes <i>Prosimia</i> { Fingered animals Long-footed Flying Lemur Lemurs	<i>Leptodactyla</i> <i>Macrotarsi</i> <i>Ptenopleura</i> <i>Brachytarsi</i>
		10. Gnawing Ani- mals <i>Rodentia</i> { Squirrel species Mouse species Porcupine species Hare species	<i>Sciuromorpha</i> <i>Myomorpha</i> <i>Hystriehomorpha</i> <i>Lagomorpha</i>
		11. Toothless <i>Edentata</i> { Digging animals Sloths	<i>Effodientia</i> <i>Bradypoda</i>
		12. Insect-eaters <i>Insectivora</i> { With Cæcum Without Cæcum	<i>Menotyphla</i> <i>Lipotyphla</i>
		13. Flying Animals <i>Chiroptera</i> { Flying Foxes Bats	<i>Pterocynetes</i> <i>Nycterides</i>
		14. Apes <i>Simia</i> { Flat-nosed Narrow-nosed Apes	<i>Platyrrhina</i> <i>Catarrhina</i>


TABLE XXIV.

Pedigree of Mammals.

(189)

TABLE XXV.

Pedigree of Apes.



CHAPTER XX.

THE HISTORY OF THE EVOLUTION OF THE EPIDERMIS AND THE NERVOUS SYSTEM.

Animal and Vegetative Organ-systems.—Original Relations of these to the Two Primary Germ-layers.—Sensory Apparatus.—Constituents of Sensory Apparatus: originally only the Exoderm, or Skin-layer; afterwards, the Skin-covering specialized from the Nerve-system.—Double Function of the Skin (as a Covering and as Organ of Touch).—Outer Skin (*Epidermis*) and Leather-skin (*Corium*).—Appendages of the Epidermis: Skin-glands (Sweat-glands, Tear-glands, Sebaceous Glands, Milk-glands); Nails and Hair.—The Embryonic Wool-covering.—Hair of the Head and of the Beard.—Influence of Sexual Selection.—Arrangement of the Nerve-system.—Motor and Sensory Nerves.—Central Marrow: Brain and Dorsal Marrow.—Constitution of the Human Brain: Large Brain (*Cerebrum*) and Small Brain (*Cerebellum*).—Comparative Anatomy of the Central Marrow.—Germ-history of the Medullary-tube.—Separation of the Medullary-tube into Brain and Dorsal Marrow.—Modification of the Simple Brain-bladder into Five Consecutive Brain-bladders: Fore-brain (Large Brain, or *Cerebrum*), Twixt-brain ("Centre of Sight"), Mid-brain ("Four Bulbs"), Hind-brain (Small Brain, or *Cerebellum*), After-brain (Neck Medulla).—Various Formation of the Five Brain-bladders in the various Vertebrate Classes.—Development of the Conductive Marrow, or "Peripheral Nervous System."

"Hardly any part of the bodily frame, then, could be found better calculated to illustrate the truth that the structural differences between Man and the highest Ape are of less value than those between the highest

the lower Apes, than the hand or the foot, and yet, perhaps, there is one which enforces the same conclusion in a still more striking manner—that is the brain."—*Man's Place in Nature*, p. 94 (1863).

As if to demonstrate, by a striking example, the impossibility of any cerebral barrier between Man and the Apes, Nature has led us, in the latter animals, with an almost complete series of gradations, from brains little higher than that of a Rodent to brains little lower than that of Man."—*Ibid.* p. 96.

investigations, up to the present, have shown us how the whole human body has developed from an entirely simple beginning, from a single simple cell. The whole human body, as well as the individual man, owes its origin to a single-celled cell. The one-celled parent-form of the former is, even reproduced in the one-celled germ-form of the latter. In conclusion, we must glance at the evolutionary history of the separate parts which constitute the human body. In this matter, I must, of course, restrict myself to the most general and important outlines; for a detailed study of the evolutionary history of the separate organs and tissues would occupy too much space, and would demand a greater amount of anatomical knowledge than the generality of my readers are likely to possess. In considering the development of the organs, and of their functions, we will retain the method previously employed, except that we will consider the germ-history and the tribal history of the various parts of the body in common. In the history of the evolution of the human body as a whole we have found that Phylogeny everywhere serves to throw light on the obscure course of ontogeny, and that the clew afforded by phylogenetic continuity alone enables us to find our way through the labyrinth of ontogenetic facts. We shall experience exactly the same in the history of the development of the separate

organs, but I shall be compelled to explain the ontogeny and the phylogenetic origin of the organs simultaneously. For the further we penetrate into the details of organ development, and the more minutely we study the organs of the separate parts, the more clearly do we see inseparably the evolution of the germ is connected with that of the tribe. The Ontogeny of the organs is intelligible and explicable only through their Phylogeny; just as the germ-history of the entire body (the "person") is intelligible only by the history of the tribe. Each form is determined by a corresponding ancestral form. It is as true of the parts as of the whole.

In endeavouring, with the help of this fundamental principle of Biogeny, to obtain a general view of the main features of the development of the separate organs of man, we must, in the first place, consider the animal, and then the vegetable organ-systems of the body. The first main group of the animal organ-systems, is formed by the sensory apparatus together with the motor apparatus. To the former belong the skin-covering, the nervous system, and the organs of senses. The motor apparatus consists of the passive apparatus of movement (the skeleton) and the active organ-systems (muscles). The second main group of organs, the vegetable organ-system, is formed by the nutritive and the excretory apparatus. To the nutritive apparatus belong especially the intestinal canal with all its appendages together with the vascular and renal systems. The excretory apparatus includes the various sexual organ-systems, germ-glands, germ-ducts, organs of copulation, etc.).

In earlier chapters (IX. and X.) it has been stated that the animal organ-systems (the instruments of sensation)

movement) proceed especially from the outer primary layer, from the skin-layer. The vegetative organs, on the other hand (the instruments of nutrition and excretion), proceed principally from the inner primary layer, from the intestinal layer. This radical contrast between the animal and the vegetative spheres of the body is true, by no means absolute either in man or in the lower animals; on the contrary, many separate parts of the animal apparatus (*e.g.*, the intestinal nerve, or sympathetic) are derived from cells which have proceeded from the entoderm; and, on the other hand, a large part of the vegetative apparatus (*e.g.*, the mouth-cavity, and probably the greater part of the urinary and sexual organs) is formed of cells which are originally derived from the exoderm. Moreover, in the bodies of all the more highly developed animals, the heterogeneous parts are so intermixed and blended that it is often extremely difficult to assign its true source to one of the constituent parts. But, on the whole, we may assume as a certain and important fact, that in Man, and in all high animals, the greater part of the animal organs may be referred to the skin-layer, or exoderm; the greater part of the vegetative organs to the intestinal layer, or entoderm. For this reason, Baer called the former the animal germ-layer, the latter, the vegetative germ-layer (op. cit. pp. 53 and 196). Of course, in making this important distinction, we pre-suppose the correctness of Baer's view, according to which the skin-fibrous layer (the "flesh layer" of Baer) must have originated (phylogenetically) from the exoderm, and, on the other hand, the intestinal layer (Baer's "vascular layer") from the entoderm. This influential view, which is yet much disputed, is, we

TABLE XXVI.

Systematic Survey of the Organ-Systems of the Human Body.

(N.B.—The origin of the separate organs from the four secondary layers is indicated by the Roman numerals (I–IV.): I. Skin-sensory II. Skin-fibrous layer; III. Intestinal-fibrous layer; IV. Intestinal ular layer.)

ANIMAL ORGAN-SYSTEM.	A. Sensory Apparatus <i>Sensorium</i>	1. Skin-covering (<i>Derma</i>)	{ Outer skin Leather skin	Epidermis, I. Corium, II.
		2. Central nerve- system	{ Brain Spinal marrow	Encephalon Medulla spinalis
		3. Peripheric nerve- system	{ Brain nerves Spinal nerves Intestinal nerves	Nervi cerebralis Nervi spinales Sympatheticus
		4. Sense-organs (<i>Organa sensuum</i>)	{ Organ of touch (skin) Organ of taste (tongue) Organ of smell (nose) Organ of sight (eye) Organ of hearing (ear)	Org. tactus Org. gustus Org. olfactus Org. visus Org. auditus
	B. Motive Apparatus <i>Locomotorium</i>	5. Muscle - system (active motive organs)	{ Skin muscles Skeleton muscles	Musculi cutanei M. skeletici
		6. Skeleton-system (passive motive organs)	{ Vertebral column Skull Limb skeleton	Vertebrarium Cranium Sk. extremitatum
VEGETATIVE ORGAN-SYSTEM.	C. Nutritive Apparatus <i>Nutritorium</i>	7. Intestinal system (<i>Gaster</i>)	{ Digestive organ Respiratory organ	O. digestiva O. respiratoria
		8. Vascular system (<i>Organa circula- tionis</i>)	{ Body cavity Lymph vessels Blood vessels Heart	Cœloma, II. Vasa lymphatica V. sanguiferum Cor, III.
		9. Renal system (<i>Organa urinaria</i>)	{ Kidneys Urinary ducts Urinary bladder	Renes } I. Ureteres } Urocystis, III
	D. Reproductive Apparatus <i>Propagatorium</i>	10. Sexual organs (<i>Organa sexualia</i>)	{ Sexual glands (I. Ovary) (II. Testes) Sexual ducts (I. Oviduct) (II. Seed duct) Copulatory organs (I. Sheath) (II. Penis)	Gonades (I. Ovaria) III (II. Testes) I. Gonophori (I. Oviduc- tus) (II. Sperma- ductus) Copulativa (I. Vagina) (II. Penis)

think, securely founded on the Gastrula—that most important of all the germ-forms of the animal kingdom—which we find recurs in similar form in the germ-history of the most different classes of animals. This significant germ-form points unmistakably to a parent-form (the Gastræa) common to all animals, the Protozoa alone excepted; in this long extinct parent-form the entire body of the animal consisted throughout life of the two primary germ-layers, as is yet the case, for a short time, in the Gastrula. In the Gastræa the simple skin-layer did *actually* represent all the animal organs and functions, and the simple intestinal layer, on the other hand, all the vegetative organs and functions; potentially, this is even yet the case in the Gastrula.

In studying the development of the first important part of the animal sphere, the sensory apparatus, or sensorium, we shall now find how well adapted this Gastræa theory is to explain, not only in a morphological but in a physiological sense, the most important facts in the history of evolution. This sensory apparatus consists of two very distinct parts, having, apparently, nothing in common: in the first place, the external skin-covering (*Derma*), together with its appendages, the hair, nails, sweat-glands, &c.; and, secondly, the nervous system, situated internally. The latter includes the central nervous system (brain and spinal chord), the peripheric brain-nerves and medullary nerves, and finally, the organs of sense. In the fully developed vertebrate body these two main constituents of the sensorium are entirely separate; the skin lying entirely externally on the body, while the central nervous system is within, and quite separate from the former. The two are connected merely by a portion of the peripheric nerve-

system and of the sense-organs. And yet, as we all know from the germ-history of man, the latter is developed from the former. Those organs of our body which discharge the highest and most perfect functions of animal life—of sensation, volition, thought—in a word, the organs of the psyche, of mental life—arise from the external covering.

This remarkable fact, considered in itself alone, seems wonderful, inexplicable, and paradoxical, that the truth of the fact was simply long denied. The most trustworthy embryological observations were met with the erroneous statement that the central nerve-system develops, not from the germ-layer, but from a special cell-layer lying under this. The ontogenetic fact would not, however, yield now that Phylogeny has thrown light on the subject; the fact seems perfectly natural and necessary. When we reflect on the historic evolution of mind and sense activities we must necessarily conceive the cells, which according to these, as originally situated on the outer surface of the animal-body. Such externally placed elementary cells could alone directly receive and deal with impressions from the outer world. Afterwards, under the influence of natural selection, the complex cell-masses which had been especially "sensitive" gradually withdrew into the shelter of the interior of the body, and there laid the first foundation of a central nervous organ. As differentiation advanced the distance and distinction between the external covering and the central nervous system detached from it became continually greater, and finally the two were permanently connected merely by the conductive peripheral nerves.

This view is fully confirmed by the results of Comparative Anatomy. Comparative Anatomy shows that many lower animals possess no nervous system, although, in common with higher animals, they exercise the functions of sensation, volition, and thought. In the Primitive Animals (*Protozoa*), which do not even form germ-layers, of course the nervous system, like the skin-covering, is wanting. Even in the second main division of the animal kingdom—in the Metazoa or Intestinal Animals—there is at first no nervous system. The functions of these are performed by the simple cell-layer of the exoderm, which the lower Intestinal Animals have inherited directly from the *Gastræa* (Fig. 209, *e*). This is the case in the lowest Plant Animals (*Zoophyta*), the *Gastræads*, Sponges, and the lowest Hydroid Polyps, which are but little higher than the *Gastræads*. Just as all the vegetative functions of these are performed by the simple intestinal layer, so all the animal functions are discharged by the equally simple skin-layer. The simple cell stratum of the exoderm is, in these, skin-covering, motive apparatus, and nervous system simultaneously.

Most probably the nervous system was also wanting in a large proportion of those Primitive Worms (*Archelminthes*) which were developed directly from the *Gastræads*. Even those Primitive Worms in which the two primary germ-layers had already split into the four secondary germ-layers (Plate V. Fig. 10), seem not to have possessed a nervous system distinct from the skin. The skin-sensory layer must, even in these long-extinct Worms, have been at once skin-covering and nerve-system. But already in the Flat Worms (*Platelmintes*), and especially in the Gliding Worms (*Turbellaria*) which of all existing forms approach nearest to the Primitive

Worms, we find an independent nerve-system, distinct and separate from the outer skin-covering. This is the "upper






FIG. 209.—Gastrula of *Gastrophysa* (Gastracae-class).

FIG. 210.—Transverse section through an embryonic Earth-worm : skin-sensory layer; *hm*, skin-fibrous layer; *df*, intestinal-fibrous layer; intestinal-glandular layer; *a*, intestinal cavity; *c*, body-cavity, or *Coelom*; *n*, nerve-ganglia; *u*, primitive kidneys.

FIG. 211.—A Gliding Worm (*Rhabdocœlum*). From the brain or upper throat ganglion (*g*) nerves (*n*) radiate towards the skin (*f*), the eyes (*e*) the organ of smell (*na*), and the mouth (*m*) : *h*, testes; *c*, ovaries.

throat ganglion," situated above the throat (Fig. 211, *g*; Plate V. Fig. 11, *m*). The complex central nervous system of all higher animals has developed from this simple rudiment. In the higher Worms, *e.g.*, the Earth-worms, according to Kowalevsky, the earliest rudiment of the central nervous system (Fig. 210, *n*) is a local thickening of the skin-sensory layer (*hs*), which afterwards, becomes entirely detached from the horn-plate. Even the medullary tube of Vertebrates has the same origin. From the germ-history of Man, we already know that this medullary tube, the commencement of the central nervous system, originally develops from the outer skin-covering.

Let us now turn aside from these very interesting features in evolution, and examine the development of the later human skin-covering, with its hairs, sweat-glands, etc. Physiologically, this outer covering (*derma*, or *tegumentum*) plays a double part. The skin, in the first place, forms the general protective covering (*integumentum commune*) which covers the whole surface of the body, and protects all other parts. As such, it, at the same time, effects a certain exchange of matter between the body and the surrounding atmospheric air (perspiration or skin-breathing). In the second place, the skin is the oldest and most primitive sense-organ, the organ of touch, which effects the sensation of the surrounding temperature and of the pressure or resistance of bodies with which it comes in contact.

The human skin, like that of all higher animals, consists essentially of two distinct parts; of the outer-skin, and of the underlying leather-skin. The outer-skin (*epidermis*) consists only of simple cells, and contains no blood-vessels (Fig. 212, *ab*). It develops from the first of the secondary

germ-layers from the skin-sensory layer, and, directly, from the horn-plate of the latter. The leather-skin (*corium*), on the contrary, consists principally of connective or fibrous



FIG. 212.--Human skin in perpendicular section (after Ecker), much enlarged: *a*, horny stratum of outer-skin (*epidermis*); *b*, mucous stratum of outer-skin; *c*, papillæ of the leather-skin (*corium*); *d*, blood-vessels of the latter; *e*, *f*, excretory ducts of the sweat-glands (*g*); *h*, fat-globules of the leather-skin; *i*, nerve, passing above into a touch-body.

tissue, contains numerous blood-vessels and nerves, and has a different origin. It develops from the outer stratum of the second secondary germ-layer, from the skin-fibrous layer. The leather-skin is much thicker than the outer-skin. In its deeper part, the "*subcutis*," lie many masses of fat-cells (Fig. 212, *h*). Its upper part, the true "*cutis*," or papillary layer, forms, over nearly the whole surface of the body, a number of microscopic cone-shaped warts, or papillæ, which fit into the overlying epidermis (*c*). These touch-warts, or sensory papillæ, contain the most delicate of all the sensory organs of the skin, the "*corpuscula tactus*." Other papillæ

contain merely the terminal loops of the nutritive blood-vessels of the skin (*cd*). All these different parts of the leather-skin originate, by differentiation, from the cells, originally homogeneous, of the leather-plate, the outer lamella of the skin-fibrous layer (Fig. 112, *hpr*, vol. i. p. 352; Plates IV. and V., *l*; Figs. 65-69, *hf*, p. 277).¹⁶⁶

Analogously, all the constituent parts and appendages of the outer-skin (*epidermis*) originate, by differentiation, from the homogeneous cells of the horn-plate (Fig. 213). At a

FIG. 213.—Cells of the outer-skin (*epidermis*) of a human embryo of two months. (After Koelliker.)



very early period, the simple cell-layer of this horn-plate splits into two distinct strata. The inner, softer stratum (Fig. 212, *b*) is called the mucous layer; the outer, harder stratum (*a*), the horn-layer of the outer-skin. The surface of this horn-layer is continually worn out and thrown off; new cell-strata, produced by the growth of the underlying mucous layer, take its place. Originally the outer-skin forms an entirely simple cover over the surface of the body. Afterwards, however, sundry appendages develop from this both internally and externally. The internal appendages are the skin-glands; the sweat-glands, the sebaceous glands, etc. The external appendages are hair, nails, etc.

The glands of the skin-covering are at first merely solid plug-shaped growths of the outer-skin (*epidermis*), which penetrate into the underlying leather-skin (*corium*) (Fig. 214₁). A canal afterwards forms inside these solid

plugs (2, 3), either owing to the softening and breaking up of the central cells, or as the result of a fluid internally secreted. Some of these skin-glands remain unbranched, as for instance, the sweat-glands (*e, f, g*). These glands, which secrete the sweat, are of great length, their ends forming a coil; they never branch, however; and the same is to be said of the glands which secrete the fatty wax of the ears.



FIG. 214.—Rudiments of tear-glands from a human embryo of four months. (After Koelliker.) 1. Earliest rudiment the shape of a simple, solid plug. 2 and 3. Further developed rudiments, which branch and become hollow: *a*, a solid offshoot; *e*, cell-covering of the hollow offshoot; *f*, rudiment of the fibrous covering, which afterwards forms the leather-skin round the glands.

most other skin-glands give out shoots and branches, as, for instance, the tear-glands, situated on the upper eyelid, which secrete the tears (Fig. 214), and also the

sebaceous glands, which produce the fatty sebaceous matter and generally open into the hair-follicles. The sweat and sebaceous glands occur only in Mammals. The tear-glands, on the contrary, are found in all the three classes of Amnion Animals, in Reptiles, Birds, and Mammals. They are not represented in the lower Vertebrates.

Very remarkable skin-glands, found in all Mammals, and in them exclusively, are the milk-glands (*glandula mammales*, Figs. 215, 216). They supply milk for the nourishment of the new-born Mammal. Notwithstanding

their extraordinary size, these important organs are merely large sebaceous skin-glands (Plate V. Fig. 16, *md*). The milk is produced by liquefaction of the fatty milk-cells within the branched milk-gland pouch (Fig. 215, *c*), just as the sebaceous matter of the skin, and the fatty matter of the hair are produced by the breaking up of fatty sebaceous cells within the sebaceous skin-glands. The excretory passages of the milk-glands enlarge into sac-like milk-ducts (*b*), which again become narrower (*a*), and open through from sixteen to twenty-four minute apertures, into the nipple of the breast. The first rudiment of this large and complex gland is a very simple conical plug in the




FIG. 215.—The breast of the female in section: *c*, grape-like glandular lobules; *b*, enlarged milk-ducts; *a*, narrow excretory ducts, opening through the breast-nipple. (After H. Meyer.)

FIG. 216.—Milk-glands of a new-born child: *a*, original central gland; *b*, smaller, and *c*, larger branches of the latter. (After Langer.)

of the ordinary epidermic cells. Within this a firm central cellular mass of conical shape soon forms. This increases considerably in length, detaches itself from the surrounding cellular mass, the "root-sheath," and finally makes its way to the outside, appearing above the outer surface as a hair-stem. The deepest part, buried in the skin, the hair follicle, is the root of the hair, and is surrounded by the root-sheath. In the human embryo the first hairs make their appearance at the end of the fifth or in the beginning of the sixth month.

During the last three or four months before birth the human embryo is usually covered by a thick coating of delicate woolly hairs. This embryonic wool-covering (*lanugo*) is often lost during the last weeks of embryonic life, and at any rate, soon after birth, when it is replaced by a thinner permanent hair-covering. These later permanent hairs grow out of hair follicles which are developed from the root-sheaths of the deciduous woolly hair. In the human embryo, the embryonic woolly hair usually covers the entire body, with the exception of the palms of the hands and the soles of the feet. These parts remain bare just as in all Apes and most other Mammals. Not infrequently the woolly coat of the embryo differs considerably in colour from the later permanent hairy covering. Thus for instance, it sometimes happens in our own Indo-Germanic race that fair-haired parents are shocked to find their children, at their first appearance, covered by a dark brown, or even black, woolly covering. It is only after the woolly coat has been shed, that the permanent fair hair, which the child inherits from its parents, makes its appearance. Occasionally the dark hair is retained for several weeks

even months, after birth. This remarkable woolly covering can only be explained as an inheritance from our primordial long-haired ancestors, the Apes.

It is equally worthy of note that many of the higher Apes resemble Man in the thin coat of hair which covers certain parts of their body. In most Apes, especially in the higher Catarhines, the face is nearly or even quite bare, or is covered with hairs as thin and as short as those of Man. In these Apes also, just as in Man, the hair on the back of the head is usually distinguished by its length, and the males often have much beard and whisker. (Cf. fig. 202, p. 175). In both cases this masculine adornment has been acquired in consequence of sexual selection. In some Apes the breast and the inner sides of the joints are very thinly covered with hair—far less abundantly than the back and the outer sides of the joints. On the other hand, we not unfrequently see the shoulders, the back, and the outer sides of the limbs thickly covered with hair in men of Indo-Germanic or Semitic race. It is a well-known fact that in some families abundant hair on the body is hereditary, as is the relative vigour and character of the hair-growth of the beard and head. These great differences in the total and partial hairiness of the body, which appear very striking not only when we compare different races of man, but even when we compare many families belonging to the same race, are very simply explained by the fact that the entire hairy covering of Man is a rudimentary organ, an unused inheritance, which has been transmitted from the more hirsute Apes. In this matter, Man resembles the Elephant, Rhinoceros, Hippopotamus, Whale, and other mammals of various orders which have also entirely or

partially lost their original coat of hair in consequence of adaptation.¹⁶⁸

The form of Adaptation which has degraded the growth of hair on most parts of the human body, while preserving it, or even greatly developing it, on certain parts, was, in all probability, sexual selection. As Darwin has very clearly shown in his work on "The Descent of Man," sexual selection has had especially great influence in this respect. In consequence of the male Anthropoid Apes, in selecting a partner, preferring those females which were least hairy, and in consequence of the females preferring those suitors which were distinguished by peculiarly fine beard or head-hair, the general hirsuteness of the body was gradually degraded, while the beard and the hair of the head were advanced to a higher degree of perfection. Climatic conditions, and other circumstances unknown to us, may, however, also have promoted the loss of the hairy coat.

In proof of the assertion that the hairy covering of Man is directly inherited from the Anthropoid Apes, we find, according to Darwin, a curious evidence in the direction, otherwise inexplicable, in which the rudimentary hairs lie on our arms. Both on the upper and on the lower arm the hairs are directed towards the elbow, where they meet at an obtuse angle. Except in Man, this striking arrangement occurs only in the Anthropoid Apes, the Gorilla, Chimpanzee, Orang, and several species of Gibbons. In other Gibbons the hairs of both the lower and the upper arm are directed towards the hand, as in other Mammals. This remarkable peculiarity of Anthropoids and of Man can only be explained on the assumption that our common ape-like ancestors were accustomed, as they are even now,

During rain, to bring their hands together over their heads, or over a branch overhanging their heads. The reverse direction of the hairs, when the arms were in this position caused the rain to run off. Thus, even yet, the direction of the hairs on our lower arm testifies to this advantageous habit of our Ape-ancestors.

If the skin and its appendages are minutely examined, Comparative Anatomy and Ontogeny supply many similar important "records of creation," showing that they are directly inherited from the skin-covering of the Ape. We obtained our skin and hair by inheritance, immediately from Anthropoid Apes, these from the lower Apes, which, in turn, inherited the same parts from lower Mammals. This is also true of the other great organ-system which is developed from the skin-sensory layer—of the nervous system and the sensory organs. This very highly developed organ system, which performs the highest vital functions—those of the mind—we have inherited immediately from the Apes, and mediately from Mammals of a lower order.

The human nervous system, like that of all other Mammals, is, in its developed condition, a very complex apparatus, the anatomical arrangement and the physiological activity of which may, in general terms, be compared to a telegraph system. The central marrow (*medulla*), or central nervous system, represents the principal station, the innumerable "ganglion cells" (Fig. 7, vol. i. p. 129) of which are connected with each other and with numerous very delicate conducting lines by their branched processes. The latter are the peripheric "nerve fibres," distributed over the whole surface of the body; these, together with their terminal apparatus, the sense-organs, etc., constitute the "conductive

marrow," the peripheric nerve-system. Some, as sensory nerve-fibres, convey the sensations of the skin and of sense-organs to the central medulla; others, as motor nerve-fibres, transmit the impulses from the central marrow to the muscles.




FIG. 217.—Human embryo of three months, in natural size, seen from the dorsal side, the brain and dorsal cord exposed (after Koelliker): *h*, large hemispheres of the cerebra (fore-brain); *m*, "four-bulbous" (mid-brain); *c*, small hind-brain, or cerebellum. Below the latter is the three-cornered "neck-marrow" (after-brain).

FIG. 218.—Central nervous system of a human embryo of three months, in natural size, seen from the dorsal side (after Koelliker): *h*, large hemispheres; *v*, "four-bulbous" small brain; *mo*, neck-marrow. Below this the dorsal cord (marrow).

The central nervous system, or central marrow (*medulla centralis*), is the actual organ of mental activities, in a stricter sense. Whatever view is taken of the intimate connection between this organ and its functions, it is at least certain that those of its special activities which call sensation, volition, and thought, are in man, as in the higher animals, inseparably connected with the development of this material organ. Hence we must necessarily take a deep interest in the history of the develop-

of this organ. As it alone can give us the most important information as to the nature of our "mind," it commands our most earnest attention. For if the central marrow develops in the human embryo exactly as in the embryos of all other Mammals, then the development of the human mental organ from the same central organ of other Mammals and, more remotely, from that of lower Vertebrates, cannot be questioned. It is, therefore, impossible to dispute the enormous significance of these phenomena of development.

In order to appreciate these rightly, a few words must first be said as to the general form and anatomical construction of the developed central marrow in Man. Like the central nervous system of all other Skulled Animals (*Craniota*), it consists of two distinct parts: firstly, of the brain or the medulla of the head (*encephalon*, or *medulla cerebri*), and, secondly, of the spinal marrow (*medulla spinalis*). The former is enclosed in the bony skull, or "brain case," the latter in the bony vertebral canal, which is composed of a consecutive series of vertebræ, shaped like signet rings. (Cf. Plate V. Fig. 16, *m*.) From the brain proceed twelve pairs of head nerves, from the spinal marrow thirty-one pairs of medullary or spinal nerves for the remainder of the body. The spinal marrow, when examined merely anatomically, appears as a cylindrical cord with a spindle-shaped swelling in the region of the neck (at the last of the neck-vertebræ) and another in the lumbar region (at the last lumbar vertebra, Figs. 217, 218). At the swelling at the throat the large nerves of the upper limbs pass off from the spinal marrow, and those of the lower limbs from the swelling in the lumbar region. The upper end of the spinal marrow passes through the neck-marrow (*medulla oblon-*

gata) into the brain. The spinal marrow appears indeed to be a dense mass of nervous substance; but along its axis passes a very narrow canal, which is continued in front into the larger cavities of the brain, and which, like those cavities, is filled with a clear fluid.

The brain forms a considerable mass of nervous substance, of very complex, minute structure, which occupies




FIG. 219.—Human brain, seen from the lower side. (After H. Meyer.) Above (in front) is the large brain (*cerebrum*), with extensively branched furrows; below (behind) is the small brain (*cerebellum*), with narrow parallel furrows. The Roman numbers indicate the roots of the twelve pairs of brain nerves in order from front to back.

the greater part of the skull-cavity; it is roughly distinguishable into two main parts—the large and small brain (*cerebrum* and *cerebellum*). The former is situated in front and over the latter, and its surface exhibits the well-known characteristic convolutions and furrows (Figs. 219, 220). On its upper surface it is divided by a deep longitudinal slit into two lateral halves, the so-called “great hemispheres,” which are connected by means of a bridge, or “cross-piece” (*corpus callosum*). A deep transverse fissure separates the large brain (*cerebrum*) from the small brain

bellum). The latter is situated more posteriorly and anteriorly, and shows on its outer surface equally numerous fowls, which are, however, much finer and more regular,



Fig. 220.—Human brain, seen from the left side. (After H. Meyer.) The fowls of the large brain are indicated by large, thick lines, those of the small brain by finer lines. Below the latter the neck-marrow is visible. f^1 – f^{100} , frontal convolutions; *Ce. a*, *Ce. p*, central convolutions; *R*, fissure of Rolando; *S*, Sylvian fissure; *T*, temporal or parallel fissure; *Pa*, parietal lobe; *An*, annectant convolutions; *PO*, parieto-occipital fissure; *Su*, supra-marginal convolution; *IP*, intra-parietal fissure; *t*, temporo-sphenoidal convolution.

between them are curved ridges (Fig. 219, lower part). The small brain is also divided into two lateral halves by a longitudinal furrow; these are the "small hemispheres," which are connected at the top by a worm-like cross-piece, the "brain-worm" (*vermis*), and at the bottom by a bridge, the *pons varolii*; Fig. 219, VI.).

Comparative Anatomy and Ontogeny show, however, that man, as in all other Skulled Animals, the brain originally consists not of two but of five distinct parts lying one behind another. These originally appear in the embryo of all

Skulled Animals (*Craniota*), from the Cyclostomi and Fishes up to Man, in exactly the same form, as five bladders placed one behind the other. Alike in their first rudiments, they, however, differ in their further development. In Man and all higher Mammals the first of these five bladders, the fore-brain, develops so excessively that, when mature, it forms, both in size and weight, by far the greater part of the whole brain. To it belong, not only the great hemispheres, but also the bridge (*corpus callosum*), which connects these two, the olfactory lobes, from which proceed the nerves of smell, and most of the processes lying on the roof and floor of the great lateral cavities of the two hemispheres; such, for instance, as the large streaked bodies (*corpora striata*). On the other hand, the "centres of sight," which lie between the streaked bodies, belong to the second main part, which develops from the twixt-brain; and to the same part belong the third brain ventricle (which is single) and the processes known as the "funnel" (*infundibulum*), the gray mass, and the "cone" (*conarium*). Behind these, and between the large brain and the small brain, we find a little mass, composed of two pairs of bosses, and called the "four bulbs," on account of two superficial furrows which cross each other at right angles, thus quartering the whole mass (Figs. 217, *m*, 218, *v*). Though these "four bulbs" are very insignificant in Man and the higher Mammalia, they constitute a distinct part of the brain, the third, or mid-brain, which is, on the contrary, especially well developed in the lower Vertebrates. The next or fourth part of the brain is the hind-brain, or small brain (*cerebellum*), in the strict sense of the term, with its single middle process, the "worm" (*vermis*), and its two lateral parts, the "small

hemispheres" (Figs. 217, c, 218, c). Behind this comes, finally, the fifth and last part, the "neck-marrow" (*medulla oblongata*, Fig. 218, mo), which includes the single fourth brain ventricle and the adjoining processes (pyramids, olives, and testiform bodies). The neck medulla passes directly down into the spinal marrow. The narrow central canal of the spinal marrow extends into the wider "fourth ventricle" of the neck medulla, which is rhomboidal in shape, and the floor of which forms the "rhomboid groove." From this proceeds a narrow duct, called the "aqueduct of Sylvius," which leads through the "four-bulbs" into the third ventricle, situated between the two "centres of sight;" and this cavity in turn is connected with the pair of lateral cavities which lie right and left in the large hemispheres. All the cavities of the central marrow are, therefore, directly connected together. Individually all these parts of the brain which we have enumerated have an infinitely complex, minute structure, which we cannot now study, and which hardly bears on our subject. This wonderful brain-structure, as it occurs only in Man and the higher Vertebrates, is of the highest importance, simply because, in all Skulled Animals (*Craniota*), it develops from the same simple rudiments, from the five brain-bladders already enumerated. (Cf. Plates VI. and VII.)


Before we direct our attention to the individual development of the complex brain from this series of simple bladders, we will, in order to understand the matter more clearly, glance for a moment at those lower animals which have no such brain. Even in the skull-less Vertebrates, in the *Amphioxus*, there is no real brain. In this case the whole central marrow is merely a simple cylindrical cord

traversing the body longitudinally, and terminating in front almost as simply as at the other end: it is a simple medullary tube (Plate XI. Fig. 15, *m*). We found, however, that the rudiment of the same simple medullary tube occurs in the ascidian larva (Plate X. Fig. 5, *m*) and in the same characteristic position, above the notochord. Moreover, when closely examined a small bladder-like swelling may be seen at the fore end of the medullary tube in these two closely allied animals; this is the first indication of a separation of the medullary tube into brain (*m*₁) and spinal marrow (*m*₂). When, however, we consider the undeniable relationship of the *Ascidia* to the rest of the Worms, it is evident that the simple central marrow of the former exactly answers to the simple nerve-ganglion which, in the lower Worms lies above the throat (*pharynx*), and which has, therefore, long been called the "upper throat ganglion" (*ganglion pharyngeum superius*). In the Gliding Worms (*Turbellaria*) the whole nerve system consists merely of this simple ganglion, which is situated on the dorsal side of the body, and from which nerve-threads radiate to the different parts of the body (Fig. 211, *gn*). This upper throat ganglion of the lower Worms is evidently the rudiment from which the more complex central marrow of the higher animals has developed. An elongation of the upper throat ganglion along the dorsal side gave rise to the medullary tube, which is characteristic of Vertebrates and the young forms of *Ascidia* alone. On the other hand, in all other animals, the central nerve system has developed in a very different manner from the upper throat ganglion; in Articulated Animals (*Arthropoda*) especially, the latter has developed into a throat (pharyngeal) ring, with a ventral marrow; this is the case, also, in the articu-

ed Ringed Worms (*Annelida*) and the Star-animals (*Echinodermata*), which originated from Arthropods. The Soft-shelled Animals (*Mollusca*) also have a throat ring, which is quite unrepresented in Vertebrates. Only in Vertebrates is the central marrow developed along the dorsal side, while in all other animals which have been named it is developed along the ventral side of the body.¹⁶⁹

Descending below the Worms we find very many animals which are entirely without a nerve-system, and in which the functions of that system are performed simply by the outer skin-covering—by the cells of the skin-layer, or derm. This is the case in many low Plant Animals (*Phytophyta*), for instance, in all Sponges, and in the common fresh-water Polyp, the Hydra. It was also undoubtedly the case in all extinct Gastræans. In all Primitive Animals (*Protozoa*) the nerve-system is, of course, unrepresented, for they have not as yet attained to the development of germ-cells.

In considering the individual development of the nerve-system in the human embryo, we must first of all start from the important fact already mentioned, that the first rudiment of the system is the simple medullary tube, which arises itself from the outer germ-layer along the middle of the lyre-shaped primitive germ. We found (Figs. 87, vol. i. p. 298) that the rectilineal primitive groove, or neural furrow, first arises in the centre of the lyre-shaped germ-disc. On each side of this rise the two parallel dorsal medullary swellings. The free margins of these bend towards each other, coalesce, and form the closed medullary tube (Figs. 88–93, vol. i. pp. 300–309). At first this tube lies directly under the horn-plate; it is, however, afterwards situate



FIGS. 221-223.—Lyre-shaped (or sole-shaped) germ-shield of a Chick, in three consecutive stages of evolution, seen from the dorsal surface: above twenty times enlarged. Fig. 221, with six pairs of primitive vertebrae. The brain a simple bladder (*hb*). The medullary furrow is wide open from the point *x*, very wide at *z*. *mp*, Marrow (or medullary) plates; *sp*, side plates; *y*, boundary between the throat cavity (*sh*) and the head-intestine (*vd*). Fig. 222, with ten pairs of primitive vertebrae. The brain consists of three bladders: *v*, fore-brain; *m*, mid-brain; *h*, hind-brain. *c*, Heart; *d*, yolk-veins. The medullary furrow is wide open behind (*z*). *mp*, Marrow plates. Fig. 223, with sixteen pairs of primitive vertebrae. The brain consists of five bladders: *v*, fore-brain; *z*, twist-brain; *m*, mid-brain; *h*, hind-brain; *n*, after-brain. *a*, Eye-vesicles; *g*, ear-vesicles; *c*, heart; *d*, yolk-veins; *mp*, marrow-plate. *uw*, primitive vertebrae.

site internally, the upper edges of the primitive vertebral plates, which penetrate, from right and left, in between the notochord-plate and the medullary tube, uniting above the latter, and thus completely embedding it in a closed canal. As Gegenbaur most aptly remarks, "This gradual embedding of the interior of the body must be regarded as an incident required in connection with progressive differentiation, and with the consequent higher capacity, by which the most important organ of the system is secured in its interior."

To every thoughtful and unprejudiced man it must appear an extremely important and pregnant fact, that our mental organ, like that of all other Skulled Animals (*Cranota*), commences in the same way and in exactly the same simple form in which this organ remains for life in the lowest Vertebrate, the *Amphioxus* (vol. i. p. 420, Fig. 151; Plate XI. Fig. 15, *m*). In the Cyclostomi, that is, in the stage above the Acrania, the anterior extremity of the cylindrical medullary tube begins to extend, at an early period, in the form of a pear-shaped bladder, which is the first distinct rudiment of a brain (Plate XI. Fig. 16, *m*₁). For the central medulla of Vertebrates thus first distinctly differentiates into its two main sections, the brain (*m*₁) and the spinal marrow (*m*₂). The first faint indication of this important differentiation is discoverable in the *Amphioxus*, perhaps even in the Ascidian larva (Plate X. Fig. 5).

The simple bladder-like form of the brain, which is retained for a considerable time in the Cyclostomi, also appears at first in all higher Vertebrates (Fig. 221, *hb*). In the latter, however, it soon disappears, in consequence of the separation of the simple brain-bladder, by transverse contractions of its circumference, into several consecutive

parts. Two of these contractions first appear, and subsequently the brain forms three consecutive bladders (



FIGS. 224-226.—Central row of human embryo in seventh week, two cm. (After Koelliker.) Fig. view of the whole embryo the dorsal side; the brain dorsal marrow laid bare. 225, the brain and upper of the dorsal marrow from

left side. Fig. 224, the brain from above: *v*, fore-brain; *z*, twist-brain; *m*, mid-brain; *h*, hind-brain; *n*, after-brain.

222, *v*, *m*, *h*). The first and third of these three primary bladders then again separate by transverse contraction each into two parts, and thus five consecutive bladder divisions are formed (Fig. 223: cf. also Plate V. 13-16; Plates VI. and VII., second cross-line). These fundamental brain-bladders, which re-occur in the same in the embryos of all the Skulled Animals (*Craniota*), first clearly recognized by Baer, who understood their importance and distinguished them, according to their relative positions, by very appropriate names, which are in general use: I., fore-brain (*v*); II., twist-brain (*z*); III., mid-brain (*m*); IV., hind-brain (*h*); and V., after-brain (*n*).

In all Skulled Animals, from the Cyclostomi to the same parts, although in very various forms, develop from these five original brain-bladders. The first bladder, the fore-brain (*protopsyche*, *v*), forms by far the largest of the so-called "great brain" (*cerebrum*); it forms the great hemispheres, the olfactory lobes, the streaked bodies (*corpora striata*), and the cross-piece (*corpus callosum*) together with the "arch" (*fornix*). From the second

adder, the twixt-brain (*deutopsyche*, *z*), proceed primarily to the "centres of sight" and the other parts which surround the so-called "third brain-ventricle," also the "funnel" (*infundibulum*), the "cone" (*conarium*), etc. The third bladder, the mid-brain (*mesopsyche*, *m*), furnishes the small group of the "four bulbs," together with the "aqueduct of Sylvius." From the fourth bladder, the hind-brain (*metapsyche*, *h*), the greater part of the so-called "little brain" (*cerebellum*) develops; the central "worm" (*vermis*), and the two lateral "small hemispheres." The fifth bladder, finally, the after-brain (*epipsyche*, *n*), forms the neck-marrow, or the "elongated marrow" (*medulla oblongata*), together with the rhomboid groove, the pyramids, olives, etc.

The very highest importance must certainly be ascribed to the fact, seen in Comparative Anatomy and Ontogeny, that the brain is originally formed in exactly the same way in the embryos of all Skulled Animals (*Craniota*), from the lowest Cyclostomi and Fishes, to Apes and Man. In all, the first rudiment of the brain is a simple bladder-like expansion at the anterior extremity of the medullary tube. In all, the five bladders are formed from this simple bladder-like expansion, and in all, these five primitive brain-bladders develop into the permanent brain, with its many complex anatomical arrangements, which afterwards appear in such extremely diverse forms in the various vertebrate classes. On comparing the mature brain of a Fish, an Amphibian, a Reptile, a Bird, and a Mammal, it is hardly conceivable that the several parts of these forms, so extremely different, both internally and externally, may be traced back to one common condition. And yet, all these various brains of *Craniota* have originated from exactly the

same rudimentary form. We need only compare the embryos of these various classes of animals at corresponding stages of development, in order to assure ourselves of the fundamental fact. (Cf. Plates VI. and VII., second cross-line.)




FIG. 227.—Brains of three embryonic Skulled Animals in vertical longitudinal sections: A, of a Shark (*Heptanchus*); B, of a Snake (*Coluber*); C, of a Goat (*Capra*); a, fore-brain; b, twixt-brain; c, mid-brain; d, hind-brain; e, after-brain; s, primitive fissure of the brain. (After Gegenbaur.)

FIG. 228.—Brain of a Shark (*Scyllium*) from the dorsal side: g, fore-brain; h, olfactory bulbs of the fore-brain, which send the large olfactory nerves to the large nose capsules (o); d, twixt-brain; b, mid-brain (behind it, the insignificant rudiment of the hind-brain); a, after-brain. (After Busch.)

FIG. 229.—Brain and dorsal marrow of a Frog: A, from the dorsal side; B, from the ventral side; a, olfactory bulbs, in front of the fore-brain (b); i, funnel at the base of the twixt-brain; c, mid-brain; d, hind-brain; s, rhomboid groove in the after-brain; m, dorsal marrow (very short in the frog); m', root-processes of the spinal nerves; t, fibre at the end of the dorsal marrow. (After Gegenbaur.)

A thorough comparison of the corresponding stages of development in the brain in the various Skulled Animals (*Craniota*) is very instructive. If it is applied to the whole series of skulled classes, the following extremely interesting facts soon become evident: in the Cyclostomi (*Myxinoidea* and *Petromyzontes*), which, as we have seen, are the lowest and earliest Skulled Animals, the whole brain remains for life at a very low and primitive stage of development, through which the embryos of the other skulled Animals pass very rapidly; the five original sections of the brain are visible throughout life in an almost unmodified form. But even in Fishes, an essential and important transformation of the five bladders takes place; it is evidently from the brain of the Primitive Fishes (*Selachii*; Fig. 228), that, on the one side, the brain of the other Fishes, and on the other, the brain of the Amphibians and also of the higher Vertebrates, must be traced. In Fishes and Amphibians (Fig. 229), the central part, the mid-brain, and also the fifth section, the after-brain, are specially developed, while the first, second, and fourth sections remain far behind. In the higher Vertebrates, the exact reverse is the case, for in these the first and fourth sections, the fore and hind brains, develop pre-eminently; on the other hand, the mid-brain remains very small, and the after-brain is also much smaller. The greater part of the "four-bulbs" is covered by the large brain (*cerebrum*) and the after-brain by the small brain (*cerebellum*). Even among the higher Vertebrates themselves, numerous gradations occur in the structure of the brain. From the Amphibians upward, the brain, and with it the mental life, develops in two different directions, of which the one is

carried out in Reptiles and Birds, the other in Mammals. The latter are especially distinguished by the very characteristic development of the first section, the fore-brain.




FIG. 230.—Brain of Rabbit: A, from the dorsal side; B, from ventral side; lo, olfactory lobules; I., fore-brain; h, hypophysis at the base of the mid-brain; III., mid-brain; IV., hind-brain; V., after-brain; 2, optic nerve; 3, motor nerve of the eye; 5-8, fifth to eighth nerves of the brain. In A, the upper surface of the right large hemisphere (I.) is removed so that the streaked bodies (*corpora striata*) can be seen in its side chamber (*ventriculus lateralis*). (After Gegenbaur.)

Mammals alone (Fig. 230) does this "great brain" develop to such an extent, that it eventually covers all the other parts of the brain from above.

There are also remarkable differences in the relative positions of the brain-bladders. In the lower Skull Animals the five brain-bladders are at first situated behind the other in the same plane. If the brain is regarded from the side, a straight line may be drawn through all the five bladders. But in the three higher vertebrate classes, in the Amnion Animals (*Amniota*), a noticeable curving of the rudimentary brain takes place, simultane-

With the head and neck curving of the whole body, owing to the fact that the whole upper dorsal surface of the brain grows much faster than the lower ventral surface. The result is that the brain is so curved that its parts are afterwards situated thus: the fore-brain lies quite in front and below, the twixt-brain somewhat higher and over it, while the mid-brain lies highest of all and projects furthest forward; the hind-brain is situated lower, the after-brain yet further back and below. This disposition occurs only in the three classes of the Amniota, in Reptiles, Birds, and Mammals. (Cf. Plates I., VI., and VII.)

Though, in general features of growth, the brains of Mammals correspond with those of Birds and Reptiles, yet striking differences very soon appear between the two. In Birds and Reptiles (Plate VI. Figs. *H* and *C*), the mid-brain (*m*) and the central part of the hind-brain develop considerably. In Mammals, on the other hand, these parts remain small, and instead, the fore-brain begins to grow so rapidly that it covers the other bladders from in front and above. As it constantly grows further back, it eventually covers the whole of the rest of the brain above, and also encloses the central part from the sides. This process is of the greatest importance, because this fore-brain is the organ of the higher mental activities,—because in it are accomplished those functions of the nerve-cells, the sum of which is generally designated as the mind, or the “spirit” in the narrower sense. The highest activities of the animal body, the wonderful manifestations of consciousness, the complex phenomena of the activities of thought, have their seat in the fore-brain. It is possible to remove the great hemispheres of a Mammal, piece by piece, without killing

the animal, thus proving that the higher mental activities, consciousness and thought, conscious volition and sensation, may be destroyed one by one, and finally entirely annihilated. If the animal thus treated is artificially fed, it may be kept alive for a long time; for the nourishment of the entire body, digestion, respiration, the circulation of the blood, secretion, in short, the vegetative functions, are in no way destroyed by this destruction of the most important mental organs. Conscious sensation and voluntary motion, the capacity for thought and the combination of the various higher mental activities, have alone been lost.

This fore-brain, the source of all these most wonderful nervous activities, reaches that high degree of perfection only in the higher Placental Animals (*Placentalia*); a fact which explains very clearly why the higher Mammals so far excel the lower in intellectual capacity. While the "mind" of the lower Placental Animals does not exceed that of Birds and Reptiles, we find among the higher *Placentalia* an uninterrupted gradation up to Apes and Man. Accordingly, their anterior brains show surprising differences in the degree of perfection. In the lower Mammals, the surface of the great hemispheres (the most important part) is entirely smooth and even. The fore-brain, too, remains so small that it does not even cover the mid-brain above (Fig. 230). One stage higher, and this latter is indeed entirely covered by the excessive growth of the fore-brain; but the hind-brain remains free and uncovered. At last, in Apes and in Man, the fore-brain covers the hind-brain also. A similar gradual advance may also be traced in the development of the peculiar furrows and protuberances which are so characteristically prominent on the surface of the large brain

(*cerebrum*) of higher Mammals (Figs. 219, 220). If the brains of the various mammalian groups are compared with reference to these convolutions and furrows, it appears that their gradual development is entirely proportionate with the development of the higher intellectual activities. Much attention has recently been devoted to this particular branch of the Anatomy of the brain, and very striking individual differences have been found even within the human race. In all human individuals distinguished by peculiar ability and great intellect, these swellings and furrows on the surface of the great hemispheres exhibit a much greater development than in common average men; while in the latter, again, they are more developed than in Cretins and others of unusually feeble intellect. There are also similar gradations in the internal structure of the fore-brain in Mammals. The great cross-piece (*corpus callosum*), especially, the bridge between the two great hemispheres, is developed only in Placental Animals. Other arrangements, for example, in the structure of the lateral cavities, which seem primarily to be peculiar to Men as such, re-appear only in the higher species of Apes. It was long believed that Man had some entirely peculiar organs in the great brain (*cerebrum*), which are wanting in all other animals. But close comparison has shown that this is not the case, but that rather the characteristic qualities of the human brain exist in a rudimentary state even in the lower Apes, and are developed to a greater or less degree in the higher Apes. Huxley, in his important and much-quoted book, "Evidence as to Man's Place in Nature" (1863), has shown, most convincingly, that within the Ape-series the differences in the formation of the brain are greater between the

higher and lower Apes than between the higher Apes and Man. This statement is, indeed, equally true of all the other parts of the body. But the fact that it is true of the central marrow is especially important. This does not become fully evident unless these morphological facts are considered in connection with the corresponding physiological phenomena; until we consider that every mental activity requires for its complete and normal exercise the complete and normal condition of the corresponding brain-structure. The extremely complex and perfect active phenomena within the nerve-cells, summed up in the word "mental life," can no more exist without their organs in the vertebrates, including man, than can the circulation of the blood without a heart or blood. As, however, the central marrow of Man has developed from the same medullary tube as in all other Vertebrates, so also must the mental life of Man have had the same origin.

All this is of course true of the conductive marrow, or the so-called "peripheric nervous system." This consists of the *sensitive* nervous fibres which convey the impressions of sensation from the skin and the organs of the senses in a centripetal direction to the central marrow; as well as of the *motor* nervous fibres, which, reversely, convey the movements of volition from the central marrow, in a centrifugal direction to the muscles. By far the greater part of these peripheric conductive nerves originates from the skin-fibrous layer, by peculiar local differentiation of the rows of cells into the respective organs.

The membranous coverings and blood-vessels of the central marrow are identical in origin with the greater part of the conductive marrow; these membranous coverings

are the inner membrane (*pia mater*), the central membrane (*meninx arachnoides*), and the outer membrane (*dura mater*). All these parts are developed from the skin-fibrous layer.

TABLE XXVII.

SYSTEMATIC SURVEY OF THE MOST IMPORTANT PERIODS IN THE PHYLOGENY OF THE HUMAN SKIN-COVERINGS.

I. *First Period: Skin of Gastræads.*

The entire skin-covering (including the nervous system, not yet differentiated from it) consists of one simple layer of ciliated cells (exoderm, or primary skin-layer); as it is at the present day in the gastrula of the *Amphioxus*.

II. *Second Period: Skin of Primitive Worms.*

The simple exoderm of the Gastræad has thickened and split into two distinct layers, or secondary germ-layers: the skin-sensory layer (rudiment of the horn-plate and nerve-system) and the skin-fibrous layer (rudiment of the leather skin (*corium*), the muscle-plate and the skeleton-plate. The skin is potentially both covering and mind.

III. *Third Period: Skin of Chordonia.*

The skin-sensory layer has differentiated into the horn-plate (*epidermis*), and the central marrow (upper throat ganglia) separated from it; the latter elongates into a medullary tube. The skin-fibrous layer has differentiated into the leather plate (*corium*) and, below this, the skin-muscular pouch (as in all Worms).

IV. *Fourth Period: Skin of Acrania.*

The horn-plate yet forms a simple epidermis. The leather-plate is fully differentiated from the muscle and skeleton plates.

V. *Fifth Period : Skin of Cyclostoma.*

The outer-skin remains a simple, soft mucous layer of cells, but forms one-celled glands (cup-cells). The leather-skin (*corium*) differentiates into *cutis* and *sub-cutis*.

VI. *Sixth Period : Skin of Primitive Fishes.*

The outer skin is still simple. The leather skin forms placoid scales or small bony tablets, as in the *Selachii*.

VII. *Seventh Period : Skin of Amphibia.*

The outer skin differentiates into an outer horn-layer, and an inner mucous layer. The ends of the toes are covered with horny sheaths (first rudiments of claws or nails).

VIII. *Eighth Period : Skin of Mammals.*

The outer skin forms the appendages characteristic of *Mammals* only; hair, and sebaceous, sweat, and milk glands.

TABLE XXVIII.

SYSTEMATIC SURVEY OF THE MOST IMPORTANT PERIODS IN THE PHYLOGENY OF THE HUMAN NERVOUS SYSTEM.

I. *First Period : Medulla of Gastræads.*

The nerve system is not yet distinct from the skin, and, together with the latter, is represented by the simple cell-stratum of the exoderm, or primary skin-layer; as it is at the present day in the gastrula of the *Amphioxus*.

II. *Second Period : Medulla of Primitive Worms.*

The central nerve system is yet, at first, a part of the skin-sensory layer, and afterwards consists of a throat medulla, a simple nerve-ganglion lying above the throat; as it is now in the lower *Worms*: the upper throat-ganglion.

III. *Third Period : Medulla of Chordonia.*

The central nerve system consists of a simple medullary tube, anterior extension of the upper throat ganglion, which is separated from the intestine by a notochord (*chorda dorsalis*).

IV. *Fourth Period : Medulla of Acrania.*

The simple medullary tube differentiates into two parts: a head, and a spinal part. The head medulla resembles a small, pear-shaped, simple rudiment (the primitive brain, or first rudiment of the brain) on the anterior end, and the spinal part is the long cylindrical spinal marrow.

V. *Fifth Period : Medulla of Cyclostoma.*

The simple, bladder-like rudiment of the brain divides into five connected brain-bladders of simple structure.

VI. *Sixth Period : Medulla of Primitive Fishes.*

The five brain-bladders differentiate into a form similar to that now generally retained by the Selachii.

VII. *Seventh Period : Medulla of Amphibia.*

The differentiation of the five brain-bladders progresses to that structure which is now characteristic of the brain in Amphibia.

VIII. *Eighth Period : Medulla of Mammals.*

The brain attains the characteristic peculiarities distinctive of Mammals. The following may be distinguished as subordinate stages of development: 1, the brain of Monotremes; 2, the brain of Marsupials; 3, the brain of Apes; 4, the brain of Apes; 5, the brain of Man-like Apes; 6, the brain of Ape-men; and 7, the brain of Man.

TABLE XXIX.

Systematic Survey of the Evolution of the Skin-covering and Nerve System.

XXIX. A. Survey of the Evolution of the Skin-covering.

Skin (<i>Derma,</i> or <i>Integumentum</i>)	Outer-skin (<i>Epidermis</i>) Product of the Skin- sensory-layer	Horn-layer of the outer skin (<i>Stratum corneum</i>) Mucous layer of the outer skin (<i>Stratum mucosum</i>)	Hair Nails Sweat glands Tear glands Sebaceous glands Milk glands
	Leather-skin (<i>Corium</i>) Product of the Skin- fibrous-layer	Fibrous layer of the leather skin (<i>Cutis</i>) Fatty layer of the leather skin (<i>Subcutis</i>)	Connective tissue Fatty tissue Muscular tissue Blood-vessels Papillae of taste and nerves of the leather skin

XXIX. B. Survey of the Evolution of the Central Marrow.

Central Marrow, or Central Nerve System (<i>Psyche, or Medulla Centralis</i>). Product of the Skin-sensory layer	I. Fore-brain (<i>Protopsyche</i>)	Great hemispheres Olfactory lobules Lateral chambers Streaked bodies Arch Cross piece	<i>Hemisphaera cerebri</i> <i>Lobi olfactorii</i> <i>Ventriculi laterales</i> <i>Corpora striata</i> <i>Fornix</i> <i>Corpus callosum</i>
	II. Twixt-brain (<i>Deutopsyche</i>)	Centre of sight Third chamber of the brain Pineal body Funnel	<i>Thalami optici</i> <i>Ventriculus tertius</i> <i>Conarium</i> <i>Infundibulum</i>
	III. Mid-brain (<i>Mesopsyche</i>)	Four bulbs Aqueduct of Sylvius Brain stalks	<i>Corpus bigeminum</i> <i>Aqueductus Sylvii</i> <i>Pedunculi cerebri</i>
	IV. Hind-brain (<i>Metapsyche</i>)	Small hemispheres Brain worm Brain bridge	<i>Hemisphaera cerebelli</i> <i>Vermis cerebelli</i> <i>Pons Varolii</i>
	V. After-brain (<i>Epipsyche</i>)	Pyramids Olivae Restiform bodies Fourth chamber of the brain	<i>Corpora pyramidalia</i> <i>Corpora olivaria</i> <i>Corpora restiformia</i> <i>Ventriculus quartus</i>
	VI. Dorsal Marrow	<i>Notopsyche</i>	<i>Medulla spinalis</i>
Medullary coverings (<i>Meninges</i>)	Enveloping mem- branes, with the nutritive blood- vessels of the brain and spinal cord	1. Soft medullary skin 2. Central medullary skin 3. Hard medullary skin (Products of the skin-fibrous layer)	<i>Pia mater</i> <i>Arachnoides</i> <i>Dura mater</i>

CHAPTER XXI.

DEVELOPMENT OF THE SENSE-ORGANS.

Origin of the most highly Purposive Sense-organs by no Preconceived Purpose, but simply by Natural Selection.—The Six Sense-organs and the Seven Sense-functions.—All the Sense-organs originally Developed from the Outer Skin-covering (from the Skin-sensory Layer).—Organs of the Pressure Sense, the Heat Sense, the Sexual Sense, and the Taste Sense.—Structure of the Organ of Scent.—The Blind Nose-pits of Fishes.—The Nasal Furrows change into Nasal Canals.—Separation of the Cavities of the Nose and Mouth by the Palate Roof.—Structure of the Eye.—The Primary Eye Vesicles (Stalked Protuberances from the Twixt-brain).—Inversion of this Eye Vesicle by the Crystalline Lens, separated from the Horn-plate.—Inversion of the Vitreous Body.—The Vascular Capsule and the Fibrous Capsule of the Eyeball.—Eyelids.—Structure of the Ear.—The Apparatus for Perception of Sound : Labyrinth and Auditory Nerve.—Origin of the Labyrinth from the Primitive Ear Vesicles (by Separation from the Horn-plate).—Conducting Apparatus of Sound : Drum Cavity, Ear Bonelets, and Drum Membrane.—Origin of these from the First Gill-opening and the Parts immediately round it (the First and Second Gill-arch).—Rudimentary Outer Ear.—Rudimentary Muscles of the Ear-shell.

"Systematic Physiology is based especially upon the history of development, and unless this is more complete, can never make rapid progress ; for the history of development furnishes the philosopher with the materials necessary for the secure construction of a system of organic life. Hence anatomical and physiological researches should be prosecuted more from the

point of view of development than is now the case ; that is, we should determine each organ, each tissue, and even each function simply with the determining *whence they have arisen*."—EMIL HUSCHKE (1832).

THE sense-organs are undeniably among the most important and most interesting parts of the human body ; their activity alone we recognize the objects in the world around us. "*Nihil est in intellectu, quod non prius fuerit in sensu.*" They are the true springs of our mental life. In no other part of the animal body can we point to such extremely delicate and complex anatomical contrivances, co-ordinated for a definite physiological aim ; and in no other part of the body do these wonderful and very apt contrivances so clearly first, to indicate a premeditated creative design so obviously. Hence it is that, in accordance with the teleological view, it has been customary to admire the "wisdom of the Creator" and the "purposiveness of His Creation" especially in this matter. On more mature consideration it will be observed that the Creator, according to this conception, does after all but the part of an ingenious mechanic or of a skilful maker ; just, indeed, as all these cherished teleological conceptions of the Creator and His Creation are but childish anthropomorphism.

We admit that at first sight this teleological conception seems to afford the simplest and fittest interpretation of these very apt contrivances. If the structure and functions of the very highly developed sense-organs are regarded, it seems as though their origin is hardly explicable except on the assumption of a supernatural act. But it is exactly on this point that the his-

ion proves most clearly that this received conception is really false. The history of evolution convinces us that highly purposive and admirably constituted sense organs, and other organs, have developed without premeditation that they originated by the same mechanical process of Natural Selection, by the same constant interaction of adaptation and Heredity, by which all the other purposive contrivances of the animal organization have been formed and gradually evolved during the "Struggle for existence."

Like most other Vertebrates, Man possesses six distinct organs of sense, which accomplish seven distinct sensations. The external skin-covering accomplishes the sensation of pressure (resistance) and of temperature (warmth and cold). It is the earliest, the lowest, and the least differentiated organ of sense; it is distributed over the entire surface of the body. The other sensorial activities are localized. The sense of touch is limited to the skin-covering of the external organs, just as the sense of taste is limited to the mucous membrane of the mouth-cavity (tongue and palate), the sense of smell to the mucous membrane of the nasal cavity. Special mechanical contrivances of great complexity exist for the two highest and most differentiated organs of sense, the eye for the sense of sight, and the ear for the sense of hearing.

Comparative Anatomy and Physiology show that in the lower animals specialized sense-organs are entirely wanting, and all sensations are transmitted through the outer surface of the skin-covering. The undifferentiated skin-layer, or exoderm of the *Gastræa* is the simple cell-layer from which the

differentiated sense-organs of all Intestines and, therefore, of all Vertebrates, originally arising from the consideration that necessary superficial parts of the body, those immediately in contact with the outer world, could have accomplished their functions. This should be justified in conjecturing *à priori* that all sense also owe their origin to the same principle. The most important part of the nervous system develops from the outermost germ-layer, the sensory layer; in part, directly from the brain, the foremost section of the nerve tube, after this has separated from the rest. Comparing the individual development of sense, we see that at first they make the simplest conceivable form: only that that wonderful perfect structure develops. Higher sense-organs eventually become more complex and the most complex mechanisms of sense. All organs of sense are, however, distributed in portions of the external skin-covering. Even these nerves are distributed. Even these nerves are homogeneous and undifferentiated in character. By division of labour, the various functions and "energies" of the different sensorial nerves develop simultaneously the simple terminal expansion of nerves in the skin-covering developed into complex organs.

The important bearings of these facts on the just appreciation of mental life will be seen. The whole philosophy of the future

Three Groups.	Sense-organs.	Sense-nerves.	Sense-functions.
A. Sense-organs in which the terminal expansions of the nerves are distributed in the outer skin-covering.	I. Skin-covering (outer skin, or <i>epidermis</i> , and leather - skin, or <i>corium</i>)	I. Skin nerves (<i>nervi cutanei</i>)	Sense of pressure Sense of warmth
	II. External sexual parts (<i>penis and clitoris</i>)	II. Sexual nerves (<i>nervi pudendi</i>)	
B. Sense-organs in which the terminal expansions of the nerves are distributed over inverted grooves of the outer skin-covering.	III. Mucous membrane of the mouth-cavity (tongue and palate)	III. Taste nerve (<i>nervus glosso-pharyngeus</i>)	4. Sense of taste
	IV. Mucous membrane of the nose-cavity	IV. Olfactory nerve (<i>n. olfactorius</i>)	5. Sense of smell
C. Sense-organs in which the terminal expansions of the nerves are distributed over vesicles separated from the external skin-covering.	V. Eye	V. Sight nerve (<i>n. opticus</i>)	6. Sense of sight
	VI. Ear	VI. Ear-nerve (<i>n. acusticus</i>)	7. Sense of hearing

Of the developmental history of the lower organs of sense I have but little to say. The development of the skin-covering, which is the organ of the sense of pressure (or of touch) and of warmth, we have already traced (p. 120). I need only add that in the leather skin (*corium*) of all higher Vertebrates, innumerable microscopic sense-organs develop, the direct relations of which to the sensations of pressure or resistance, of warmth and of cold are not yet ascertained. These organs, in or upon which sensitive skin-nerves terminate, are the so-called "Pacini bodies" and the "Pacini bodies," named after their discoverer.

everer, Pacini. Similar bodies are also found in the organs of the sexual sense, in the penis of the male and in the clitoris of the female; these are processes of the integument, the development of which we shall consider presently, in connection with that of the other organs of generation. In the development of the organ of taste, the tongue and the palate, we will also consider presently, in connection with that of the intestinal canal, to which these parts belong. On one point, however, I will now call particular attention, viz., the mucous membrane of the tongue and palate, in which the taste-nerve terminates, is also in its origin a portion of the external skin-covering. For, as we found, the entire mouth-cavity originates, not as a part of the actual intestinal canal, but as a groove-like inversion of the external skin (vol. i. p. 338). Its mucous membrane, therefore, is formed, not from the intestinal layer, but from the skin-layer, and the taste-cells on the upper surface of the tongue and palate arise, not from the intestinal-glandular layer, but from the skin-sensory layer.

This is equally true of the mucous membrane of the organ of smell, the nose. The history of the development of this sense-organ is, however, of far higher interest. Although the human nose, externally viewed, seems simple and single, yet in Man, as in all higher Vertebrates, it consists of two perfectly distinct halves, of a right and a left nasal cavity. These two cavities are entirely separated by a vertical partition, so that the passage into the right nasal cavity lies only through the right nostril, and into the left cavity only through the left nostril. Posteriorly the two nasal cavities open separately through the two posterior nasal apertures into the head of the pharynx, so that the

pharynx may be entered without touching the cavity of the mouth. This is the passage by which air is usually inhaled, the mouth being shut, it enters the pharynx, and then passes through the windpipe into the lungs. Both the nasal and oral cavities are separated from the mouth-cavity by the horizontal bony palate roof, to the back of which the soft palate and the uvula is attached, like a hanging curtain. In the upper and hinder portion of both nasal cavities the olfactory nerve extends over the mucous membrane, and supplies lines these parts. This is the first pair of brain nerves which issue from the skull-cavity through the sieve-plate. Its branches extend partly over the partition wall and partly over the inner side-walls of the nasal cavities, which are attached the "shells," or spongy bones of the nose—complex bony structures. These "shells" are further developed in many of the higher Mammals and are present in Man. In all Mammals there are three of these "shells" in each of the two nasal cavities. The sensation of smell is produced by a current of air, containing odorous matters, passing over the mucous membrane of the cavities and there coming in contact with nerve-ends.

The peculiar characters which distinguish the olfactory organ of Mammals from that of lower Vertebrates are represented in Man. In all specific points the human nose exactly resembles that of the Catarrhine Apes, some of which indeed possess an entirely human external nose (see Fig. of the Nose-ape, Fig. 202, p. 175). The first rudimentary olfactory organ in the human embryo does not, however, show any signs of the fine form of the future catarrhine nose. Indeed, it first appears in the same form as in Fishes; in the form of two simple

grooves in the skin of the upper surface of the head. In Fishes two of these mere blind nose-pits are found in the upper surface of the head; sometimes they are situated on the back, near the eyes, sometimes near the snout, or, again, near the mouth-opening (Fig. 191, *n*, p. 113). They are covered by mucous membrane in folds, over which the end branches of the olfactory nerves spread.

In this its original condition the double nose of all *Aphirhina* (p. 101) is entirely unconnected with the primitive mouth-cavity. The connection, however, begins to

FIG. 231.—Head of a Shark (*Scyllium*), from the ventral side: *m*, mouth opening; *o*, nose grooves, or pits; *r*, nasal furrow; *n*, nose-flap in its normal position; *n'*, nose-flap turned up. (The dots are openings of mucous glands.) (After Gegenbaur.)



appear even in some Primitive Fishes (*Selachii*); a superficial skin-furrow extends on each side from the nose-groove down to the adjacent corner of the mouth. This furrow, the nasal channel, or furrow (Fig. 231, *r*), is of great significance. In many Sharks (e.g., *Scyllium*) a special process of the frontal skin, the nasal flap, or "inner nasal process," overlaps the nasal furrow (*n*, *n'*). Opposite to this the outer edge of the furrow rises and forms the "outer nasal process." In Dipneusta and Amphibia these two nasal processes meet over the furrow and coalesce, thus forming a canal, the "nasal canal." There is now a passage from the external nasal groove through this canal directly into the mouth-

cavity, which latter was developed independently of the upper lip groove. In the Dipneusta and the lower Amphibia the internal opening of the nasal canal lies well forward (near the lips); in the higher Amphibia it lies further back. In the three highest vertebrate classes, the Amniota, the primary mouth-cavity is separated by the formation of the horizontal palate roof into two perfectly distinct cavities, the superior (or secondary) nasal cavity, and the inferior (or secondary) mouth-cavity. The nasal cavity is further separated by the vertical partition into two distinct cavities, into a right and a left nasal cavity.

Comparative Anatomy thus still shows us similarities in the ascending series of the double-nostrilled Vertebrates from Fishes up to Man, all the various stages of development of the nose which the very highly developed organ of the higher Mammals has passed through successively in the different periods of its tribal history. The first rudiment of the organ of smell in the embryo of Fishes, and in that of all the higher Mammals, makes its appearance in the same entirely simple form which is retained throughout life by the nose of Fishes. At a very early stage, while no trace of the characteristic facial structure is yet visible, a pair of small grooves appear on the sides of the head, and before the primitive mouth-cavities were first discovered by Baer, and by him properly named "olfactory grooves" ("Riechgruben," Figs. 233, n). These primitive nasal grooves are quite separated from the primitive mouth-cavity, or mouth indentation, which, as we found, likewise makes its appearance as a groove-like indentation of the external skin-covering at the front of the blind anterior extremity of the intestinal

This pair of nasal grooves, as well as the single mouth groove (Fig. 235, *m*), is lined by the horn-plate. The



FIG. 232.



FIG. 233.



FIG. 234.



FIG. 235.



FIG. 236.

FIGS. 232, 233.—Head of an embryonic Chick, on the third day of incubation: 232, from the front; 233, from the right side. *n*, Nose-rudiment (olfactory grooves); *l*, eye-rudiment (sight-grooves); *g*, ear-rudiment (auditory grooves); *v*, fore-brain; *gl*, eye-slits; *o*, upper jaw process; *u*, lower jaw process of the first gill arch.

FIG. 234.—Head of an embryonic Chick, on the fourth day of incubation, from below: *n*, nose-groove; *o*, upper jaw process of the first gill arch; *u*, lower jaw process of the same; *k''*, second gill-arch; *sp*, choroidal fissure of the eye; *s*, throat (pharynx).

FIGS. 235, 236.—Two heads of embryonic Chicks: 235, at the end of the fourth day; 236, at the end of the fifth day of incubation. The letters as in g. 234. Additional letters are *in*, inner, and *an*, outer nasal process; *nf*, nasal furrow; *st*, frontal process; *m*, mouth-cavity. (After Koelliker.)

All these figures are proportionately enlarged.

original separation of the nasal groove from the mouth groove is, however, soon interrupted, for the frontal process (Fig. 235, *st*, Rathke's "Nasenfortsatz der Stirnwanne") is immediately formed above the mouth groove. Right and left the edges of this process project in the form of lateral processes: these are the inner nasal processes or nasal flaps (Fig. 235, *in*). On each side, opposite to the eye, arises a parallel ridge between the eye and the nasal groove. These ridges are the outer nasal processes (Rathke's "Nasendächer," Fig. 235, *an*). Between the inner and outer nasal process a channel-like depression thus extends on each side from the nose groove toward the mouth groove (*m*), and this channel is, of course, the same nasal furrow or channel which we found in the Shark (Fig. 231, *r*). Between the two parallel edges of the inner and the outer nasal processes bend towards each other and coalesce above the nasal channel, the latter becomes a small tube—the primitive "nasal canal." In this stage of its Ontogeny, therefore, the nose of Man and of all other Amnion Animals consists of two small narrow tubes—the "nasal canals"—leading from the outer surface of the frontal skin into the simple primitive mouth-cavity. This transient condition resembles the permanent condition of the nose in Dipneustic Amphibia. (Cf. Plate I., Frontispiece, with explanation.)

Specially significant in the modification of the open nasal channel into the closed nasal canal, is a plug-shaped process, the upper jaw process, which extends from below up to the lower extremities of both the nasal processes on each side, and unites them. This is the upper jaw process (Figs. 232, *o*, Plate I., *o*). Below the mouth groove lie the gill openings, which are separated from one another by the gill opercula.

I., VI., and VII., *k*). The first of these gill arches, at the most interesting to us, which we may call the *h*, develops the jaw-skeleton of the mouth (Plate I., *u*). The first process first grows out from the base of the front *h*: this is the upper jaw process. The first gill-arch develops a cartilage on its inner side, called after its discoverer, "Meckel's cartilage," on the outer surface of the lower jaw forms (Figs. 232, *u*, 236, *u*). The upper process forms the principal part of the entire framework of the upper jaw, viz., the palate bone and the wing bone. On the outer side the upper jaw bone, in the narrower sense, arises, while the middle portion of the upper jaw bone, the twist jaw (intermaxillary bone) develops the anterior portion of the frontal process. (See development of the face in Plate I.)

In the further characteristic development of the face in the higher vertebrate classes, the two upper jaw processes are of the highest importance. From them proceeds the palate roof, the important horizontal partition which divides the primitive mouth-cavity, separating it into two quite distinct cavities. The upper cavity, into which the two nasal cavities open, now develops into a nasal cavity—a respiratory air passage and an olfactory organ. The lower cavity, on the other hand, forms, by itself, a permanent secondary mouth-cavity (Fig. 237, *m*)—the cavity for the food passage and the organ of taste. Both the upper cavity and the lower taste-cavity open at the back into the pharynx. The palate roof, separating these two cavities, is formed by the coalescence of two lateral portions of the horizontal plates of the two upper jaw processes (Fig. 237, *p*). When these do not perfectly

adhere in the middle line, the result is a permanent longitudinal cleft, through which there is an open passage from the mouth-cavity directly into the nasal cavity. The




FIG. 237.—Diagrammatic transverse section through the mouth and nose cavity. When the palate-plates (*p*) separate the original mouth-cavity into the lower secondary mouth-cavity (*m*) and the upper nasal cavity, the latter is parted by the vertical partition wall of the nose (*e*) into two halves (*n, n*). (After Gegenbaur.)

called "wolf's jaws" are thus caused. The "hare-lip" or "split lip" is a slighter degree of this arrested development.¹¹⁶

Simultaneously with the horizontal partition of the palate roof, a vertical wall by which the single nasal cavity is divided into two, a right and a left cavity, develops (Fig. 237, *n, n*). This vertical partition of the nose is formed by the middle part of the frontal process: this gives rise by ossification to the vertical lamella or sieve bone (cubiform plate), and below the great vertical bony partition wall—the "plough-share" (*vomer*), as front to the twixt-jaw (*os intermaxillare*). Goethe was the first to show that in Man, just as in all the other Sk Animals, the twixt-jaw appears as an independent bone between the two halves of the upper jaw. The vertical partition wall of the nose finally coalesces with the horizontal palate roof. The two nasal cavities are now as entirely separate from one another as from the secondary mouth-cavity. These three cavities open, however, at the bottom into the pharynx, or jaw-cavity.

double-nostrilled nose has now attained the structure characteristic of Man in common with all other Mammals. Further development is very easily intelligible: it is to the formation of internal and external processes in the walls of both nasal cavities. Within the cavities are the "nose shells," spongy bony structures, over which the olfactory mucous membrane spreads. The first brain nerve, the olfactory nerve, with its delicate branches, passes



238, 239.—Upper part of the body of a human embryo (16 mm. in length) during the sixth week: Fig. 238, from the left side; Fig. 239, from the front.

The origin of the nose in two lateral halves, separate, is still plainly visible. The nose and mouth are disproportionately great in comparison with the rest of the face, especially with the lower lip. (Hillman.)

240.—Face of a human embryo of eight weeks. (Hillman.) Cf. Frontispiece, Plate I. Fig. M—



from the large brain through the roof of both nasal cavities into the cavities, and extends over the olfactory mucous membrane. At the same time, by inversion of the mucous membrane, the minor cavities of the nose, which afterwards filled with air, and which communicate directly with the two nasal cavities, arise (frontal cavities, cavities in the sphenoid bone, jaw cavities, etc.). In this special mode of development they occur only in Mammals.¹⁷¹

The external nose is not developed until long after these essential internal parts of the olfactory organ have been formed. The first trace in the human embryo appears at the end of the second month (Figs. 238-240). The human embryo during the first month shows that originally there is no trace of the external nose. It afterwards grows out from the anterior nasal portion of the primitive mouth. The form of nose which is characteristic of Man does not appear till a period far later. Much stress is usually laid on the shape of the external nose as a noble characteristic occurring exclusively in Man; but there are Apes which have very human noses, as, for instance, the *Nasalis* already mentioned. On the other hand, the external shape of the nose, the fine shape of which is so extremely important to the beauty of the facial structure, possesses in certain races of Man a shape anything but beautiful. In Apes the external structure of the nose remains undeveloped. Especially remarkable is the important fact already mentioned that it is only in the Apes of the Old World, in the *rhinopithecines*, that the nasal partition wall (*septum*) remains small as it is in Man; in Apes of the New World it is considerably enlarged at the base, so that the nostrils open out (Platyrrhini, p. 175).

TABLE XXX.

SYSTEMATIC SURVEY OF THE CHIEF PHYLOGENETIC STAGES OF THE
HUMAN NOSE.*First Stage: Nose of the earlier Primitive Fishes.*

The nose is formed by a pair of simple skin-grooves (nose-pits) in the outer surface of the head (like those which are now permanently retained by the lower Selachians).

Second Stage: Nose of the more recent Primitive Fishes.

Each of the two blind nasal grooves becomes connected by a furrow (nasal-furrow) with one end of the mouth (as is yet permanently the case in the higher Selachians).

Third Stage: Nose of the Dipneusta.

The two nasal furrows change, in consequence of the coalescence of their edges, into closed canals (primary nose-canals), which open at their front ends, within the soft edges of the lip, into the primary mouth-cavity; as is yet permanently the case in the Dipneusta and the earlier lower Amphibia (Saxobranchia).

Fourth Stage: Nose of Amphibia.

The inner openings of the nasal canals penetrate further back into the primary mouth-cavity, so that they are surrounded by hard bony portions of the jaw (as is yet permanently the case in the higher Amphibia).

Fifth Stage: Nose of the Protamnia.

The primitive mouth-cavity, into which both nasal canals open, separates, in consequence of the formation of a horizontal partition (the palate-roof), into an upper nasal cavity and a lower (secondary) mouth-cavity. The formation of the spongy bones of the nose commences (as in the earlier Amnion Animals).

Sixth Stage: Nose of the earlier Mammals.

The simple nose-cavity separates, in consequence of the development of a vertical partition wall (the "plough," vomer), into two distinct nose-cavities, each of which is occupied by one of the nasal canals (as is yet the case in all Mammals). The spongy nose-bones differentiate.

Seventh Stage: Nose of the more recent Mammals.

Within both nose-cavities the development of the spongy bones proceeds farther, and an external nose begins to form.

Eighth Stage: Nose of the Catarhine Apes.

The internal and the external nose attain the full development exclusively characteristic of Catarhine Apes and of Man.

The history of the development of the eye is equally remarkable and instructive. For although the eye, owing to its exquisite optical arrangement and wonderful structure, is one of the most complex and most nicely adapted organs, yet it develops, without a preconceived design, from a very simple rudiment in the outer skin-covering.




FIG. 241.—The human eye in transverse section: *a*, protective membrane (*sclerotica*); *b*, horn membrane (*cornea*); *c*, outer membrane (*conjunctiva*); *d*, circular veins of iris; *e*, vascular membrane (*choroidea*); *f*, ciliary muscle; *g*, *corona ciliaris*; *h*, rainbow membrane (*iris*); *i*, optic nerve (*n. opticus*); *k*, anterior limit of the retina; *l*, crystalline lens (*lens crystallina*); *m*, inner cover of the horn membrane (water membrane, *membrana Descemeti*); *n*, pigment membrane (*pigmentosa*); *o*, retina; *p*, "petits-canal"; *q*, yellow spot of the retina. (After Helmholtz.)

When fully developed, the human eye is a globular capsule (the eyeball, *bulbus*, Fig. 241). This lies in th

ny orbit of the skull, surrounded by protective fat and motor muscles. The greater part of this eyeball is occupied by a semi-fluid, clear gelatinous substance, the vitreous body (*corpus vitreum*). The crystalline lens (fig. 241, *l*) is embedded in the anterior surface of the vitreous body. It is a lentil-shaped, bi-convex, transparent body—the most important of the light-refracting media of the eye. Among these media is, in addition to the lens and vitreous body, the aqueous humour (*humor aqueus*, at *a* in Fig. 241), in front of the lens. These three pellucid, light-refracting media—the vitreous body, the crystalline lens, and the aqueous humour—by which the rays of light, incident on the eye, are refracted and concentrated, are enclosed in a firm globular capsule consisting of several distinct membranes, comparable with the concentric layers of an onion. The outer and thickest of these forms the white protective membrane of the eye (*sclerotica*, *a*). It consists of firm, compact white connective tissue. In front of the lens a circular, very convex, transparent plate, resembling a watch glass, is inserted in the white protective membrane; this is the horny membrane (*cornea*, *b*). On its outer surface the horny membrane is covered by a very thin coating of outer skin (*epidermis*); this coating is called the connecting membrane (*conjunctiva*); it extends from the horny membrane over the inner surface of both eyelids—the upper and lower folds of skin which on closing the eyes are drawn together over them. At the inner corner of our eye there is, as a sort of rudimentary organ, the remnant of a third (inner) eyelid, which, as the “nictitating membrane,” is highly developed in the lower vertebrates (vol. i. p. 110). Below the upper eyelid are lodged

the tear-glands, the secretion of which keeps the surface of the eye smooth and clean.

Directly under the protective membrane is a delicate dark-red, highly vascular membrane, the vascular membrane (*choroidea, e*), and within this the retina (*o*), which is a dilatation of the optic nerve (*i*). This latter is the second brain nerve. It extends from the "centre of vision" (the second brain-bladder) to the eye, penetrates the coats of this, and then extends, as the retina, between the vascular membrane (*choroidea*) and the vitreous body (*corpus vitreum*). Between the retina and the vascular membrane lies another very delicate membrane, which is commonly, but wrongly, considered as part of the retina. This is the black pigment membrane (*pigmentosa, l. n. pigmenti, n*), or the "black carpet" (*tapetum nigrum*). It consists of a single layer of beautiful hexagonal cells accurately joined together and filled with black pigment granules. This pigment membrane lines, not only the inner surface of the actual vascular membrane, but also the inner surface of its anterior muscular prolongation, which, as a circular ring-like membrane, covers the edge of the iris and prevents the penetration of lateral rays. This is the well-known "rainbow membrane" (*iris, h*), which is differently coloured in different persons (blue, gray, brown, etc.). The "rainbow membrane" is the limit towards the front of the vascular membrane. The round hole in the iris is the pupil, through which the rays of light pass into the interior of the eye. Where the iris proceeds from the edge of the actual vascular membrane, the latter is much thickened and forms a beautiful ciliated crown (*corona ciliaris, g*), which surrounds the edge of the lens with about seventy and many smaller rays.

the embryo of Man, as in that of all other Amphibians, two pear-shaped vesicles grow out laterally, at a very early period, from the foremost part of the first brain-ventricle (Fig. 223, *a*, p. 218). These bladder-like protuberances are the primary eye-vesicles. At first they are directed laterally and forward, but they soon make their way further backward, so that after the specialization of the five brain-ventricles, they lie at the base of the twixt-brain. The spaces within the two pear-shaped vesicles, which attain a considerable size, communicate through their stalks with the cavity of the twixt-brain. Their covering is formed by the outer skin-covering (horny and leather-plate). Where, on each side, the latter comes directly in contact with the most curved portion of the primary eye-vesicles, a thickening (*l*) arises, and at the same time a groove-like indentation (*o*) in the horn-plate (Fig. 224, 1). This groove, which we will call the lens groove, deepens into a closed sac, the thick-walled lens vesicle (2, *l*), on the fact that the edges of the groove coalesce above



42.—Eye of an embryonic Chick in longitudinal section (1, of a chick sixty-five hours of incubation; 2, of a somewhat older germ; 3, of a chick four days old): *h*, horn-plate; *o*, lens groove; *l*, lens (in 1, as a part of the epidermis, while in 2 and 3 it has separated); *x*, point of the horn-plate at the point from which the lens separated; *b*, lens body; *r*, retina; *u*, pigment membrane. (After Remak.)

it. Exactly as the medullary tube originally separates from the outer germ-layer does this lens-sac separate from the horn-plate, in which it originated. The space within the sac is afterwards entirely filled by the cells of its thick wall and the solid crystalline lens is thus formed. The latter is therefore, purely a formation of the epidermis. Together with the lens the small fragment of the leather-plate (cornea) lying below the lens separates from the outer skin-covering. This small piece of the leather-skin very soon forms a highly vascular sac round the lens (*capsula vasculosa lens*). Its anterior portion at first covers the pupillary orifice and is then known as the pupillary membrane (*membrana pupillaris*). Its back portion of the same membrane is called the "*membrana capsulo-pupillaris*." This "vascular capsule, which merely serves to nourish the growing lens, afterwards entirely disappears. The later, permanent capsule contains no vessels, and is a structureless secretion of the lens cells.

As the lens thus separates from the horn-plate and grows inward, it must necessarily indent the adjacent primary eye-vesicles from without (Fig. 242, 1-3). This process may be compared to the inversion of the germ-layer vesicle (*blastula*), which in the *Amphioxus* and many low animals gives rise to the gastrula (vol. i. p. 192). In both instances the inversion of one side of the closed vesicle proceeds until finally the inner, inverted portion touches the outer, uninverted portion of the wall of the vesicle, so that the cavity disappears. Just as in the gastrula the former part changes into the intestinal layer (*entoderma*) and the latter into the skin-layer (*exoderma*), so in the inverted primary eye-vesicle the retina develops from the f

ner) part (Fig. 242, *r*), and the black pigment membrane from the latter (the outer, uninverted part). The hollow of the primary eye-vesicle changes into the optic nerve.

The lens (*l*) which enacts so important a part in this inverting process of the primary eye-vesicle, lies at first directly upon its inverted part, that is, on the retina (*r*). Very soon, however, the two separate, a new body, the vitreous body (*corpus vitreum*, *gl*), coming in between them. While the lens-sac is detaching itself, and the primary eye-vesicle is being inverted from without, another inversion simultaneously proceeds from beneath—from the superficial portion of the skin-fibrous layer, *i.e.*, from the leather-plate of the head. At the back of the lens and below it, a ledge-like process of the leather-plate arises (Fig. 243, *g*), which inverts the primary eye-vesicle (now shaped like a cup) from below, and presses in between the lens (*l*) and the retina (*r*) thus the primary eye-vesicle assumes the form of a hood. The opening of this hood, answering to the face, is covered by the lens; but the opening, through which the neck could pass, answers to the indentation through which the leather-skin passes in between the lens and the retina (the inner wall of the hood). The space within this secondary eye-vesicle is almost filled by the vitreous body, which answers to the head wrapped in this hood. The hood itself, properly speaking, double: the inner hood itself is the retina, and the outer one, directly surrounding the former, is the pigment membrane. The comparison with a hood renders this process of inversion, which is sometimes hard to explain, more clearly understood. The rudiment of the vitreous body (*corpus vitreum*) is at first very incon-

siderable (Fig. 243, *g*), and the retina disproportionately thick. As the former expands, the latter becomes much thinner, till at last the retina appears only as a very delicate




FIG. 243.—Horizontal transverse section through the eye of a human embryo of four weeks; 100 times enlarged (after Koelliker): *t*, lens (the dark wall of which is equal to the diameter of the central cavity); *g*, vitreous body (connected with the leather-plate by a stalk, *g'*); *r*, vascular loop (penetrating through the stalk (*g'*) into the vitreous body behind the lens); *r*, retina (inner, thicker, inverted lamella of the

primary eye-vesicle); *a*, pigment membrane (outer, thinner, uninverted lamella of the same); *h*, intermediate space between the retina and the pigment membrane (remnant of the cavity of the primary eye-vesicle).

coat of the thick, almost globular vitreous body, which fills the greater part of the secondary eye-vesicle. The outer layer of the vitreous body changes into a highly vascular capsule, the vessels of which afterwards disappear.

The slit-like passage through which the rudiment of the vitreous body grows from below in between the lens and the retina, of course causes a break in the retina and the pigment-membrane. This break, which appears on the inner surface of the vascular membrane as a colourless streak, has been inaptly called the choroidal cleft, though the true vascular membrane is not cleft at all at this point (Fig. 234, *sp*, 235, *sp*, p. 243). A thin process of the vitreous body passes inward on the under surface of the optic nerve, which it inverts in the same way as the primary eye-vesicle was inverted. The hollow cylindrical optic nerve (the stalk of

primary eye-vesicle) is thus transformed into a channel, growing downward. The inverted lower surface attaches to the uninverted upper surface of the hollow stalk, so that the hollow space within the stalk, forming the communication between the cavity of the twist-brain and of the primary eye-vesicle, now disappears. The two edges of the channel now grow downward toward each other, forming the band-like process of the leather-plate, and lying beneath it. Thus this process now lies within the folds of the solid secondary optic nerve. It develops into a mass of connective tissue carrying the central blood-vessel (retina (*vasa centralia retinæ*)).

An entirely fibrous covering, the fibrous capsule of the eye, now finally forms round the outside of the secondary optic nerve and its stalk (the secondary optic nerve). It arises from the head-plate, from that part of the skin-layer which immediately encloses the eye-vesicle. The fibrous covering takes the form of a completely-closed sac, which surrounds the whole ball of the eye, and the outer side of this, grows in between the lens and the head-plate. The globular wall of the capsule soon separates, by division of the surface, into two distinct membranes. The inner membrane becomes the choroidea, or vascular layer; and it forms the ciliated crown (*corona ciliaris*) and iris. The outer membrane, on the other hand, becomes the white enveloping, or protective membrane (*sclerotica*), and in front, forms the transparent horny membrane (cornea). The rudiments of all the essential parts of the eye are now formed, and its further development is only in the complex differentiation and combination of several parts.

TABLE XXXI.

Systematic Survey of the Development of the Human Eye.

I. Systematic Survey of those parts of the Human Eye which develop from the first of 6 Secondary Germ-layers, the Skin-sensory Layer.			
A. Products of the Marrow-plate	1. Stem of the primary eye-vesicle	1. Optic nerve	<i>Nervus opticus</i>
	2. Inner (inverted) part of the primary eye-vesicle	2. Retina	<i>Retina</i>
	3. Outer (uninverted) part of the primary eye-vesicle	3. Screen, or pigment-coat	<i>Pigmentosa (as pigment)</i>
B. Products of the Horn-plate	4. Vesicle separated from the horny plate	4. Crystalline lens	<i>Lens crystallina</i>
	5. Outer epidermic skin	5. Connective membrane	<i>Conjunctiva</i>
	6. Inverted portions of the epidermic skin	6. Tear-glands	<i>Glandula lacrym</i>
II. Systematic Survey of those parts of the Human Eye which develop from the second of 6 Secondary Germ-layers, the Skin-fibrous Layer.			
C. Products of the Leather-plate	7, 8. Ledge-like process of the corium on the lower side of the primary eye-vesicle	7. Vitreous body	<i>Corpus vitreum</i>
		8. Vascular membrane of the vitreous body	<i>Capsula vasculosa corporis vitrei</i>
	9. Continuation of the corium process	9. Central vessels of the retina	<i>Vasa centralia retinae</i>
	10. Pupillary membrane, with its capsule	10. Vascular membrane of the lens	<i>Capsula vasculosa lentis crystallinae</i>
D. Products of the Skull-plate	11. Folds of the leather skin (corium)	11. Eyelids	<i>Palpebrae</i>
	12, 13. Vascular membrane of the eyeball (<i>capsula vasculosa bulbi</i>)	12. Vascular membrane	<i>Choroides</i>
		13. Rainbow membrane	<i>Iris</i>
	14, 15. Fibrous membrane of the eyeball (<i>capsula fibrosa bulbi</i>)	14. Protective membrane	<i>Sclerotica</i>
		15. Horny membrane	<i>Cornes</i>

The most important fact in this remarkable process of development is the circumstance that the optic nerve, retina, and the pigment-membrane originate from a part of the brain, from a protuberance of the twixt-brain, while the crystalline lens, the most important refracting medium, develops from the outer skin (*epidermis*). From the outer skin—the horny lamina—originates also the delicate connecting membrane (*conjunctiva*) which afterwards envelopes the outer surface of the eyeball. The tear-ducts proceed, as branched processes, from the conjunctiva (g. 214, p. 202). All the other parts of the eye originate from the skin-fibrous layer; the vitreous body and the ocular lens-capsule from the leather-plate, the choroid together with the iris, and the protective membrane (*sclerotica*) with the horny membrane (*cornea*) from the head-plates.

The outer protective organs for the eye, the eyelids, are merely simple folds of skin, which, in the human embryo, appear in the third month. In the fourth month the upper eyelid adheres to the lower, and the eye then remains covered by them till birth. (Plate VII. Fig. *M* III., *R* III., &c.) The two eyelids usually again separate shortly before birth, but sometimes not till after. Our skulled ancestors, in addition to these, a third eyelid, the nictitating membrane, which was drawn over the eye from the inner corner. Many Primitive Fishes (*Selachii*) and Ammonites yet retain this. In Apes and in Man it has atrophied, and only a small remnant of it exists in the inner corner of the eye as the "crescent-shaped fold," as a useless "rudimentary organ." (Cf. vol. i. p. 109.) Apes and Man have also lost the "Harder gland," opening below the nictitating membrane, which appears in other Mammals, and in Birds, Reptiles, and Amphibians.

The ear of Vertebrates develops in many important points similarly to the eye and nose, but yet in respects very differently.¹⁷³ The organ of hearing of a developed human being resembles that of other Man in all essential particulars, and is especially similar to that of Apes. As in the latter, it consists of two principal parts: an apparatus for the conveyance of sound (external and internal ear) and an apparatus for producing the sensation of sound (internal ear). The outer ear opens in the ear-shell (a)



FIG. 244.—Auditory organ of man (left ear, seen from the front; size): *a*, ear-shell; *b*, external ear-canal; *c*, drum, or tympanic membrane; *d*, cavity of drum; *e*, ear-trumpet; *f*, *g*, *h*, the three ear bonelets (*f*, *h*, *g*, anvil; *h*, stirrup); *i*, ear-pouch (*utricle*); *k*, the three semi-circular canals; *l*, ear-sac (*sacculus*); *m*, snail (*cochlea*); *n*, auditory nerve.

auris), situated at the side of the head (Fig. 224, *a*). This is the outer ear-canal, which is usually about an inch long. It leads to the inside of the head (*b*). The inner end of the

It is closed by the well-known tympanic membrane or drum (*tympanum*); a thin membrane of oval form (*c*), placed in a vertical position, but slightly inclined. This membrane separates the outer ear-canal from the so-called cavity of the drum (*carum tympani*). This is a small cavity enclosed in the petrous part of the temporal bone, which is filled with air and connected by a special tube with the mouth-cavity. This tube is somewhat longer, but much narrower than the outer ear-canal; it leads inward and upward in an oblique direction from the inside wall of the tympanum and opens behind the inner nostrils (or Choana) into the upper part of the cavity of the throat (*pharynx*). This canal is called the Eustachian tube (*tuba Eustachii*).

It equalizes the pressure of the air in the tympanic cavity, and the outer atmospheric air which enters by the ear canal. Both the Eustachian tube and the tympanic cavity are lined by a thin, mucous membrane, which is a direct continuation of the mucous membrane of the throat. Within the tympanic cavity are the three bonelets of the ear, which, from their characteristic shape, are called the hammer, the anvil, and the stirrup (Fig. 244 *f, g, h*). The hammer (*f*) lies furthest outward, just within the tympanic membrane; the anvil (*g*) is wedged in between the two others, above the hammer, and further in than the hammer; and, lastly, the stirrup (*h*) lies next to the anvil toward the inside, and touches with its base the outer wall of the internal ear, or the auditory sac. All these parts of the middle and external ear belong to the sound-conducting apparatus. Their principal office is to convey the waves of sound from without through the thick side-wall of the head, to the internal ear. In Fishes these parts are entirely unre-

presented. In them, the sound-waves are conveyed directly through the wall of the head itself to the internal ear.

The inner apparatus, that which produces the sensation of sound, receiving the sound-waves thus conveyed to it consists in Man, as in all other Vertebrates (with the single exception of the *Amphioxus*), of a closed auditory sac filled with fluid, and of an auditory nerve, the ends of which are distributed over the wall of this sac. The vibrations of the waves of sound are conveyed by that medium to the nerve-ends. In the auditory fluid (*endolymph*), which fills the labyrinth, and opposite the places at which the auditory nerves enter, are some small stones, composed of a mass of microscopic calcareous crystals (*otoliths*). The organs of hearing of most Invertebrates have essentially the same construction. In them, also, it usually consists of a closed sac filled with fluid, containing otoliths, and having the auditory nerve distributed over its wall. But while in Invertebrates the auditory vesicle is usually of a simple spherical or oval form, in all Amphirrhina, on the contrary, that is, in all Vertebrates above the Fishes and Man, it is distinguished by a very characteristic and singular form known as the auditory labyrinth. This thin membranous labyrinth is enclosed in a bony envelope of the same form, the osseous labyrinth (Fig. 245), which lies within the petrous bone of the skull. The labyrinth in all Amphirrhina is divided into two sacs. The larger sac is called the auditory pouch (*utricle*), and has three curved appendages called the semi-circular canals (*c, d, e*); the smaller sac is called the auditory sac (*sacculus*), and is connected with the larger by a peculiar appendage, which in Man and the higher Mammals is distinguished by a spiral form, like the shell of a snail.

ace is called the "snail" (*cochlea*, *b*). On the thin wall of this delicate membranous labyrinth, the auditory nerve, which passes from the after-brain to the labyrinth, is distributed in a very complex manner. It divides into two main branches, the nerve of the cochlea, and the nerve of the vestibule, for the remaining part of the labyrinth. The former seems specially to determine the quality of the sound heard, the latter its quantity. The nerve of the cochlea

FIG. 245.—The bony labyrinth of the human ear (right side): *a*, vestibule; *b*, cochlea; *c*, upper semi-circular canal; *d*, posterior semi-circular canal; *e*, anterior semi-circular canal; *f*, fenestra ovalis; *g*, fenestra rotunda. (From Meyer.)



is us the pitch and quality of sounds, the nerve of the vestibule their strength.

The first rudiment of this extremely complex organ of hearing is very simple in the human embryo, as in those of all other Skulled Animals (*Craniota*); it is a groove-like depression of the outer skin (*epidermis*). At the back of the head, near the after-brain, at the upper end of the second hind-opening, a little wart-like thickening of the horn-plate arises on each side (Figs. 246, *A*, *fl*; 248, *g*). This deepens into a small groove, and separates from the outer-skin, just as does the lens of the eye. (Cf. p. 253.) A small vesicle filled with fluid, the primitive ear-vesicle, is thus formed on each side, immediately below the horn-plate of the back of the head; this is also called the "primary labyrinth" (Plates VI. and VII.). As this separates from its original site, the horn-plate, and grows inward and downward in the skull, it changes from a globular to a pear-shaped form (Figs. 246, *B*, *lv*; 249, *o*). The outer part has

elongated into a thin stalk, which at first opens outward in a narrow canal. (Cf. Fig. 137, *f*, vol. i. p. 382.) This is called the appendage of the labyrinth *recessus labyrinthi*, Fig. 246, *lr*.




FIG. 246.—Development of the ear-labyrinth of a Chick, in five consecutive stages (*A-E*) (cross-sections through the rudimentary skull): *e*, ear-groove; *lv*, ear-vesicle; *lr*, labyrinth appendage; *c*, rudiment of the cochlea; *csp*, hind semi-circular canal; *cse*, outer semi-circular canal; *jr*, jugular vein. (After Reissner.)

FIGS. 247, 248.—Head of an embryonic Chick, on the third day of incubation: 247 in front, 248 from the right; *n*, rudimentary nose (olfactory groove); *l*, rudimentary eye (ocular groove); *g*, rudimentary ear (auditory groove); *v*, fore-brain; *gl*, eye-slit; *o*, process of the upper jaw; *u*, process of the lower jaw of the first gill-arch. (After Koelliker.)

FIG. 249.—Primitive brain of human embryo of four weeks, in vertical section, and the left half observed from within: *v, z, m, h, n*, the five grooves of the skull cavity in which the five brain bladders are situated (fore, twin, mid, hind, and after brains); *o*, primary, pear-shaped auditory vesicle (showing through); *a*, eye (showing through); *no*, optic nerve; *p*, canal of the hypophysis; *t*, central skull-pieces. (From Koelliker.)

lower Vertebrates, this develops into a peculiar cavity filled with calcareous crystals, which in some Primitive fishes (*Selachii*) remains permanently open, and opens directly on the skull (*ductus endolymphaticus*). In Mammals, on the contrary, the appendage of the labyrinth degenerates into a blind pouch. In these, it is of interest only as a rudimentary structure, which has no longer any physiological significance. A useless remnant traverses the osseous wall of the petrous part of the temporal bone in the form of a narrow canal, and is called the aqueduct of the vestibule (*aquæductus vestibuli*).

Only the inner and lower part (extended like a bladder) of the detached ear-vesicle develops into the differentiated and extremely complex structure which is afterwards known as the "secondary labyrinth." This vesicle separates at an early stage into an upper, larger section, and a lower, smaller section. The former gives rise to the ear-pouch (*utriculus*) with the three semi-circular canals; from the lower part proceeds the ear-sac (*sacculus*) with the "snail" (*cochlea*, Fig. 246, c). The three semi-circular canals originate as simple pocket-like processes from the ear-pouch (Fig. 246, E, *cse* and *csp*). In the centre of each of these processes, the two walls coalesce, and separate themselves from the utricle, while their extremities still communicate with its cavity. In all Double-nostrils (*Amphirhina*) there are three semi-circular canals, as in Man, while of the Single-nostrils the Lampreys have but two, and the Myxinoides one (p. 103). The highly-developed structure of the "snail" (*cochlea*), which is one of the most delicate and admirable products of adaptation in the mammalian body, originally develops very simply as a bottle-like process from the ear-sac (*sacculus*). As Hasse has shown, the

various stages in its ontogenetic development still exist permanently side by side in the ranks of the lower Vertebrates.¹⁷⁸ Even in Monotremes the snail-like spiral curving of the cochlea is not present; it is exclusively characteristic of the other Mammals and Man.

The auditory nerve (*nervus acusticus*), or the eighth brain-nerve,—one of the main branches of which distributes itself over the “snail” (*cochlea*), the other over the other parts of the labyrinth,—is, as Gegenbaur has shown, the sensory dorsal branch of a spinal brain-nerve, the motor ventral branch of which is the motor nerve of the facial muscles (*nervus facialis*). Phylogenetically it has, therefore, originated from an ordinary skin-nerve, and is, consequently, of wholly different origin from the optic and olfactory nerves, which represent the two direct processes of the brain. In this respect the organ of hearing differs essentially from the organs of sight and of smell. The auditory nerve originates from the cells of the head-plate; therefore, from the skin-fibrous layer. From this also develop all the membranous, cartilaginous, and bony coverings of the ear-labyrinth.

The development of the apparatus for the conveyance of sound, situated in the middle and external ear of Mammals, is entirely distinct from that of the apparatus of auditory sensation. It must be regarded, phylogenetically as well as ontogenetically, as an independent, secondary formation, which only afterwards connects itself with the primary internal ear. Its development is, however, not less interesting, and is equally clearly explained by Comparative Anatomy. In all Fishes, and in the yet lower Vertebrates, there is no special apparatus for the conveyance of sound,

TABLE XXXII.

SYSTEMATIC SURVEY OF THE CHIEF STAGES IN THE DEVELOPMENT
OF THE HUMAN EAR.I. *First Stage.*

The auditory nerve is an ordinary sensitive skin-nerve, which, during the differentiation of the horn-plate, appears at a certain point on the skin of the head.

II. *Second Stage.*

The differentiated place of the horn-plate, at which the auditory nerve appeared, forms a small special auditory groove in the skin, which has an outer orifice in the appendage called the "labyrinth."

III. *Third Stage.*

The auditory groove has detached itself from the horn-lamina, and forms a small closed auditory vesicle filled with fluid. The "labyrinth-appendage" becomes rudimentary (*Aquæductus vestibuli*).

IV. *Fourth Stage.*

The auditory vesicle differentiates into two connected parts, the ear-pouch (*utricleus*) and the ear-sac (*sacculus*). Each of the two vesicles receives a special main branch of the auditory nerve.

V. *Fifth Stage.*

Three semi-circular canals grow from the ear-pouch (as in all *Amphibia*).

VI. *Sixth Stage.*

The "snail" (*cochlea*) grows from the ear-sac in Fishes and Amphibia; it is very insignificant, and is only developed as an independent part in the Amniota.

VII. *Seventh Stage.*

The first gill-opening (the blow-hole of Selachians) changes into the tympanic cavity and the Eustachian tube; the former is externally closed by the tympanic membrane (Amphibia).

VIII. *Eighth Stage.*

The small bones of the ear (*ossicula auditus*) (the hammer (*malleus*) and anvil (*incus*) from the first gill-arch, the stirrup (*stapes*) from the second) develop from parts of the first and second gill-arches.

IX. *Ninth Stage.*

The external ear is developed, together with the bony ear-canal. The shell of the ear is pointed and movable (as in most lower Mammals).

X. *Tenth Stage.*

The ear-shell, with its muscles, becomes disused and a rudimentary organ. It is no longer pointed, but, on the contrary, has a curved rim with a small ear-flap (as in Anthropoid Apes and Men).

TABLE XXXIII.

Systematic Survey of the Development of the Human Ear.

I. Survey of the parts of the Internal Ear. (Apparatus perceptive of sound.)				
A. Products of the Horn-plate	{	1. Stalk of the primary ear-vesicle	1. Aqueduct of the vestibule (<i>Ductus endolymphaticus</i>)	<i>Aqueductus vestibuli</i> <i>s. Rocessus labyrinthi</i>
		2, 3. Upper part of the primary ear-vesicle	2. Ear-pouch 3. Three semi-circular, or curved canals	<i>Utriculus</i> <i>Canales semi-circulares</i>
		4, 5. Lower part of the primary ear-vesicle	4. Ear-sac 5. "The small"	<i>Sacculus</i> <i>Cochleæ</i>
B. Products of the Head-plate	{	6. Auditory nerve	6. Auditory nerve	<i>Nervus acusticus</i>
		7. Bony covering of the membranous labyrinth	7. Osseous labyrinth	<i>Labyrinthus osseus</i>
		8. Bony covering of the whole internal ear	8. "The stony bone"	<i>Os petrosum</i>
II. Survey of the parts of the Intermediate and External Ear. (Apparatus for the conveyance of sound.)				
C. Products of the first Gill-opening	{	9. Inner part of the first gill-opening	9. Eustachian tube	<i>Tuba Eustachii</i>
		10. Central part of the first gill-opening	10. Tympanic cavity (Interior of the drum)	<i>Cavum tympani</i>
		11. Closed part of the first gill-opening	11. Tympanic membrane (Head of the drum)	<i>Membrana tympani</i>
D. Products of the first two Gill-arches	{	12. Upper part of the second gill-arch	12. Stirrup bonelet (First of the ear)	<i>Stapes</i>
		13. Upper part of the first gill-arch	13. Anvil (Second bonelet of the ear)	<i>Incus</i>
		14. Central part of the first gill-arch	14. Hammer (Third bonelet of the ear)	<i>Malleus</i>
E. Product of the Head-plate	{	15. Tympanic circle (<i>Annulus tympanicus</i>)	15. Bony outer auditory passage	<i>Meatus ossæus auditorius</i>
F. Product of the Skin-covering	{	16. Circular membranous fold at the closed part of the first gill-opening	16. Ear-shell	<i>Concha auris</i>
			17. Rudimentary ear-muscles	<i>Musculi conchæ</i>

no external and middle ear; in these animals there is only a labyrinth, an internal ear, situated within the skull. The tympanic membrane, its cavity, and all the connected parts are unrepresented. The middle ear first develops in the Amphibian class, in which a tympanic membrane, a tympanic cavity, and an Eustachian tube are first found. All these essential parts of the middle ear develop from the first gill-opening, with its surrounding parts, which in the Primitive Fishes (*Selachii*) remains through life as an open blow-hole, situated between the first and second gill-arches. In the embryos of higher Vertebrates it closes in the centre, the point of concrescence forming the tympanic membrane. The remaining outer part of the first gill-opening is the rudiment of the outer ear-canal. From the inner part originates the tympanic cavity, and further inward, the Eustachian tube. In connection with these, the three bonelets of the ear develop from the first two gill-arches; the hammer and anvil from the first, and the stirrup from the upper end of the second gill-arch.¹⁷⁴

Finally, as regards the external ear, the ear-shell (*concha auris*), and the outer ear-canal, leading from the shell to the tympanic membrane—these parts develop in the simplest way from the skin-covering which borders the outer orifice of the first gill-opening. At this point the ear-shell rises in the form of a circular fold of skin, in which cartilage and muscles afterwards form (Fig. 238, p. 247). This organ is also limited to Mammals. Among them, it is originally wanting only in the lowest division, in the Beaked Animals, (*Monotrema*). In the others, on the contrary, it appears in very different stages of development and partly also of atrophy. The ear-shell has atrophied in most aquatic

Mammals. Most of these have even lost it entirely; this is so, for example, in the Sea-cows and Whales, and most Seals. On the other hand, in the great majority of Pouched Animals (*Marsupialia*) and Placental Animals (*Placentalia*), the ear-shell is well developed, receives and concentrates the waves of sound, and is provided with a highly-developed muscular apparatus, by means of which it can be turned freely to all sides, and at the same time can be changed in form. Every one must have noticed how strongly and freely our domestic Mammals, Horses, Cows, Dogs, Rabbits, etc., can "prick" their ears, erect them and turn them in different directions. Most Apes yet retain the power of doing this, and our ancient Ape progenitors could also do it. The more



FIG. 250.—Rudimentary ear-muscles on the human skull: *a*, upward muscle (*m. attollens*); *b*, forward muscle (*m. attrahens*); *c*, backward muscle (*m. retrahens*); *d*, larger muscle of the helix (*m. heliciis major*); *e*, smaller muscle of the helix (*m. heliciis minor*); *f*, muscle of the tragus (*m. tragicus*); *g*, muscle of the antitragus (*m. antitragicus*). (After H. Meyer.)

recent Ape ancestors, common to Men and to the Anthropoid Apes (Gorilla, Chimpanzee, etc.), discontinued the habit of moving their ears, and hence the motor muscles gradually

became rudimentary and useless. We still, however, possess them (Fig. 250). A few individual men can even move their ears forward or backward a little by the use of the forward muscle (*b*) and the backward muscle (*c*); and by long practice these motions can be gradually increased. On the other hand, no man is able to erect the ear-shell by the upward muscle (*a*), or to change its form by the little inner muscles of the ear (*d, e, f, g*). These muscles, which were very useful to our ancestors, have become entirely unimportant to us. This is equally true of Anthropoid Apes.

We also share only with the higher Anthropoid Apes—the Gorilla, Chimpanzee, and Orang—the characteristic form of our human ear-shell, especially the rolled edge, the helix, and the ear-flap. The lower Apes, like all other Mammals, have pointed ears without the helix, and without ear-flaps. Darwin has, however, shown that in some men a short, pointed process, not occurring in most individuals, is perceptible at the upper part of the folded rim of the ear. In some few individuals, this process is very well developed. It can only be explained as the remnant of the original point of the ear which, in consequence of the folding of the edge of the ear, has been bent forward and inward. (Cf. the similarly folded ear in the embryo of the Pig and Cow, Plate VII. Fig. *H* III. and *C* III.) On carefully comparing the ear-shells of Man and of the various Apes in this particular, we find that they form a connected series of retrograde steps. In the common catarrhine ancestors of the Anthropoids and of Man, this retrogression began with the folding down of the ear-shell. In consequence of this, the ear-edge was formed on which that significant corner appears, the last trace of the free prominent point of the ear

in our older Ape ancestors. Thus it is possible even here, with the help of Comparative Anatomy, to trace this human organ from the similar but more highly-developed organ of the lower Mammals, with certainty. At the same time, Comparative Physiology shows us that this organ is of more or less high physiological value to the latter, while in Anthropoids and Man it is a useless rudimentary organ. Men with their ears cut off can hear as well as they did before. The conveyance of sound is not affected by the loss of the ear-shell. This explains the great diversity in the form and size of the ear-shell in different persons; it shares this high degree of variability with other rudimentary organs.¹⁷⁵

CHAPTER XXII.

DEVELOPMENT OF THE ORGANS OF MOTION.

The Motive Apparatus of Vertebrates.—These are constituted by the Passive and Active Organs of Motion (Skeleton and Muscles).—The Significance of the Internal Skeleton of Vertebrates.—Structure of the Vertebral Column.—Formation and Number of the Vertebrae.—The Ribs and Breast-bone.—Germ-history of the Vertebral Column.—The Notochord.—The Primitive Vertebral Plates.—The Formation of the Metamera.—Cartilaginous and Bony Vertebrae.—Intervertebral Discs.—Head-skeleton (Skull and Gill-arches).—Vertebral Theory of the Skull (Goethe and Oken, Huxley and Gegenbaur).—Primitive Skull, or Primordial Cranium.—Its Formation from Nine or Ten Coalescent Metamera.—The Gill-arches (Ribs of the Head).—Bones of the Two Pairs of Limbs.—Development of the Five-toed Foot, adapted for Walking, from the Many-toed Fin of the Fish.—The Primitive Fin of the Selachians (*Archipterygium* of Gegenbaur).—Transition of the Pinnate into the Semi-pinnate Fin.—Atrophy of the Rays or Toes of the Fins.—Many-fingered and Five-fingered Vertebrates.—Comparison of the Anterior Limbs (Pectoral Fins) and the Posterior Limbs (Ventral Fins).—Shoulder Girdle and Pelvis Girdle.—Germ-history of the Limbs.—Development of the Muscles.

"In forming his estimate of my entire theory, the reader may begin with the details and examine the fundamental facts on which I base my conclusions. But it is equally necessary to connect the detached facts, and estimate their bearing on the whole. He who in the world of organisms sees only disconnected existences, in which some organic similarities appear as

accidental coincidences, will remain a stranger to the results of the investigation; not merely because he does not comprehend the conclusions, but principally because the significance of the facts on which they are grounded, escapes him. A fact in itself is no more a scientific result than a mere collection of facts is a science. That which makes a science of these facts, is their combination by that organizing mental faculty which determines the relations of the facts to each other."—KARL GROENBACH (1872).

AMONG those features of the organization which are specially characteristic of the vertebrate tribe as such, the peculiar arrangement of the motive apparatus, or "locomotorium," undoubtedly occupies a principal place. As in all the higher animals, the active organs of motion, the muscles, form the most important part of this apparatus; these are the fleshy bands which, by means of their peculiar contractibility, of their power of contracting and shortening, move the various parts of the body, and thus change the position of the entire body. The arrangement of these muscles is, however, entirely peculiar in Vertebrates, and differs from the arrangement common to all Invertebrates.

In most lower animals, especially in Worms, we find that the muscles form a simple, thin flesh-layer immediately below the outer skin-covering. This "skin-muscle pouch" is most intimately connected with the skin itself, and the same feature occurs in the tribe of the Soft-bodied Animals (*Mollusca*). In the great group of the Articulated Animals (*Arthropoda*), in the Crab, Spider, Centipede, and Insect classes, we also find a similar feature, but with the difference that in these the skin-covering forms a hard coat of mail or an inflexible skin-skeleton, formed of chitine, and often of carbonated chalk. This outer chitinous coat of mail is jointed in a great variety of ways both on the trunk and

on the limbs of Articulated Animals, and the muscular system, the contractile fleshy bands of which are attached to the inside of the chitinous tubes, is correspondingly pointed in an extremely varied manner. The case is exactly reversed in Vertebrates. In these alone an internal hard skeleton develops; an inner cartilaginous or bony frame to which the fleshy muscles are externally attached, and in which they find a firm support. This bony frame forms a combined lever-apparatus, a passive apparatus of motion. The hard parts of this, the arms of the lever, or the bones, are moored against each other by the active movable muscular bands, as by hawsers. This admirable locomotive apparatus, and especially its firm central axis, the vertebral column, is quite peculiar to Vertebrates, on account of which the whole group has long been called that of Vertebrates.

This internal skeleton, notwithstanding the similarity of its first rudiment, has, however, developed so variously and characteristically in the different vertebrate classes, and in the higher classes forms so complex an apparatus, that Comparative Anatomy finds one of its richest mines in this feature. This was recognized as long ago as the beginning of the century by the older Natural Science, which at once seized these very welcome materials with peculiar pleasure. That science also, which is now called in the higher and more philosophical sense, "Comparative Anatomy," has reaped its richest harvest from this field. The Comparative Anatomy of the present day has studied the skeleton of Vertebrates more thoroughly, and revealed the laws of its formation more successfully, than has been the case with any other system of organs of the animal body. Here the well-known and oft-quoted passage, in which Goethe

summed up the general result of his investigations in Morphology is especially appropriate :

" All forms have a resemblance ; none is the same as another,
And their chorus complete points to a mystical law." *

Now that, by the Theory of Descent, we have discovered this " mystical law," have solved this " sacred enigma," and that we can explain the similarity of forms by Heredity and their dissimilarity by Adaptation, we can find a new weapon in the whole rich arsenal of Comparative Anatomy which defends the truth of the Theory of Descent more powerfully than the comparison of the internal skeleton of the various Vertebrates. We may, therefore, explain *à priori* that such comparison is of special importance in our History of the Evolution of Man. The inner vertebral skeleton is one of those organs as to the Phylogeny which Comparative Anatomy affords us conclusions more important and deeper than those to be gained from its Ontogeny.¹⁷⁶

More than any other system of organs, the internal skeleton of Vertebrates, when studied comparatively, clearly and immediately impresses the observer with the necessity of the phylogenetic connection between these allied and very varied forms. A thoughtful comparison of the bony frame of Man with that of other Mammals, and of that again with that of lower Vertebrates, is alone sufficient to afford conviction of the true tribal relationship of the Vertebrates. All the separate parts of which this bony frame is composed appear in other Mammals, in a greater or less degree.

* " Alle Gestalten sind ähnlich, doch keine gleicht der andern,
Und so deutet der Chor auf ein geheimes Gesetz."

variety of forms indeed, but yet in the same characteristic arrangement and relative position; and if the comparison of the anatomical conditions of the skeleton is carried out below Mammals, we can prove that a direct and uninterrupted connection exists throughout between these various forms which are apparently so utterly unlike, and can finally be traced from a most simple, common, fundamental form. These facts alone must fully convince every adherent of the Theory of Development that all Vertebrates, including Man, must be traced from a single common parent-form, from a Primitive Vertebrate; for the morphological features of the inner skeleton, and of the muscular system which stands in the closest correlative relations to it, are of such a kind that it is quite impossible to conceive a polyphyletic origin, a descent from several different root-forms. It is impossible, on mature reflection, to accept the theory that the vertebral column with its various appendages, or the skeleton of the limbs with their variously differentiated parts, could have originated on several occasions during the course of the earth's history, and that, consequently, the various Vertebrates must be referred in various lines of descent from Invertebrates. Indeed, it is exactly in this point that Comparative Anatomy and Ontogeny irresistibly drive us to the monophyletic conclusion, that the human race is a very recent offshoot of the same great single trunk, from branches of which all other Vertebrates have also sprung.

In order to obtain a view of the outlines of the development of the human skeleton, we must first take a general survey of its arrangement in the developed Man. (Cf. Table XXXIV. and Fig. 251, the human skeleton from the

TABLE XXXIV.

Systematic Survey of the Arrangement of the Human Skeleton.

A. Central Skeleton, or Axial Skeleton. Spine.			
A.a. Vertebral Bodies and Upper Arches.		A.b. Lower Vertebral Arches.	
1. Skull (Cranium)	1 a. Pre-vertebral skull	1. Products of the gill-arches	Products of branchiostele
	2 b. Vertebral skull		
	7 Neck vertebrae		
2. Vertebral column (columna vertebralis)	12 Chest "	2. Ribs and breast-bone	Costal et sternum
	5 Hip "		
	5 Vertebrae of the sacrum		
	4 " " " tall (coccyx)		
B. Bones connecting the Extremities.			
B.a. Bones connecting the Anterior Limbs: Bones of the Shoulder.		B.b. Bones connecting the Lower Limbs: Bones of the Pelvis.	
1. Shoulder-blade	Scapula	1. Intestinal bone	Os ileum
(2. Primitive key-bone	Procoracoides †)	2. Pubic bone	Os pubis
(3. Raven bone	Coracoides †)	3. Hip-bone	Os ischi
4. Collar-bone, or key-bone	Clavicula		
C. Jointed Skeleton of the Limbs.			
C.a. Skeleton of the Fore Limbs.		C.b. Skeleton of the Hind Limbs.	
I. FIRST DIVISION: UPPER ARM.		I. FIRST DIVISION: THIGH.	
1. Upper arm bone	Humerus	1. Thigh-bone	Femur
II. SECOND DIVISION: LOWER ARM.		II. SECOND DIVISION: LEG.	
2. Spoke-bone	Radius	2. Shin-bone	Tibia
3. Ell-bone	Ulna	3. Calf-bone	Fibula
III. THIRD DIVISION: HAND.		III. THIRD DIVISION: FOOT.	
III. A. Wrist		III. Ankle	
Original parts.	Modified parts.	Original parts.	Modified parts.
a. Radical	= Scaphoideum	a. Tibial	} = Astragalus
b. Intermedium	= Lunatum	b. Intermedium	
c. Ulnar	= Triquetrum	c. Fibular	
d. Central	= Intermedium †)	d. Central	= Naviculare
e. Carpal I.	= Trapezium	e. Tarsal I.	= Cuneiform I
f. " II.	= Trapezoides	f. " II.	= " II
g. " III.	= Capitatum	g. " III.	= " III
h. " IV. + V.	= Hamatum	h. " IV. + V.	= Cuboides
III. B. Palm of the Hand	Metacarpus (5)	III. B. Sole of the Foot	Metatarsus (5)
III. C. Five Fingers	Digiti	III. C. Five Toes	Digiti
(14 bones)	Phalanges)	(14 bones)	Phalanges)



FIG. 251.



FIG. 252.



FIG. 253.—
Human vertebral
column (in an up-
right position;
from the right
side). (After H.
Meyer.)

right side (without arms); Fig. entire skeleton from the front.) as in all other Mammals, the skeleton is primarily distinguishable into the axial skeleton, or spine, and the skeletal appendages, or the bony frame of the body. The spine consists of the vertebrae and of the skull; the latter being a specially modified anterior part of the axial skeleton. The ribs are the appendages of the thoracic column; the tongue-bone (*os linguae*), the lower jaw, and the other products of the first gill-arches, are those of the skull. The skeletons of the two pairs of limbs, the forelimbs, are composed of two parts: of the bony frame of the appendages, or the prominent extremities, and of the intrinsic skeleton, by which the limbs are attached to the vertebral column. The girdle of the arms (or forelimbs) is the shoulder girdle; the girdle skeleton of the legs (or the hind limbs) is the pelvic girdle.

The bony vertebral column in man is the *columna vertebralis*, or *vertebral column*, Fig. 253) is composed of thirty or thirty-four circular pieces of bone, the vertebrae, which lie one behind the other (one above the other in the usual upright position of man). These bones (*vertebrae*) are separated from each other by elastic

the intervertebral discs (*ligamenta intervertebralia*), and at the same time, are connected by joints, so that the entire vertebral column forms a firm and solid axis, which is however, flexible and elastic, capable of moving freely in all directions. In the various regions of the trunk, the vertebrae differ in form and connection, so that the following are distinguished in the human vertebral column, beginning from above: seven neck-vertebrae, twelve breast-vertebrae, five lumbar-vertebrae, five cross-vertebrae, and four to five tail-vertebrae. The uppermost, those directly in



FIG. 254.—Third neck-vertebra of man.

FIG. 255.—Sixth breast-vertebra of man.

FIG. 256.—Second lumbar-vertebra of man.

in contact with the skull, are the neck-vertebrae (Fig. 254), and are distinguished by a hole found in each of the two lateral processes. There are seven neck-vertebrae in man, as in nearly all other Mammals, whether the neck is long, as in the Camel and the Giraffe, or short, as in the mole and the Hedgehog. The fact that the number of these neck-vertebrae is always seven,—and there are but few exceptions (explicable by adaptation),—is a strong argument for the common descent of all Mammals; it can only be accounted for as a strict transmission from a common

parent-form, from some Promammal which had seven vertebrae. If each animal species had been a distinct type, it would have been far more to the purpose to have furnished the long-necked Mammalia with a larger, and the short-necked with a smaller number of neck-vertebrae. The neck-vertebrae are immediately followed by those of the thorax, which, in Man and most other Mammals, number twelve or thirteen (usually twelve). Attached to the side of each breast-vertebra (Fig. 255) is a pair of ribs—curved processes of bone lying in and supporting the wall of the thorax. The twelve pairs of ribs, with the connecting intercostal muscles and the breast-bone (*sternum*) constitute the breast-body (*thorax*, Fig. 252, p. 279). In this elastic and yet firm thorax lie the double lung, and between the two halves of this, the heart. The chest-vertebrae are followed by a short but massive section of the vertebral column, formed by five large vertebrae. These are the lumbar-vertebrae (Fig. 256), which bear no ribs and no perforations in their lateral processes. Next come the sacral-vertebrae, which are fused into a single cross-bone (*sacrum*), which is inserted between the two halves of the pelvic girdle. This cross-bone consists of five fused and amalgamated cross-vertebrae. Last comes a rudimentary tail-vertebral column, the rump-bone (*caudal vertebrae*). This bone consists of a varying number (usually four, rarely three or five) of small aborted vertebrae; it is a useless rudimentary organ, retaining no physiological significance either in Man or in the Tail-less Apes or Anthropoids. (Cf. Figs. 204–208.) Morphologically it is, however, very interesting, as affording incontrovertible evidence of the descent of Man and of Anthropoids from Long-tailed Apes. For this assumption affords the only possible

explanation of this rudimentary tail. In the human embryo, indeed, during the earlier stages of germ-history, the tail projects considerably. (Cf. Plate VII. Fig. *M* II., and Figs. 123, *s*, 124, *s*, vol. i. p. 370.) It afterwards becomes adherent, and is no longer externally visible. Yet traces of the aborted tail-vertebræ, as well as of the rudimentary muscles, which formerly moved them, persist throughout life. According to the earlier anatomists the tail in the female human being has one vertebra more than that of the male (four in the latter, five in the former).¹⁷⁷

Number of Vertebra in various Catarkini.		Neck Verte- bræ.	Chest or tho- racic Verte- bræ.	Lum- bar Verte- bræ.	Cross or sacral Verte- bræ.	Tail Verte- bræ.	Total.
Tail- less	Man (Fig. 208)	7	12	5	5	4	33
	Orang (Fig. 205)	7	12	5	4	5	33
	Gibbon (Fig. 204)	7	13	5	4	3	32
	Gorilla (Fig. 207)	7	13	4	4	5	33
	Chimpanzee (Fig. 206)	7	14	4	4	5	34
Tailed	Mandril (<i>Mormon choras</i>)	7	13	6	3	5	34
	Drill (<i>Mormon leucophæus</i>) ...	7	12	7	3	8	37
	Rhesus (<i>Inuus rhesus</i>)	7	12	7	2	18	46
	Sphinx (<i>Papio sphinx</i>)	7	13	6	3	24	53
	Simpai (<i>Semnopithecus melus</i>)	7	12	7	3	31	60

The number of vertebræ in the human vertebral column is usually thirty-three in all; but it is an interesting fact that this number frequently varies, one or another vertebra failing, or a new, supernumerary vertebra inserting itself. Not unfrequently, also, a rib, capable of free motion, forms on the last neck-vertebra or on the first lumbar-vertebra, so that thus there are thirteen breast, and six neck, or four lumbar vertebræ. In this way contiguous vertebræ in the different sections of the vertebral column may replace each

other. On the other hand, the above comparison of number of vertebræ in different tail-less and tailed Cats shows considerable fluctuations in these numbers even in this one family.¹⁷⁸

To understand the history of the development of the human vertebral column, we must now study the formation and combination of the vertebræ in somewhat greater detail. The main outline of each vertebra is that of a signet ring (Figs. 254-256). The thicker part, which faces the ventral side, is called the body of the vertebra, and it forms a disc of bone; the thinner part forms a semi-circular arch, the vertebral arch, which is turned toward the dorsal side of the body. The arches of all the consecutive vertebræ are connected by thin ligaments (*ligamenta intercruralia*) the space enclosed by them all in common forms a long canal. In this spinal, vertebral canal lies, as we have seen, the portion of the central nervous system, the spinal marrow. The front part of this, the brain, is enclosed in the skull cavity, and hence the skull itself is merely the anterior section of the vertebral column, modified in a peculiar manner. The base or ventral side of the bladder-shaped brain-cavity was originally formed by a number of coalescent vertebral bodies, the amalgamated upper vertebral arches of which formed the arched or ventral side of the skull.

While the firm, massive vertebral bodies constitute the true central axis of the skeleton, the dorsal arches surround and enclose and protect the central marrow. Analogous arches also develop on the ventral side as a protection for the thoracic and abdominal viscera. These inferior or ventral vertebral arches, proceeding from the ventral side of the vertebral bodies, form a canal in many low Vertebræ.

ich are enclosed the large blood-vessels on the under face of the vertebral column—the aorta and the tail vein. Higher Vertebrates most of these inferior vertebral arches are lost or become merely rudimentary. But in the breast region of the vertebral column they develop into strong, independent bony arches, the ribs (*costæ*). The ribs are, in fact, merely large vertebral arches which have become independent, and have broken their original connection with the vertebral bodies. The gill arches, of which we have spoken so often, are of similar origin; they are actual head-ribs in the strictest sense—processes which have actually originated from the lower arches of the skull-vertebræ, and which correspond with the ribs. Even the mode of connection of the right and left halves of the arches on the ventral side is the same in both instances. The breast is closed in front by the intervention, between the upper ribs, of the breast-bone (*sternum*)—a single bone originating from two corresponding side-halves. The gill-plate is also closed in front by the intervention of a single piece of bone—the *copula lingualis*.

In now turning from this anatomical examination of the constitution of the vertebral column to the question of its development, I may, as regards the first and most important features in the evolution, refer the reader to the explanation already given of the germ-history of the vertebral column (chap. XII, vol. i. pp. 369–378). In the first place, it is necessary to recollect the important fact that in Man, as in all other Vertebrates, a simple, unarticulated cartilaginous rod at first occupies the place of the articulated vertebral column. This firm but flexible and elastic cartilaginous rod is the well-known notochord (*chorda dorsalis*). In the lowest Ver-

tebrate, the *Amphioxus*, this persists throughout life in a very simple form, and permanently constitutes the internal skeleton (Fig. 151, *i*, vol. i. p. 420; Plate XI. F). But even in the Mantle Animals (*Tunicata*), the invertebrate allies of Vertebrata, we find this same chord; transitorily in the transient larval tail of (Plate X. Fig. 5, *ch*); permanently in the Appendicularia (Fig. 162). The Mantle Animals, as well as the Vertebrates, have undoubtedly inherited the notochord from a common worm-like parent-form, and these primæval worm animals are the Chorda Animals (*Chordonia*, p. 91).

Long before any trace of a skull, limbs, etc., appears in the human embryo or in that of any of the higher Vertebrates, in that early stage when the whole body is represented





FIG. 257.—Portion of notochord (*chorda dorsalis*) of an embryo sheep: *a*, sheath; *b*, cells. (After Koelliker.)

by the lyre-shaped germ-disc—in the central line of this latter, directly under the primitive groove or medullary furrow, appears the simple *chorda dorsalis*. (See Fig. 84-87, vol. i. pp. 297, 298, surface view; Figs. 66-70, 89-93, transverse sections; Plates IV., V., *ch*.) As a cylindrical rod it traverses the longitudinal axis of the embryo and is equally pointed at both ends. The cells which compose the chord (Fig. 257) come, in common with all the other cells of the skeleton, from the skin-fibroblasts. They most resemble certain cartilage cells; a special "chordal tissue" is often said to exist; but this must not be regarded as more than a special form of cartilaginous tissue. In the early period the notochord envelopes itself in a strong sheath (*a*) as clear as glass, which is secreted by its

This perfectly simple, inarticulate, primary axial skeleton is soon replaced by an articulated, secondary skeleton, called the "vertebral column." On each side of the notochord the primitive vertebral bands or primitive vertebral plates (vol. i. p. 306, Fig. 92, *uv*) differentiate from the inner portion of the skin-fibrous layer. The inner part of these primitive vertebral bands, which immediately surrounds the notochord, is the skeleton-plate, or skeleton-tum (*i.e.*, the cell-layer forming the skeleton), which furnishes the tissue for the rudiments of the permanent vertebral column and of the skull. In the anterior half of the body the primitive vertebral plate remains a simple, continuous, unbroken layer of tissue, and soon expands into a thin-walled vesicle, which surrounds the brain; this is the notochordal skull. In the posterior half, on the contrary, the primitive vertebral plate breaks up into a number of homologous cube-shaped pieces, lying one behind the other; these are the several primitive vertebræ. The number of these is at first very small, but soon increases, as the animal grows in the posterior direction (Figs. 258-260, *uv*). The first and earliest primitive vertebræ are the foremost neck-vertebræ; the posterior neck-vertebræ then originate; then the anterior breast-vertebræ, etc. The lowest of the tail-vertebræ arise last. This successive ontogenetic growth of the vertebral column in a direction from front to rear may be explained phylogenetically by regarding the many-membered vertebrate body as a secondary product, which originated from an originally inarticulate parent-form by progressive metameric development, or articulation. Just as the many-membered Worms (Earth-worm, Leech) and the closely allied Arthropods (Crabs, Insects) originally



FIGS. 258-260.—Lyre-shaped germ-shield of a Chick, in three consecutive stages of development; seen from the dorsal side; enlarged about two times. Fig. 258, with six pairs of primitive vertebrae. The brain is a single bladder (*hb*). The spinal furrow from *x* remains wide open; behind *z*, it is much enlarged. *mp*, Marrow-plates; *sp*, side-plates; *y*, limit between the pharynx cavity (*sh*) and the head-intestine (*vd*). Fig. 259, ten pairs of primitive vertebrae. The brain has separated into three bladders: *v*, fore-brain; *m*, mid-brain; *h*, hind-brain; *c*, heart; *dv*, dorsal veins. The spinal furrow is still wide open (*z*). *mp*, Marrow-plates. Fig. 260, with sixteen pairs of primitive vertebrae. The brain has separated

five bladders; *v*, fore-brain; *z*, twist-brain; *m*, mid-brain; *h*, hind-brain; *n*, after-brain; *a*, eye-vesicles; *g*, ear-vesicles; *c*, heart; *dv*, yolk-sacs; *mp*, marrow-plate; *uw*, primitive vertebra.

Developed from an inarticulate worm-form by terminal budding, so the many-membered vertebrate body has originated from an inarticulate parent-form. The nearest tant allies of this parent-form are the Appendicularia (Fig. 162) and the Ascidian (Plate XI. Fig. 14).

As has been repeatedly pointed out, this primitive vertebral, or metameric structure has a very important bearing on the higher morphological and physiological development of Vertebrates. (Cf. vol. i. p. 346.) For the articulation is by no means confined to the vertebral column, but usually affects the muscular, nervous, vascular, and other stems. As is shown by the *Amphioxus*, the metameric structure appeared much earlier in the muscular than in the skeleton system. Each so-called primitive vertebra is in fact far more than the mere rudiment of a future vertebra. In each primitive vertebra exists the rudiment of a segment of the dorsal muscles, of a pair of spinal nerves, etc. Only the inner portion—that which lies directly next to the notochord and the medullary tube—is employed, the skeleton-plate, in the formation of actual vertebræ. We have already seen how these true vertebræ develop from the skeleton-plate of the primitive vertebræ or metamera. The right and left lateral halves of each primitive vertebra, originally separate, unite. The ventral edges, meeting below the medullary tube, surround the chord and thus form the rudiments of the vertebral bodies; the dorsal edges, meeting above the medullary tube, form the first rudiments of the vertebral arches. (Cf. Figs. 95-98, and Plate IV. Figs. 3-8.)

In all Skulled Animals (*Craniota*), most of the undifferentiated cells which originally constitute the skeleton-plate, afterwards change into cartilage cells



FIG. 261.—Three breast-vertebrae of a human embryo of eight weeks, in lateral longitudinal section: *v*, cartilaginous vertebral bodies; *li*, intervertebral discs; *ch*, notochord. (After Koeliker.)

secrete a firm, elastic "intercellular substance," and thus produce cartilage tissue. Like most other parts of the skeleton, the rudimentary vertebrae pass into a cartilaginous condition. In the higher Vertebrates, the cartilaginous tissue is afterwards replaced by the rigid bony tissue with its radiate bone-cells (Fig. 5, vol. i.). The original axis of the vertebral column, the notochord, is more or less completely surrounded by the cartilaginous tissue which vigorously round it. In lower vertebrates (*i.e.*, in Primitive Fishes) a more or less considerable portion of the notochord remains within the vertebral bodies. In Mammals, on the contrary, it disappears entirely. In the human embryo, even at the end of the second month, the notochord is seen only as a thin line which passes through the axis of the thick cartilaginous vertebral column (Fig. 261, *ch*). In the cartilaginous vertebral bodies themselves, which afterwards ossify, the thin remnant of the notochord (Fig. 262, *ch*) soon disappears entirely. A remnant remains, however, throughout life in the "intervertebral discs" which develop, from the skeleton-plate, between each pair of vertebral bodies (Fig. 263). In a new-born child, a large, pear-shaped cavity, filled with a gelatinous cell-mass, is visible in each intervertebral

(Fig. 263, a). This "gelatinous nucleus" of the elastic vertebral disc becomes less sharply defined, but persists throughout life in all Mammals, while in Birds and Reptiles, even the last remnant of the notochord vanishes.



FIG. 262.—A breast-vertebra of the same embryo in lateral cross-section: *cv*, cartilaginous vertebral bodies; *ch*, notochord; *pr*, square process; *a*, vertebral arch (upper); *c*, upper end of rib (lower arch). (After Koelliker.)

FIG. 263.—Intervertebral disc of new-born child in cross-section: *a*, remnant of the notochord. (After Koelliker.)

When the cartilaginous vertebræ afterwards ossify, the first deposit of bone-substance (the first "bone-nucleus") in the vertebral bodies is formed immediately round the remnant of the notochord, and soon completely displaces the latter. A special bone kernel or nucleus is then formed in each half of the cartilaginous vertebral arch. It is not till after birth that the ossification progresses so far that the three bone-nuclei approach each other. The two bony halves of the arch unite during the first year, but it is not till much later, till between the eighth and the twelfth year, that they unite with the bony vertebral body.

The bony skull (*cranium*), which must be regarded as

the foremost, peculiarly modified section of the vertebral column, develops in an exactly similar manner. Just as, in the spinal column, the vertebral canal envelopes and protects the dorsal marrow, so the skull forms a bony covering round the brain; and, as the brain is merely the anterior, peculiarly differentiated portion of the dorsal marrow, we might conclude on *a priori* grounds, that the bony envelope of the brain is a peculiar modification of that of the dorsal marrow. It is true, that if the developed human skull (Fig. 264) is considered by itself, it is impossible to understand how it can be merely the modified anterior portion of the vertebral column. It is a complex, capacious bony structure, consisting of no less than twenty bones, differing




FIG. 264.—Human skull, from the right side.

widely in form and size. Seven of these skull-bones constitute the spacious case which encloses the brain, and in which we distinguish the strong, massive floor of the skull (*basis cranii*) below, and the boldly arched roof of the skull (*fornix cranii*) above. The other thirteen bones form the "facial skull," which especially provides the bony envelopes of the higher sense-organs, and at the same time as the jaw-skeleton, encircles the entrance to the intestinal canal. The lower jaw (usually regarded as the twenty-first skull-bone) is jointed to the skull-floor, and behind this, embedded in the roots of the tongue, we find the tongue-bone, which, like the lower jaw, has originated from the gill-arches, together with a portion of the lower arch, which originally developed as "skull-ribs" from the ventral side of the skull-floor.

Although, therefore, the developed skull of the higher Vertebrates, in its peculiar form, its very considerable size, and its complex structure, seems to have nothing in common with ordinary vertebræ, yet the old comparative anatomists at the close of the eighteenth century correctly believed that the skull is originally merely a series of modified vertebræ. In 1790, Goethe "picked up out of the sand of the Jews' burying-ground among the downs near Venice, a dismembered skull of a sheep; he at once perceived that the face bones (like the three vertebræ of the back of the skull) are also derivable from vertebræ." And, in 1806, Oken (without knowing of Goethe's discovery), at Ilsestein, on the way to the Brocken, "found a beautifully bleached skull of a hind; the thought flashed through him, it is a vertebral column!"¹⁷⁹

For the last seventy years, this celebrated "Vertebral Theory of the Skull," has interested the most prominent zoologists; the most important representatives of Comparative Anatomy have exercised their ingenuity in attempting to solve this philosophical skull-problem; and the question has engaged attention in yet wider circles. It was not till 1872 that the solution was found, after seven years of labour, by the comparative anatomist, who, both in the wealth of his real empirical knowledge and in the profundity of his philosophic speculations, surpasses all other students of this science. Karl Gegenbaur, in his classic "Researches in the Comparative Anatomy of Vertebrates" (third part), showed that the skull skeleton of the Selachii is the only record which affords definite proof of the vertebral theory of the skull. Earlier comparative anatomists erred in starting from the developed mammalian skull, and

in comparing the several component bones with the separate parts of vertebræ; they supposed that in this way they could prove that the developed mammalian skull consisted of from three to six original vertebræ. The hindmost of these skull-vertebræ was, according to them, the occipital bone. A second and a third vertebra were represented by the sphenoid bone, with the parietal bones, and by the frontal bone, etc. The elements of anterior skull vertebræ were even supposed to exist in the face bones. In opposition to this view, Huxley first called attention to the fact that in the embryo this bony skull originally develops from a simple cartilaginous vesicle, and that in this simple cartilaginous "primitive skull" not the slightest trace of a constitution of vertebrate parts is visible. This is equally true of the skulls of the lowest and most ancient Skulled Animals (*Craniota*), the Cyclostomi and the Selachii. In these the skull retains throughout life the form of a simple cartilaginous capsule—of an inarticulate "primitive or primordial skull." If the older skull-theory, as it was accepted from Goethe and Oken by most comparative anatomists, were correct, then in these lowest Skulled Animals especially, and in the embryos of the higher Skulled Animals, the constitution of the "primitive skull" by a series of "skull-vertebræ" would be very clearly evident.

This simple and obvious consideration, first duly emphasized by Huxley, indeed overturns the famous "Vertebrate Theory of the Skull," as held by the older comparative anatomists. Yet the entirely correct fundamental idea holds good, *i.e.*, the hypothesis that the skull develops from the anterior portion of the spinal column by differentiation and peculiar modification, just as the brain develops from

the anterior portion of the dorsal marrow. But the true mode of empirically establishing this philosophic hypothesis is yet to be discovered; and this discovery we owe to Gegenbaur.¹⁸⁰ He was the first to employ the phylogenetic method, which, in this as in all morphological questions, leads most surely and quickly to the result. He showed that the Primitive Fishes (*Selachii*, Figs. 191, 192, p. 113), as the parent-forms of all Amphirhina, yet retain permanently in their skull-structure that form of primordial skull, from which the modified skull of the higher Vertebrates, and therefore that of Man, has developed phylogenetically. He also pointed out that the gill-arches of the *Selachii* show that their primordial skull was originally formed of a considerable number—at least nine or ten—primitive vertebræ, and that the brain-nerves, which branch from the base of the brain, entirely confirm this. These brain-nerves—with the exception of the first and the second pairs (the olfactory and the optic nerves)—are merely modified spinal nerves, and, in their peripheric distribution, essentially resemble the latter. The Comparative Anatomy of these brain-nerves is one of the strongest arguments for the newer vertebral theory of the skull.

It would lead us too far aside if we were to enter into the particulars of this ingenious theory of Gegenbaur, and I must content myself with referring to the great work already quoted; in it the theory is fully demonstrated by empirical and philosophical arguments. The same author has given a brief abstract in his "Outlines of Comparative Anatomy" (1874), the study of which it is impossible to recommend too highly. In this work Gegenbaur indicates as original "skull-ribs," or "lower arches of skull-

vertebræ," in the selachian skull (Fig. 265), the following pairs of arches: I. and II. are two lip cartilages, of which the anterior (*a*) consists only of an upper, and the inferior (*bc*) of an upper and a lower piece; III., the jaw-arch, which also consists of two pieces on each side,—viz. the primitive upper jaw (*os palato-quadratum*, *o*) and




FIG. 265.--Head skeleton of a Primitive Fish: *n*, nose-groove; *eth*, of the sieve-bone; *orb*, eye-cavity; *la*, wall of ear-labyrinth; *occ*, occipital region of the primitive skull; *cv*, vertebral column; *a*, front; *bc*, hind part of the jaw; *o*, primitive upper jaw (*palato quadratum*); *u*, primitive lower jaw; II., tongue-arch; III.-VIII., first to sixth gill-arches. (After Huxley.)

primitive lower jaw (*u*); IV., the tongue arch (II.), and V., six true gill arches, in the stricter sense of that (III.-VIII.). The anatomical features of these nine arches, or "lower vertebral arches," and of the nerves distributed over them, show that the apparatus is a simple, cartilaginous "primordial skull" of the Primitive Fishes originally develops from an equal number (nine at the least) of primitive vertebræ. The base of the skull is formed by the vertebral bodies; the roof of the skull by the upper vertebral arches. The coalescence and amalgamation of these into a single capsule is, however, so ancient,

their primordial separate condition now appears effaced by the action of the "law of abridged heredity," and is no longer demonstrable in the Ontogeny.

In the human primitive skull (Fig. 266), and in that of all higher Vertebrates, which has been modified, phylogenetically, from the primitive skull of the Selachii, five consecutive divisions are visible at a certain early period of development; these one might be tempted to refer to five

FIG. 266.—Primitive skull of human embryo of four weeks; vertical section, the left half seen from the inside: *v*, *z*, *h*, *n*, the five grooves in the skull cavity, in which lie the five brain-bladders (fore-brain, twixt-brain, mid-brain, hind-brain, after-brain); *o*, pear-shaped primary ear-vesicle; *a*, eye; *no*, optic nerve; *h*, canal of the hypophysis; *t*, central part of the cranial basis. (After Koelliker.)



original primitive vertebræ; they are, however, merely the result of adaptation to the five primitive brain-bladders, and, like the latter, they rather correspond to a larger number of metamera. The fact that the primitive vertebrate skull is a much modified and profoundly transformed organ, and by no means a primitive structure, is also evident in the circumstance that its rudiment, originally a soft membrane, commonly assumes the cartilaginous state only at its base and on the sides, while it remains membranous at the skull-roof. Here the bones of the later bony skull develop in the soft membranous rudiment as an external bony roof, without a previous intermediate cartilaginous state, as in the base of the skull. Thus a great part of the skull-bones originally developed as roof-bones from the

leather-skin (*corium*), and only secondarily, come into relations with the skull. How, in Man, this most simple primordial rudiment of the primitive skull develops, genetically, from the head-plates, and how, in the meantime, the anterior extremity of the notochord is enclosed the base of the skull, has already been explained. (vol. i. p. 378; Figs. 145 and 146, p. 393.)

The main features in the history of the development of the gill-arches, which must now be regarded as skull, has been told. Of the four original rudimentary gill-arches in Mammals (Plates I. and VII., Figs. 232–236, p. 243), the first lies between the primitive mouth-opening and the first gill-opening. From the base of this gill-arch the “upper process” develops, and this unites, in the manner already described, with the internal and the external nasal process on each side, and forms the chief parts of the upper jaw skeleton—palate-bones, wing-bones, etc. (Cf. p. 245 and 268.) The rest of the first gill-arch, now distinguished as the “lower process,” forms out of its base two ear bonelets—the hammer (*malleus*) and the anvil (*incus*); the rest of its mass becomes a long strip of cartilage, called, after its discoverer, “Meckel’s cartilage.” On the external surface of this cartilage ossifies, as a surface-bone (formed of cellular matter from the leather-plate), the permanent bony lower jaw. From the base of the second gill-arch in Mammalia originate the third ear bonelet, the stirrup (*stapes*), and from the subsequent parts, in order, the stirrup-muscle, the styloid process, the temporal bone, the styloid band, and the small horn of the tongue-bone. Finally, the third gill-arch becomes cartilaginous only at its anterior portion, and here, by the union of its two halves, is formed the body of the tongue-bone (or

dea) and its great horn on each side. The fourth gill-appears in the mammalian embryo only as a transient, elementary embryonic organ, and does not develop into final parts. Of the posterior gill-arches (the fifth and sixth pairs), which are permanent in the Primitive Fishes, none is visible in the embryo of higher Vertebrates. The first two have long been lost. The four gill-openings in the mammalian embryo are also only interesting as transient rudimentary organs, which soon disappear entirely by concretions. The first gill-opening (between the first and second arches) alone is of permanent importance; from it develops the drum, or tympanic cavity of the ear, and the Eustachian tube. (Cf. p. 269, and Plate I., with explanation.)

Not only did Gegenbaur, in his model "Researches into Comparative Anatomy of Vertebrates," first correctly explain the skull and its relation to the vertebral column, but he also first performed the no less weighty and interesting task of showing the phylogenetic derivation of the origin of the limbs in all Vertebrates from one primordial pair. Few parts of the body in the different Vertebrates are subjected, by adaptation to various circumstances, to such an infinite variety of modifications as the limbs, in respect of size, form, and special fitness for certain purposes, and we are now able to refer them all to one common hereditary form. Vertebrates are distinguishable as regards the structure of their limbs into three large main groups. The most and most ancient Vertebrates, the skull-less and jawless classes, like all their invertebrate ancestors, had no true limbs; this condition is yet represented in the *Amphioxus* and in the *Cyclostomi* (Figs. 189, 190). The second

main group consists of the two classes of true Fishes, the Dipneusta; in these, two pairs of lateral limbs, shape of many-fingered swimming-fins—one pair of p fins (the fore legs) and one pair of abdominal fins (hind are originally always present (Figs. 191, 192, Plate Finally, the third main group embraces the four vertebrate classes: Amphibia, Reptiles, Birds, and Ma in these the same two pairs of legs exist originally, the form of five-fingered feet. The digits or fing often fewer than five; sometimes, also, the feet are aborted. But the original parent-form of the entire had anteriorly and posteriorly five digits (Pentadac p. 123).

As regards the Phylogeny of the limbs, from Comparative Anatomy it appears, therefore, that the mities originated in the Fishes, in the Primitive (*Selachii*), and were transmitted from these to all Vertebrates (all the *Amphirhina*), first in the f many-fingered fins, and afterwards as five-finger (Figs. 267–272). The anterior extremity—the pect (or the fore leg)—is originally shaped precisely l posterior extremity—the ventral fin (or the hind le the one, as in the other, the true limb, externally nent, is distinguishable from the internal, concealed by which the limb is attached to the spinal column shoulder-girdle above, the pelvic girdle below.

The genuine primitive form of the paired limb existed in the most ancient of the Primitive Fishes the Silurian Period, occurs to this day in perfect pr tion in the ancient *Ceratodus*, and very curious Mud Australia (p. 119, Plate XII). In this, both the pecto

Ventral fin is a flat, oval paddle, in which we find a feathered or biserial cartilaginous skeleton (Fig. 267). This skeleton consists firstly of a strong, articulated fin-rod or "stem" (Fig. 267, *A B*), which extends from the base to the tip of the fin, and secondly, of a double row of thin, feathered rays (*rr*), which are attached to both sides of the central rod, like the pinnæ of a pinnate leaf. This primitive fin, first recognized by Gegenbaur, and by him called *Archipterygium*, is attached to the spinal column by means of a simple girdle in the shape of a cartilaginous

In some Sharks and Rays, especially when very young, the same primitive fin also occurs in a more or less modified form. But in most Primitive Fishes the fin is already partially modified, in that the rays on one side of the stem are partly or altogether lost, and are retained only on the other side (Fig. 268). Hence arises the half-feathered, or uniserial fish-fin, inherited by the other fishes from the chii (Fig. 269).

Gegenbaur first showed how the five-fingered leg of the Amphibia is developed from this uniserial fin (Fig. 270) and inherited by three classes of Amniota. In those Diplopoda which were the ancestors of the Amphibia, the fin rays on the other side of the stem also were gradually degraded in development, and were in a great measure lost (the light-colored cartilages in Fig. 269). Only the four lowest rays (labeled in Fig. 269) were retained; and these are the four digits of the foot (second to fifth digits). The first, great digit (toe), on the contrary, originated from the upper part of the fin-rod. From the middle and upper parts of this fin-rod developed the long main stem of the limbs




FIG. 267.




FIG. 268.




FIG. 269.




FIG. 270.




FIG. 271.



FIG. 272

267.—Bones of pectoral fins of *Ceratodus* (Archipterygium, or pinnate skeleton): *AB*, series of cartilaginous pieces forming the stem of the fin; *rr*, rays of the fin. (After Günther.)

268.—Bones of pectoral fin of an earlier Primitive Fish (*Acanthias*). The rays of the medial edge of the fin (*B*) have disappeared; only *R'* remain. *RR*, rays of the lateral edge of fin; *mt*, Metapterygium; *ms*, Mezoptyrgium; *p*, Propterygium. (After Gegenbaur.)

269.—Bones of pectoral fin of a more recent Primitive Fish, or *Elasmobranch*. The rays of the medial edge of the fin have entirely disappeared.

The shaded part on the right is that portion which develops into the fore-limb of higher Vertebrates (*b*, the three basal pieces of *mt*, Metapterygium; rudiment of the humerus; *ms*, Mezoptyrgium-Propterygium). (After Gegenbaur.)

270.—Bones of the fore-limb of an Amphibian: *h*, upper arm; *r, u*, lower arm (*r*, radius; *u*, ulna); *r, c, i, c, u*, root-bones of the first row (*r*, radial; *i*, intermediate; *c*, central; *u*, ulnary); 1, 2, 3, 4, 5, bones of the hand, second row. (After Gegenbaur.)

271.—Bones of hand of Gorilla. (After Huxley.)

272.—Bones of human hand, seen from the back. (After H. Meyer.)

is so prominent in the higher Vertebrata as the upper leg) (Fig. 270, *r* and *u*) and the lower arm (or

many-fingered fish-fins thus gave rise, by a process of gradual reversion and differentiation, to the five-fingered mammalian foot, which occurs first in the Sozobranchia, and from them, has been transmitted on the one hand to the Reptiles, and to Mammals, up to Man, on the other (Fig. 272). Simultaneously with the reduction of the number of the fin-rays, a further differentiation affected the fin-stem or pterygium, which became transversely divided into the upper and lower arms (or legs), and a modification took place in the pectoral girdle, which in the higher Mammals originally consists, anteriorly and posteriorly, of three bones. The simple pectoral girdle of the original shoulder-girdle separates, on each side into an upper (dorsal) piece—the shoulder-blade (*scapula*)

and a lower (ventral) piece; the anterior portion of the latter constitutes the pro-key (or collar) bone (*procoracium*) and its posterior part the raven-bone (*coracoid*). The simple arch of the pelvic girdle breaks up, correspondingly, into an upper (dorsal) piece—the intestinal bone (*os ilium*), and a lower (ventral) piece; the anterior portion of the latter becomes the pubic bone (*os pubis*), and the posterior portion the hip-bone (*os ischii*). Table XX p. 278, shows the correspondence of these three pieces of the pelvic girdle with those of the shoulder-girdle. The latter, however, in the key-bone or collar-bone (*clavicula*) possesses a fourth, wanting in the former. (Cf. Gegenbaur.)

As in the girdle, so in the trunk of the limbs there is originally an absolute agreement between the anterior and posterior limbs. The first section of the trunk is supported by a single strong bone—in the anterior limbs, the humerus (*humerus*); in the posterior, the upper leg (femur, *femur*). The second section, on the other hand, contains two bones: on the anterior extremity the spoke-bone (*radius*, Fig. 270, *r*), and the ell-bone (*ulna*, Fig. 270, *u*); in the posterior the two corresponding bones, the shin-bone (*tibia*) and the calf-bone (*fibula*). (Cf. skeletons in Fig. 196 and 204–208). Moreover, the subsequent small and numerous bones of the wrist (*carpus*) and of the ankle (*tarsus*) correspond; so do the five bones of the middle of the hand (*metacarpus*) and of the middle of the foot (*metatarsus*). Finally, the same is true of the five digits attached to these parts, which in their characteristic structure of a series of small bone-pieces correspond in the anterior and posterior limbs. Charles Martins, of Montpellier, an excellent morphologist, has shown that, in detail, the anterior and posterior limbs correspond.¹⁸³

Comparative Anatomy thus shows that the skeleton of the limbs in Man is composed of the same bones, and in the same manner as the skeleton in the four higher vertebrates, we may justly infer their common descent from a single parent-form. This parent-form was the most ancient Amphibian possessing five digits both on the fore and the hind limbs. The outermost part of the limbs has been very much modified by adaptation to the conditions of life. The diversities in this point in the mammalian class are enormous. The slender legs of the swift Deer and the strong, springy legs of the Hare, the climbing feet of the Sloth and the digging feet of the Mole, the fins of the Whale and the wings of the Bird are all instances. It will, of course, be admitted by all that these organs of locomotion are as diverse as possible in form, of size, form, and special function. And yet the bony skeleton is substantially the same in them all. In these different forms of limbs the same characteristic is always represented in essentially the same strongly marked combination; and here we have a weighty confirmation of the theory of descent, such as is hardly afforded by the Comparative Anatomy of any other organ. (Cf. Plate I., vol. ii. of "History of Creation.") True, in the different Mammals, the skeleton is subject to arrests of development and reversions, in addition to those due to special adaptation (Fig. 273). Thus, in the forelimb (or hand) of the Dog the first digit, or thumb, is rudimentary (Fig. 273 II.). In the Pig (III.) and the Tapir (V.) it has entirely disappeared. So, too, in the Ruminants, the Ox, Fig. IV.) the second and fifth digits are rudimentary, and only the third and fourth are well developed.

loped. Finally, in the Horse, only one digit, the third, is perfectly developed (Fig. VI, 3). And yet all these diverse fore-feet, as also the hand of the Ape (Fig. 271) and the human hand (Fig. 272), have originated from the same common five-fingered parent-form. This is proved, not only by the rudiments of the aborted digits, but also by the homologous disposition of the wrist-bones (Fig. 273, a-f) (*Vide supra*, p. 124.)

The same story is also told by the germ-history of the limbs, which is originally identical, not only in all Mammals but in all Vertebrates. However different the limbs of the various Skulled Animals (*Craniota*) afterwards appear in their fully developed state, they nevertheless all originate from the same simple rudiment. (Cf. Plates VI and VII)




FIG. 273.—Skeleton of hand or fore-foot of six Mammals. I. Man; Dog; III. Pig; IV. Ox; V. Tapir; VI. Horse. *r*, Radius; *u*, ulna; *a*, scaphoid; *b*, semi-lunar; *c*, triquetrum (cuneiform); *d*, trapezium; *f*, capitatum (unciform process); *g*, hamatum (unciform bone); *p*, pisiform; 1, thumb; 2, digit; 3, middle finger; 4, ring finger; 5, little finger. (After Gegenbaur.)

i. p. 362; *f*, fore-leg, *b*, hind-leg.) In all, the first rudiment of each limb in the embryo is a simple wart, or small protuberance, which grows from the side of the body between the dorsal and ventral sides (Figs. 119 and 120, vol. i. pp. 357, 359; and 137, pp. 381, 382). The cells composing these knobs belong to the skin-fibrous layer. The outer surface is coated with the horn-plate, which is rather thicker at the apex of the protuberance (Plate IV. Fig. 5, *x*). The two anterior protuberances appear at a rather earlier period than the posterior. By differentiation of the cells, these simple rudiments develop immediately, in Fishes and in the Annelida, into fins. In the higher vertebrate classes, on the contrary, each of the four protuberances, in the course of its development, assumes the form of a stalked plate, the outer portion of which being narrower and thicker, the inner broader and thinner. The inner portion, or the handle of the plate, then divides into two sections: the upper and lower legs (or arms). Four notches then appear in the free edge of the plate, and these gradually become deeper; these are the divisions between the five digits (Plate VIII. Fig. 1). The latter soon become more prominent. At first, however, the five digits, both on the fore and on the hind limbs, are joined by a thin connecting web-like membrane; this recalls the original adaptation of the foot as a swimming-fin. The further development of the limbs from this most simple rudiment takes place in the same way in all Vertebrates; that is, by the modification of certain groups of the cells of the skin-fibrous layer into cartilage, of other groups into muscles, yet others into blood-vessels, nerves, etc. Probably the differentiation of all these various tissues occurs actually in the limbs. Like the vertebral column and the skull, the

bony parts of the limbs are also formed at first from undifferentiated cell-groups of the skin-fibrous layer. They afterwards change into cartilage, and from these the permanent bones originate by a tertiary process.¹⁸⁴

The development of the muscles, or the active organs of locomotion, is, as yet, of much less interest than that of the skeleton, or the passive instruments of motion. The comparative Anatomy of these is, indeed, of much higher importance than their Embryology. But as very little attention has, as yet, been paid to the Comparative Anatomy and Ontogeny of the muscular system, we have only general ideas of its Phylogeny also. The muscular system as a whole has developed in the most intimate reciprocal correlation with the bone system.¹⁸⁵

TABLE XXXV.

STEMATIC SURVEY OF THE MOST IMPORTANT PERIODS IN THE PHYLOGENY OF THE HUMAN SKELETON.

I. *First Period : Skeleton of the Chordonia* (Fig. 187, p. 90).

The entire skeleton is formed by the notochord.

II. *Second Period : Skeleton of the Acrania* (Fig. 189, p. 91).

A notochord-membrane, the dorsal continuation of which forms a cover-
 round the medullary tube, is formed round the notochord.

III. *Third Period : Skeleton of the Cyclostomi* (Fig. 190, p. 103).

A cartilaginous primordial skull develops round the anterior extremity
 the notochord, from the notochord-membrane. An outer cartilaginous
 skeleton forms round the gills.

IV. *Fourth Period : Skeleton of the older Selachii* (Fig. 268, p. 302).

A primitive vertebral column, with upper and lower arches (the gill-
 arches and ribs) forms round the notochord. The remnant of the outer gill-
 skeleton remains with the inner. Two pairs of limbs, with pinnate (biserial)
 skeletons, appear.

V. *Fifth Period : Skeleton of the more recent Selachii* (Fig. 269, p. 302).

The anterior gill-arches change into lip-cartilage and jaw-arches. The
 external gill-skeleton is lost. The skeleton of the two pairs of fins becomes
 uniserial (semi-pinnate).

VI. *Sixth Period : Skeleton of the Dipneusta* (Fig. 2, Plate XII.).

The skull becomes partially ossified ; as does the shoulder-girdle.

VII. *Seventh Period : Skeleton of the Amphibia* (Fig. 270, p. 302).

The gill-arches are modified into parts of the tongue-bone, and of the jaw-
 apparatus. On the semi-pinnate skeletons of the fins the rays diminish in
 number to four, thus giving rise to the five-toed foot. The vertebral
 column ossifies.

VIII. *Eighth Period: Skeleton of the Monotremata* (Fig. 196, p. 148)

The vertebral column, skull, jaws, and limbs, acquire the characteristics of Mammals.

IX. *Ninth Period: Skeleton of the Marsupialia* (Fig. 197, p. 152).

The coracoid bone of the shoulder-girdle becomes atrophied, as remnant of it amalgamates with the shoulder-blade.

X. *Tenth Period: Skeleton of the Semi-apes* (Fig. 199, p. 164).

The pouch-bones, which distinguish Monotremes and Mars disappear.

XI. *Eleventh Period: Skeleton of the Anthropoid Apes*
(Figs. 204-208, p. 179).

The skeleton acquires the peculiar development shared by exclusively with the Anthropoid Apes.

CHAPTER XXIII.

DEVELOPMENT OF THE INTESTINAL SYSTEM.

The Primitive Intestine of the Gastrula.—Its Homology, or Morphological Identity in all Animals (excepting the Protozoa).—Survey of the Structure of the Developed Intestinal Canal in Man.—The Mouth-cavity.—The Throat (*pharynx*).—The Gullet (*oesophagus*).—The Wind-pipe (*trachea*) and Lungs.—The Larynx.—The Stomach.—The Small Intestine.—The Liver and Gall-bladder.—The Ventral Salivary Gland (*pancreas*).—The Large Intestine.—The Rectum.—The First Rudiment of the Simple Intestinal Tube.—The Gastrula of the Amphioxus and of Mammals.—Separation of the Germ from the Intestinal Germ Vesicle (*Gastrocystis*).—The Primitive Intestine (*Protogaster*) and the After Intestine (*Metagaster*).—Secondary Formation of the Mouth and Anus from the Outer Skin.—Development of the Intestinal Epithelium from the Intestinal-glandular Layer, and of all other parts of the Intestine from the Intestinal-fibrous Layer.—Simple Intestinal Pouch of the Lower Worms.—Differentiation of the Primitive Intestinal Tube into a Respiratory and a Digestive Intestine.—Gill-intestine and Stomach-Intestine of the Amphioxus and Ascidian.—Origin and Significance of the Gill-openings.—Their Disappearance.—The Gill-arches and the Jaw-skeleton.—Formation of the Teeth.—Development of the Lungs from the Swim-bladder of Fish.—Differentiation of the Stomach.—Development of the Liver and Pancreas.—Differentiation of the Small and Large Intestines.—Formation of the Cloaca.


“Cautious people require us to confine ourselves to gathering materials, and to leave it to posterity to raise a scientific structure from those materials; because only in that way can we escape the ignominy of having the theories we believed in overthrown by the advance of knowledge. The unreasonableness of this demand is apparent enough from the fact that

Comparative Anatomy, like every other science, is endless; and the the endlessness of the accumulation of materials would never allow them to comply with this demand, to reap any harvest from this field further than this, history teaches us clearly, that no age in which scientific inquiry has been active, has been able so to deny itself, as, setting it of its researches in the future, to refrain from drawing conclusions from its larger or smaller treasury of observations, and from trying to gaps with hypotheses. It would, indeed, be a hopeless proceeding order to avoid losing any part of our possessions, we should re acquire any possessions whatever."—KARL ERNST BAER (1819).

AMONG the vegetative organs of the human body, the development of which we now turn our attention, the intestinal canal is the most important. For the intestine is the oldest of all the organs of the animal body carries us back to the earliest time of organological entiation, to the first period of the Laurentian Epoch we have already seen, the result of the first division of labour in the homogeneous cells of the earliest many animal body must have been the formation of a new intestinal canal. The first duty and the first need of organism is self-support. This task is accomplished by two functions of nutrition and of the covering of the body. When, therefore, in the primæval collection of homogeneous cells (*Synmæbium*), of the phylogenetic existence of which we yet have evidence in the ontogenetic development of the mulberry-germ (*Morula*), the several members of the community began to divide the work of life, they were first obliged to engage in two separate tasks. One half modified into nutritive cells, enclosing a digestive cavity, the intestinal canal; the other half, on the contrary, developed into covering cells, forming the outer integument of this intestinal canal, and, at the same time, the whole body. Thus arose the first two germ-layers

nutritive, or vegetative layer, and the outer, covering, mal layer.

we try to construct for ourselves an animal body of simplest conceivable form, possessing such a primitive inal canal, and the two primary germ-layers forming all, the result is necessarily the very remarkable form of the gastrula, which we have shown to exist onderful uniformity throughout the whole animal



274.—Gastrula of a Chalk-sponge (*Olynthus*): *A*, from outside; longitudinal section through the axis; *g*, primitive intestine; *o*, primitive mouth; *i*, intestinal layer, or entoderm; *e*, skin-layer, or exoderm.

in the Sponges, Sea-nettles (*Acalephæ*), Worms, bodied Animals (*Mollusca*), Articulated Animals (*Arthro-* and Vertebrates (Figs. 174–179, p. 65). In all these s animal tribes the gastrula reappears in the same y simple form (Fig. 274). Its whole body is really y the intestinal canal; the simple cavity of the body, gestive intestinal cavity, is the primitive intestine

(*protogaster*, *g*); its simple opening, the primitive mouth (*protostoma*, *o*), is at once mouth and anus; and the two cell-strata which compose its wall, are the two primary germ-layers: the inner, the nutritive, or vegetative germ-layer, is the intestinal layer (*entoderma*, *i*); and the outer, covering layer, which, by means of its cilia, is also the agent of motion, is the animal layer, or skin-layer (*exoderma*, *e*). This highly important fact, that the gastrula appears as a early larval condition in the individual development of the most varied animals, and that this gastrula always exhibits the same structure, and that the very differently developed intestinal canals of the most varied animals, arises, ontogenetically, from the same extremely simple gastrula-intestine, this very important fact justifies, in accordance with the fundamental law of Biogeny, two conclusions, which involve important results, and of which one is general and one special. The general conclusion is an inductive one, and may be stated thus: The very variously formed intestinal canal of all the different Intestinal Animals has developed, phylogenetically, from one common and extremely simple primitive intestine, from the intestinal cavity of the *Gastrea*, that primæval common parent-form which is at the present reproduced, in accordance with the fundamental law of Biogeny, in the gastrula. The second, the special conclusion, which is connected with the former, is deductive, and may be stated thus: The intestinal canal in Man as a whole is homologous with the intestinal canal in all other animals; it has the same original significance, and has developed from the same rudimentary form.¹⁸⁸

Before proceeding to trace the history of the development of the human intestinal canal in detail, it will be

necessary briefly to get a correct idea of the more general conditions of the formation of the intestinal canal in the developed Man. Not until this is known can the development of the several parts be correctly understood. (Cf. Plates IV. and V., vol. i. p. 321.) The intestinal canal in the developed Man is, in all essential points, exactly similar in form to those of all other higher Mammals, and, especially, to that of the Catarhines, the Narrow-nosed Apes of the Old World. The entrance to the intestinal canal is the mouth-opening (Plate V. Fig. 16, o). Food and drink pass first through this into the mouth-cavity, in the lower part of which is the tongue. The human mouth-cavity is hedged with thirty-two teeth, attached in two rows to the two jaws, the upper and lower. It has already been stated that the series of teeth is formed in Man exactly as in the Catarhine Apes, but differs from the corresponding part in all other animals (p. 173). Above the mouth-cavity is the double nose-cavity; the two parts of this are separated by the partition-wall of the palate. But, as we have seen, the nasal cavity is not originally separated at all from the mouth-cavity, a common nasal and mouth cavity being primarily formed in the embryo, and this separates at a later period into two separate stories by the hard palate-roof: the upper is the nasal cavity, the lower is the mouth-cavity. The nasal cavity is connected with certain air-filled bony cavities; the jaw-cavities in the upper jaw, the frontal cavities in the frontal bone, and the sphenoid cavities in the sphenoid bone. Numerous glands of various kinds open into the mouth-cavity, particularly many small mucous glands and three pairs of large salivary glands.

The human mouth-cavity is half closed at the back by

the vertical curtain which we call the soft palate, the centre of the lower part of which is situated the uvula. A glance with the mouth open into a mirror is sufficient to show the form. The uvula is of importance because it occurs only in Men and in Apes. On both sides of the soft palate are the tonsils (*tonsillæ*). Through a gate-like arched opening situated beneath the soft palate we pass into the throat-cavity (*pharynx*; Plate 16, *sh*), which lies behind the mouth-cavity. This is partly visible in the open mouth when reflected in a mirror. Into the throat-cavity a narrow passage opens on each side (the Eustachian tube of the ear), which leads directly into the tympanic cavity of the ear (Fig. p. 260). The throat-cavity is continued into a narrow tube, the gullet (*œsophagus*, *sr*). Through this the masticated and swallowed food passes down into the stomach. The wind-pipe (*trachea*, *lr*) also opens into the upper part of the throat, and leads thence to the lungs. The opening of this is protected by the epiglottis, through which the food passes. The respiratory organs, the lungs (Plate IV. Fig. 8, *lu*), are situated, in Man, as in Mammals, in the right and left sides of the breast (*thorax*), and midway between them is the heart (Fig. 8, *hr*, *hl*). At the upper end of the wind-pipe (trachea) below the epiglottis just spoken of, is a peculiarly differentiated section, the larynx, which is protected by a cartilaginous frame. The larynx is the most important part of the human voice and speech, and also develops into the part of the intestinal canal. In front of the larynx is the thyroid gland (*thyreoidea*), which occasionally enlarges into the so-called "goitre."

The gullet (*œsophagus*) passes downward through the thorax, along the vertebral column, behind the lungs and the heart, and enters the ventral cavity, after penetrating the diaphragm. The latter (Fig. 16, *z*) is a membranous, muscular, transverse partition, which in all Mammals (and only in these) completely separates the chest-cavity (*thorax*, *c*) from the ventral cavity (*c*_v). As has been said, this division does not originally exist; at first a common chest and ventral cavity, the *cœloma*, or the pleuro-peritoneal cavity, is formed in the embryo. It is only afterwards that the diaphragm forms a muscular, horizontal partition between the chest and the ventral cavities. This partition then completely separates the two cavities, and is penetrated only by separate organs passing through the



FIG. 275.—Human stomach and gall-intestine in longitudinal section: *a*, cardia (limit of the *œsophagus*); *b*, fundus (blind sac of the left side); *c*, pylorus fold; *d*, pylorus valve; *e*, pylorus-cavity; *fgh*, gall-intestine; *i*, mouth of the gall-duct and of the pancreas duct. (After H. Meyer.)

chest-cavity into the ventral cavity. One of the most important of these organs is the gullet (*œsophagus*). After this has passed through the diaphragm into the ventral cavity it enlarges into the stomach in which digestion

especially takes place. The stomach of an adult (Fig. 275, Plate V. Fig. 16, *mg*) is an oblong sac, placed somewhat obliquely, the left side of which widens into a blind-sac, the base of the stomach or fundus (*b*), while the right side narrows, and passes at the right end, called the pylorus (*e*), into the small intestine. Between these two parts of the intestine is a valve, the pyloric valve (*d*), which only opens when the food-pulp (chyme) passes from the stomach into the small intestine. The stomach itself is the most important digestive organ, and serves especially to dissolve the food. The muscular wall of the stomach is comparatively thick, and, on the outside, has strong muscular layers, which effect the digestive movements of the stomach;—on the inside, it has a great number of small glands, the gastric glands, which secrete the gastric juice.

Next to the stomach follows the longest part of the whole intestinal canal, the central, or small intestine (*chylogaster*). Its principal function is to effect the absorption of the fluid mass of digested food, or the food-pulp (chyme), and it is again divided into several sections, the first of which, the one immediately following the stomach, is called the gall-intestine, or "twelve-finger intestine" (*duodenum*, Fig. 275, *fgh*). The gall-intestine forms a short loop curved like a horse-shoe. The largest glands of the intestinal canal open into it: the liver, the most important digestive gland, which furnishes the bile, or gall, and a very large salivary gland, the ventral salivary gland, or pancreas, which secretes the digestive saliva. Both of these glands pour the juices they secrete, the bile and pancreatic juice, into the duodenum (*i*) near each other. In adults the liver is a very large gland, well supplied with blood, lying on the

side immediately below the diaphragm, and separated from the latter from the lungs (Plate V. Fig. 16, *lb*). The stomach lies somewhat further back and more to the right (Fig. 16, *p*). The small intestine is so long that it has to lie in many folds in order to find room in the limited space of the ventral cavity; these coils are the bowels. They are divided into an upper intestine, called the empty intestine (*jejunum*), and a lower, the crooked intestine (*ileum*). In this latter part lies that part of the small intestine at which, in the embryo, the yolk-sac opens into the intestinal tube. This long, thin intestine then passes into the large intestine, from which it is separated by a peculiar valve. Directly behind this "Bauhinian valve" the first part of the large intestines forms a broad pouch-like expansion, the blind intestine (*cæcum*), the atrophied extremity of which is a well-known rudimentary organ, the vermiform process (*processus vermiformis*). The large intestine (*colon*) consists of three parts; an ascending part on the right, a transverse central part, and a descending part on the left. The latter finally curves like an S, called the "sigmoid flexure," into the last part of the intestinal canal, above the rectum, which opens at the back by the anus (Plate V. Fig. 16, *a*). Both the large intestine and the small intestine are furnished with numerous glands, most of them very small, and which secrete mucous and other juices.

Along the greater part of its length the intestinal canal is attached to the inner dorsal surface of the ventral cavity, or to the lower surface of the vertebral column. It is fastened by means of the thin, membranous plate, called the mesentery, which develops directly under the notochord

from the intestinal-fibrous layer, at the point where it curves into the outer lamina of the side-layer, into the skin-fibrous layer (Plate IV. Fig. 5, *g*). The curving was distinguished as the middle-plate (Fig. 99, *mp*). The mesentery is, at first, very short (Plate V. Fig. 14, *g*); soon lengthens considerably at the central part of the intestinal canal, and takes the form of a thin, transverse membranous plate, which has to be extended further the farther the folds of the intestine diverge from the vertebral column where they are first attached to the vertebral column. Blood-vessels, lymphatic vessels, and nerves which the intestinal canal traverse this mesentery.

Although, therefore, the intestinal canal, in the human being forms an extremely complex organ though it shows in its details so many intricate and diversified structural arrangements,—into which we cannot here,—this entire structure has developed, historically from that simplest form of primitive intestine which was possessed by our gastrea ancestors, and which the extant gastrula now exhibits. We have already shown (Chapter VIII.) that the peculiar Hood-gastrula (*A. hoodi*) of Mammals (Fig. 277) may be referred to the original Bell-gastrula (*Archigastrula*) form, among Vertebrates, is now accurately retained solely in the *Amphioxus* (Fig. 276; Plate X. Fig. 10).

Like the latter, the gastrula of Man and of all Mammals must be regarded as the ontogenetic reproduction of that phylogenetic evolution-form which we call the Gastraea, and in which the whole body of the animal is intestine.

The peculiar form and mode in which the co

an intestinal canal develops from the simple gastrula which is similar to that in other Mammals, can therefore only correctly understood when it is considered in light of Phylogeny. We must, accordingly, distinguish




FIG. 276.—Archigastrula of *Amphioxus* (in longitudinal section): *d*, primitive intestine; *o*, primitive mouth; *i*, intestinal layer; *e*, skin-layer.

FIG. 277.—Amphigastrula of Mammal (in longitudinal section). The primitive intestine (*d*) and primitive mouth (*o*) are filled up by the cells of the intestinal layer (*i*); *e*, skin-layer.

between the original primary intestine ("the primitive intestine, or *protogaster*") of the Skull-less Animals (*Acrania*), and the differentiated or secondary intestine ("after intestine, or *metagaster*") of the Skulled Animals (*Craniota*). The intestine of the *Amphioxus* (the representative of the *Acrania*) forms no yolk-sac, and develops, palingenetically, from the entire primitive intestine of the gastrula. The intestine of the Skulled Animals, on the other hand, has a modified, kenogenetic form of evolution, and differentiates at a very early period into two different parts: into the permanent secondary intestine, which alone

gives rise to the various parts of the differentiated intestinal system, and the transient yolk-sac, which serves only as a storehouse of materials for the building of the embryo. The yolk-sac attains its greatest development in Primitive Fishes (*Selachii*), Bony Fishes (*Teleostei*), Reptiles, and Birds. In Mammals, and especially in Placental Animals, it is atrophied. The peculiar intestinal development of the Cyclostomi, Ganoids, and Amphibia must be regarded as an intermediate form, between the palingenetic intestinal development of the Skull-less animals, and the kenogenetic intestinal development of the Amnion Animals (*Amniota*).¹⁸⁷

We have already seen in what a peculiar way the development of the intestine takes place ontogenetically in the human embryo and in that of other Mammals. Immediately from the gastrula of these originates a globular intestinal germ-vesicle (*gastrocystis*), filled with fluid (Figs. 72, 73, vol. i. p. 289). In the wall of this is formed the lyre-shaped germ-shield, on the lower side of which, along the middle line, appears a shallow groove, the first rudiment of the future, secondary intestinal tube.

This intestinal groove grows constantly deeper, and its edges curve toward each other, to grow together at last and form a tube (Fig. 100, vol. i. p. 333). The wall of this secondary intestinal tube consists of two membranes of the inner, intestinal-glandular layer, and of the outer, intestinal-fibrous layer. The tube is completely closed at the ends, having only an opening in the centre of the lower wall, by which it is connected with the intestinal germ-vesicle (Plate V. Fig. 14). The latter, in the course of development, becomes continually smaller, as the intestinal canal continues

to grow larger and more perfect. While, at first, the intestinal tube appears only as a little appendage on one side of the great intestinal germ-vesicle (Fig. 278), the remnant of the latter afterwards forms only a very inconsiderable appendage of the great intestinal canal. This appendage is the yolk-sac, or navel-vesicle. It entirely loses its importance, and at length disappears, while the intestinal canal is finally closed at the original central opening, where it forms the so-called intestinal navel (Fig. 94, vol. i. p. 312).

It has also been said that this simple cylindrical intestinal tube, in Man as in all Vertebrates, is at first entirely closed at both ends (Plate V. Fig. 14), and that the two permanent openings of the intestinal canal—at the anterior extremity, the mouth, at the posterior, the anus—form only secondarily, and from the outer skin. At the fore end, a shallow mouth-furrow originates in the outer skin, and this grows toward the blind, anterior end of the head intestinal cavity, into which it finally breaks. In the same way a shallow furrow for the anus is formed behind in the skin, and this soon grows deeper, and grows toward the blind posterior end of the pelvic intestinal cavity, with which it finally unites. At both extremities there is, at first, a thin partition between the outer skin-furrow and the blind end of the intestine, and this disappears when the opening is made.¹⁸⁸

Directly in front of the anus the allantois grows out of the posterior intestine; this is the important embryonic appendage which develops, in Placental Animals, and only in these (thus in Man too) into the placenta (Figs. 278, 279, *l*; Plate V. Fig. 14, *al*). In this more developed form—represented in the diagram (Fig. 94, *4*, vol. i. p. 312)—the intestinal

canal of Man, like that of all other Mammals, now forms a slightly-curved, cylindrical tube, which has an opening at both ends, and from the lower wall of which depend two sacs; the anterior navel-bladder, or yelk-sac, and the posterior allantois, or primitive urinary sac.

Microscopic observation shows that the thin wall of the simple intestinal tube and of its two bladder-like appendages is composed of two distinct cell-strata. The inner stratum which coats the entire cavity, consists of larger, darker cells



FIG. 278.—Human embryo of the third week, with the amnion and allantois. The great globular yelk-sac is below, the bladder-like allantois on the right; there are as yet no limbs. The germ, with its appendages, is enclosed in the tufted membrane (*chorion*).

FIG. 279.—Human embryo, with amnion and allantois, in the fourth week. (After Krause.) The amnion (*w*) lies pretty close to the body. The greater part of the yelk-sac (*d*) has been torn away. Behind this the allantois appears as a small pear-shaped bladder. Arms (*f*) and legs (*l*) are already commenced: *v*, fore-brain; *z*, twixt-brain; *m*, mid-brain; *n*, after-brain; *a*, eye; *k*, three gill-arches; *c*, heart; *s*, tail.

and is the intestinal-glandular layer. The outer stratum consists of lighter, smaller cells, and is the intestinal fibrous-layer. The cavities of the mouth and the anus are the only exceptions to this, because they originate from the outer skin. The inner cell-coating of the entire mouth-cavity is therefore furnished, not by the intestinal glandular-layer, but by the skin-sensory layer, and its muscular lower layer, not by the intestinal-fibrous layer, but by the skin-fibrous layer. This is equally true of the wall of the anal cavity (plate V. Fig. 15).

If the question be asked, what relation these component germ-layers of the primitive intestinal wall bear to the infinitely varied tissues and organs which we afterwards find in the developed intestine, the answer is extremely simple. The relations of these two layers to the formation and differentiation of the tissues of the intestinal canal with all its parts, may be condensed into a single sentence: The intestinal epithelium, that is, the inner, soft cell-stratum which coats the cavities of the intestinal canal and of all its appendages, and which directly accomplishes the nutritive process, develops solely from the intestinal-glandular layer; on the contrary, all other tissues and organs belonging to the intestinal canal and its appendages, proceed from the intestinal-fibrous layer. From this latter, therefore, originates the entire outer covering of the intestinal tube and its appendages; the fibrous connective tissue and the smooth muscles which compose its fleshy skin; the cartilages which support these, for example, the cartilage of the larynx and of the trachea; the numerous blood and lymph vessels which absorb nutrition from the wall of the intestine; in short, everything belonging to the intestine, with the

exception of the intestinal epithelium. From the intestinal fibrous layer originates also the entire mesentery and the adjacent parts, the heart, the large blood-vessels, body, etc. (Plate V. Fig. 16).

Let us now turn aside for a moment from this rudimentary intestine of Mammals, in order to insure a comparison between it and the intestinal canal of lower Vertebrates and Worms, which we have learned to recognize as the ancestors of Man. In the simplest form, the planarian worm, or Turbellaria (*Rhabdocœlum*, Fig. 280), we find a very simple intestinal form. As in the gastrula, the intestine in these Worms is a simple pouch with a single opening, which latter acts both as mouth and anus (*m*). In the intestinal pouch has, however, differentiated into two portions, an anterior throat-intestine (*sd*) and a posterior stomach-intestine (*d*). This differentiation becomes important in the Ascidia (Fig. 281) and in the Amphioxus (Fig. 282), which connects the Worms with the Vertebrates. In these two animal forms the intestine is essentially identical; the anterior portion forms the respiratory intestine, the posterior forms the digestive stomach-intestine. In both it develops, palæogenetically, directly from the primitive intestine of the gastrula (Plate XI. Figs. 1 and 2). But the original mouth-opening of the gastrula, the primitive mouth, afterwards closes, and in its place is formed the later anus. In the same way, the mouth-opening of the Amphioxus and of the Ascidian is transformed into a respiratory opening, as is the mouth-opening of Man, and generally of all Skulled Animals (*Craniota*). The secondary transformation of the mouth of the Lancelet is connected, as I have conjectured with some probability, with the formation

the gill-openings, which appear directly behind it on the intestine. The front portion of the intestine has thus





FIG. 280.—A simple Gliding Worm (*Rhabdocalum*): *m*, mouth; *sd*, throat-epithelium; *sm*, throat muscle-mass; *d*, stomach-intestine; *nc*, renal ducts; *f*, ciliated outer-skin; *nm*, openings of the latter; *au*, eye; *na*, nose-pit.

FIG. 281.—Structure of an Ascidian (seen from the left side, as in Plate XI. Fig. 14). The dorsal side is turned toward the right, the ventral side to the left; the mouth-opening (*o*) is above; at the opposite, tail end, the ascidian has become adherent. The gill-intestine (*br*), perforated by many openings, extends into the stomach-intestine. The terminal intestine opens through the anus (*a*) into the gill-cavity (*cl*), from which the excrement is passed out with the respired water through the gill-pore, or cloaca opening (*a'*); *m*, mantle. (After Gegenbaur.)



the tribal history of Vertebrates.

The most important process with in the further ontogenetic development of the intestinal canal in the human is the origin of the gill-opening at the head of the human embryo, the throat very early unites with the rest of the body, and four openings on the right and left sides of the body, and these lead from without into the throat-cavity. These openings are the gill-openings, and the partitions separating them are the gill-arches (Figs. 116-118, vol. i. p. 356; and V., Fig. 15, *ks*). These embryonic developments are very interesting; for

FIG. 282.—Lancelet (*Amphioxus lanceolatus*) the natural size, seen from the left side. The longitudinal axis is perpendicular, the mouth at the head end, the tail end below (as in Plate XI. Fig. 1).

produce, in accordance with the fundamental principle of Biogeny, the same process which was originally of the greatest importance to the development of the whole vertebrate tribe. This process was the differentiation of the intestinal canal into two sections: an anterior, respiratory part, the gill-intestine, which serves only for breathing, and a posterior, digestive part, the stomach-intestine, which serves only for digestion. As we meet with this very characteristic differentiation of the intestinal tube into two, physiologically, very distinct main sections, not only in the Amphioxus, but also in the Ascidian and the Appendicularia, we can safely conclude that it also existed in our common ancestors, the Chorda Animals (*Chordonia*), especially as even the Acorn Worm (*Balanoglossus*) has it (Fig. 186, p. 86). All other Invertebrate Animals are entirely without this peculiar arrangement.

The number of the gill-openings is still very large in the Amphioxus, as in Ascidians and in the Acorn Worm. In the Skulled Animals it is, on the contrary, very much lessened. Fishes mostly have from four to six pairs of gill-openings. In the embryos of Man and the higher Vertebrates also, only three or four pairs are developed, and these appear at a very early period. The gill-openings are permanent in Fishes, and afford a passage to the water which has been breathed in through the mouth (Figs. 191, 192, p. 113; Plate V. Fig. 13, *ks*). On the other hand, the Amphibians lose them partially, and all the higher Vertebrates entirely. In the latter, only a single vestige of the gill-openings remains, the remnant of the first gill-opening. This changes into a part of the organ of hearing; from it originates the outer ear-canal, the tympanic cavity, and the Eustachian tube.

TABLE XXXVI.

Systematic Survey of the Development of the Human Intestinal
N.B.—The parts marked thus † are processes from the intestina

I. First main section of the Intestinal System: the Respiratory Intestine (Gill Intestine). PNEOGASTER. (<i>Tractus respiratorius.</i>)	1. Mouth-cavity (<i>Orum oris</i>)	{ Mouth-opening Lips Jaws Teeth Tongue Tongue bone † Salivary glands Soft palate Uvula	<i>Rima oris</i> <i>Labia</i> <i>Maxilla</i> <i>Dentes</i> <i>Lingua</i> <i>Os hyoides</i> <i>Glandulae saliv.</i> <i>Velum palatinum</i> <i>Uvula</i>
	2. Nose-cavity (<i>Orum nasi</i>)	{ Nose canal † Jaw cavities † Frontal cavities † Ethmoid cavity	<i>Meatus narium</i> <i>Sinus maxillaris</i> <i>Sinus frontalis</i> <i>Sinus ethmoidalis</i>
	3. Throat-cavity (<i>Orum pharyngis</i>)	{ Isthmus of the throat Tonsils Pharynx † Eustachian tube † Tympanic cavity † Brain-appendage † Thyroid gland	<i>Isthmus faucium</i> <i>Tonsillae</i> <i>Pharynx</i> <i>Tuba Eustachiana</i> <i>Cavum tympani</i> <i>Hypophysis</i> <i>Thyreoidea</i>
	4. Lung-cavity (<i>Orum pulmonis</i>)	{ † Larynx † Windpipe † Lungs	<i>Larynx</i> <i>Trachea</i> <i>Pulmones</i>
II. Second main section of the Intestinal System: Digestive Intestine (Stomach Intestine). PEPTOGASTER. (<i>Tractus digestivus.</i>)	5. Anterior Intestine (<i>Prosogaster</i>)	{ Gullet Stomach-opening Stomach Stomach exit	<i>Oesophagus</i> <i>Cardia</i> <i>Stomachus</i> <i>Pylorus</i>
	6. Central Intestine (<i>Mesogaster</i>)	{ Gall-intestine † Liver † Pancreas Empty intestine († Yolk-sac, or navel-bladder) Crooked intestine	<i>Duodenum</i> <i>Hepar</i> <i>Pancreas</i> <i>Jejunum</i> (<i>Vesicula umbilicalis</i>) <i>Ileum</i>
	7. Posterior Intestine (<i>Epigaster</i>)	{ Large intestine † Blind intestine † Vermiform process of the caecum Rectum Anal opening	<i>Colon</i> <i>Caecum</i> <i>Processus vermiformis</i> <i>Rectum</i> <i>Anus</i>
	8. Urinary Intestine (<i>Urogaster</i>)	{ († Primitive urinary sac † Urinary tube † Urinary bladder	<i>Allantois</i> <i>Urethra</i> <i>Urocystis</i>

have already considered this remarkable formation, and only call attention once more to the interesting fact that the human middle and external ear is the last remnant of the gill-opening of a Fish. The gill-arches, also, which form the gill-openings, develop into very various parts. In Fishes they remain permanently as gill-arches, carrying the respiratory gill-tufts; so also in the lowest Amphibia; in the higher Amphibia they undergo various modifications in the course of development, and in all the three higher vertebrate classes, thus also in Man, the tongue-bone (*hyoides*) and the bonelets of the ear originate from the gill-arches. (Cf. Plates VI. and VII.)

From the first gill-arch, from the centre of the inner side of which the muscular tongue grows, proceeds the rudimentary jaw-skeleton; the upper and lower jaws which enclose the cavity of the mouth and carry the teeth. The Acrania and Monorhina are entirely destitute of these important parts. They first appear in the genuine Fishes, and have been transmitted by these to the higher vertebrates. The original formation of the human mouth-skeleton, of the upper and lower jaws, can thus be traced back to the earliest Fishes, from which we have inherited it. The teeth originate from the outer skin-covering which covers the jaws; for, as the formation of the whole mouth-cavity takes place from the outer germ-layer, the teeth, of course, also have developed originally from the skin-covering. This can be actually proved by close microscopic examination of the most delicate structural features of the teeth. The scales of Fishes, especially of Sharks, are, in every respect, exactly similar to their teeth (Fig. 283). Thus the human teeth, in their earliest origin, are modified fish-

scales.¹⁸⁸ On similar grounds we must regard the skin glands, which open into the mouth-cavity, as really a skin (epidermic) glands, which have not developed, like other intestinal glands, from the intestinal-glandular part of the intestinal canal, but from the outer skin, from the horn-plate of the outer germ-layer. It is evident that when the mouth develops in this way, the salivary glands must be placed genetically in the same series with the sebaceous, and milk glands of the epidermis.



FIG. 288. — Scales of a Shark (*Centrophorus calceus*). On each rhomboid bone-tablet, lying in the leather-skin, rises a small, three-cornered tooth. (After Gegenbaur.)

form of gill-intestine, it surrenders entirely its function as a respiratory

The human intestinal canal is therefore quite as simple in its original formation as the primitive intestine of the gastrula. It resembles that of the lowest Vertebrate. It then differentiates into two divisions, an anterior gill-intestine and a posterior stomach-intestine, the intestinal canal of the Lancelet and the Ascidian. By the development of the jaws and gills, it is modified into a true intestine. Afterwards, however, the gill-intestine, which is a remnant of the Fish-ancestors, as such is entirely lost. The parts that take a wholly different form notwithstanding that the anterior section of our intestinal canal surrenders entirely its original function as a respiratory intestine; for the extreme

ing and remarkable discovery is now made that even permanent respiratory organ of the higher Vertebrates, air-breathing lungs, has also developed from this anterior end of the intestinal canal. Our lungs, together with wind-pipe (*trachea*) and the larynx, develop from the lateral wall of the anterior intestine. This entire breathing-apparatus, which occupies the greater part of the chest (*thorax*) in the developed Man, is at first merely a very small and simple vesicle or sac, which grows out from the intestinal canal immediately behind the gills, and separates into two lateral halves (Figs., 284, *c*, 285, *c*; Pl. V. Figs. 13, 15, 16, *lu*). This vesicle occurs in all Vertebrates except in the two lowest classes, the Acrania and Protozoa. In the lower Vertebrates, however, it develops, not into lungs, but into an air-filled bladder of considerable size, occupying a great part of the body-cavity (*coeloma*), which is of quite a different significance from the lungs. It serves, not for breathing, but as an hydrostatic apparatus: for vertical swimming movements it is the swimming-bladder of Fish; but the lungs of Man and of other air-breathing Vertebrates develop from the same bladder-like appendage of the anterior intestine, which, in Fishes, becomes the swimming-bladder.

Originally this sac also has no respiratory function, but serves only as an hydrostatic apparatus, augmenting or diminishing the specific gravity of the body. Fishes, in which the swimming-bladder is fully developed, are able to compress it, and thus to condense the air contained in it. In some air sometimes also escapes from the intestinal canal through an air-passage which connects the swimming-bladder with the throat (*pharynx*), and is expelled through

the mouth; in this way the circumference of the swimming-bladder is diminished, and the fish becomes heavy and sinks. When the animal is again about to rise the swimming-bladder is distended by remitting the



FIG. 284.—Intestine of an embryonic Dog (which is represented 137, vol. i. p. 382; after Bischoff), from the ventral side: *a*, gill-arches (pairs); *b*, rudimentary throat and larynx; *c*, lungs; *d*, stomach; *f*, lateral walls of the opened yolk-sac, into which the central intestine opens by a wide aperture; *h*, rectum.

FIG. 285.—The same intestine, seen from the right side: *a*, liver; *c*, stomach; *d*, yolk-sac; *e*, rectum.

pressing force. This hydrostatic apparatus begins to be transformed into a respiratory organ in the Mud-puppy (*Dipneusta*), the blood-vessels in the wall of the swimming-bladder no longer merely separating air, but inhaling fresh air, which has come in through the passage. This process is fully developed in all Amphibians. The original swimming-bladder here generally beco-

lung, and its air-passage a wind-pipe. The amphibian lung has been transmitted to the three higher vertebrate classes, and even in the lowest Amphibia the lung on either side is as yet a very simple, transparent, thin-walled sac—as, for instance, in our common Water-Newts, or Tritons, and very like the swimming-bladder of Fishes. The Amphibia have, it is true, two lungs, a right and a left; but in many Fishes also (in the ancient Ganoids) the swimming-bladder is double, the organ being divided into a right and a left half. On the other hand, the lung of the *Ceratodus* is single (p. 119). The earliest rudiment of the lung in the human embryo and in the embryo of all higher Vertebrates is also a simple, single vesicle, which does not separate till afterwards into a pair of halves—the right and the left lung. At a later period, the two vesicles grow considerably, occupy the greater part of the chest cavity, and lie one on each side of the heart; even in Frogs we find that the simple sac, in the course of its development, is transformed into a spongy body of a peculiar, froth-like texture. This lung-tissue develops as a tree-like, branched gland, bearing berry-like appendages. The process by which the lung-sac was attached to the anterior intestine, which was originally very short, lengthens, by simple growth, into a long thin tube; this tube is the wind-pipe (*trachea*); it opens above into the throat (*pharynx*), and below divides into two branches which pass into the two lungs. In the wall of the wind-pipe ring-shaped cartilages develop, which keep the whole distended; at the upper end of this wind-pipe, below its entrance into the throat, the larynx, the organ of voice and speech, develops. The larynx occurs even in Amphibia in very various stages of development, and with the aid of

Comparative Anatomy we can trace the progressive development of this important organ from its very simple rudiment in the lower Amphibia up to the complex and vocal apparatus represented by the larynx of Birds and Mammals.

Though these organs of voice, speech, and air-respiration develop so differently in the various higher Mammals, they yet all arise from the same simple original rudiment—from a vesicle which grows out of the wall of the anterior intestine. We have thus satisfied ourselves of the interesting fact that both the respiratory apparatus of Vertebrates develop from the fore part of the intestinal canal; first, the primary and more primitive water-respiring apparatus, the gill-body, which is altogether lost in the three higher vertebrate classes; and, afterwards, the secondary and more recent air-breathing apparatus, which acts in Fishes only as a swimming-bladder, but as a lung from the Dipneusta upwards.

We must say a few words about an interesting rudimentary organ of the respiratory intestine, the thyroid gland (*thyreoidea*), the large gland situated in front of the larynx, and below the so-called "Adam's apple," and which, especially in the male sex, is often very prominent; it is produced in the embryo by the separation of the lower wall of the throat (*pharynx*). This thyroid gland is of no use whatever to man; it is only aesthetically interesting, because in certain mountainous districts it has a tendency to enlarge, and in that case it forms the "goitre" which hangs from the neck in front. Its dysteleological interest is, however, far higher; for as Wilhelm Müller of Jena has shown, this useless and unsightly organ is the last remnant of the "hypobranchial groove," which we have

eady considered, and which, in the *Ascidia* and in the *ophioxus*, traverses the middle of the gill-body, and is of great importance in conducting the food into the stomach (Pl. I. p. 420; Plate XI. Figs. 14-16, *y*).¹⁸⁹

The second main section of the intestinal canal, the stomach or digestive intestine, undergoes modifications no less important than those affecting the first main section. In tracing the further development of this digestive section of the intestinal tube, we again find a very complex and composite organ eventually produced from a very simple filament. For the sake of rendering the matter more intelligible, we may distinguish the digestive intestine into three parts: the fore intestine (with the gullet and stomach); the middle intestine, the gall-intestine (with the liver and pancreas); the empty intestine (*jejunum*), and the hooked intestine (*ileus*); and the hind intestine (large intestine and rectum). Here we again find protuberances or appendages of the originally simple intestinal tube which change into very various structures. We have already discussed two of these appendages—the yolk-sac, which protrudes from the middle of the intestinal tube (Fig. 286, *c*), and the allantois, which grows out of the posterior portion of the pelvic intestine as a large sac-like protuberance (*u*). The protuberances from the middle of the intestine are the two great glands which open into the duodenum, the liver (*h*) and the ventral salivary gland.

Immediately behind the bladder-like rudiment of the lungs (Fig. 286, *l*) comes that portion of the intestinal tube which forms the most important part of the digestive apparatus, viz., the stomach (Figs. 284, *d*, 285, *b*). This sac-

shaped organ, in which the food is especially dissolved and digested, is not so complex in structure in the lower Vertebrates as in the higher. Thus, for instance, in many Fishes it appears as a very simple spindle-shaped expansion at the



FIG. 286.—Longitudinal section through an embryonic Chick on the fifth day of incubation: *d*, intestine; *o*, mouth; *a*, anus; *l*, lungs; *g*, mesentery; *v*, auricle of heart; *k*, ventricle of heart; *b*, arterial arches; *l*, aorta; *c*, yolk-sac; *m*, yolk-duct; *u*, allantois; *r*, stalk of allantois; *w*, amnion; *s*, serous membrane. (After Baer.)

beginning of the digestive section of the intestine, while the latter passes from front to rear in a straight line under the spinal column in the central plane of the body. In Mammals the rudiment of this organ is as simple as it then is permanently in Fishes; but at a very early period the various parts of the stomach-sac begin to develop unequally. As the left side of the spindle-shaped pouch grows much more vigorously than the right, and as, at the same time

occurs a considerable obliquity of its axis, it soon assumes an oblique position. The upper end lies more to the right and the lower end more to the left. The anterior end is so placed as to form the long straight canal of the gullet (oesophagus); below the latter, the blind-sac of the stomach (pylorus) bulges out to the left, thus the later form of the stomach is gradually developed (Fig. 275, p. 317). The stomach which was originally verti-

287.—Human embryo of five months from the ventral side; opened up. The breast wall, abdominal wall and liver, have been removed. 3, nasal process; 4, upper jaw; 5, lower jaw; *z*, tongue; *v*, right, *v'*, left ventricle of heart; *o'*, left auricle of heart; *b*, origin of aorta; *b' b'' b'''*, 1st, 2nd, 3rd aorta-arches; *c c' c''*, hollow heart; *y*, lungs (y, lung-arteries); *e*, esophagus; *m*, primitive kidneys (*j*, left, *k*, right); *s*, pylorus; *a*, right yelk-sac; *n*, navel-artery; *u*, navel-vein; *d*, duct; *i*, terminal intestine; 8, fore-limb 9', hind-limb. (After



now inclines from a higher point on the left to a lower point on the right, and continually acquires a more transverse position. In the outer stratum of the stomach-wall, and in the intestinal-fibrous layer, develop the strong muscles which perform the powerful digestive movements. In

the inner stratum, on the contrary, innumerable ~~min~~ glands develop from the intestinal-glandular layer. These are the peptic glands, which supply the most important digestive fluid—the gastric juice. At the lower extremity of the pouch of the stomach a valve develops, which, as the pylorus, separates the stomach from the small intestine (Fig. 275, *d*).

The disproportionately long middle intestine, or small intestine, now develops below the stomach. The development of this section is very simple, and is essentially caused by a very rapid and considerable longitudinal growth. Originally this section is very short, straight, and simple; but immediately below the stomach a horseshoe bend, or loop, begins to appear at a very early period in the intestinal canal, simultaneously with the separation of the intestinal tube from the yolk-sac and with the development of the mesentery. (Cf. Plate V. Fig. 14, *g*, and Fig. 136, vol. i. p. 381.) Before the abdominal wall closes, a horseshoe-shaped loop of intestine (Fig. 136, *m*) protrudes from the ventral opening of the embryo, and into the curve of this the yolk-sac or navel-bladder opens (*n*). The thin, delicate membrane which secures this intestinal loop to the ventral side of the vertebral column, and occupies the inside of this horseshoe curve, is the first rudiment of the mesentery (Fig. 286, *g*). The most prominent part of the loop into which the yolk-sac opens (Fig. 287, *x*), and which is afterwards closed by the intestinal navel, represents that part of the small intestine which is afterwards called the crooked intestine (*ileum*). Soon a very considerable growth of the small intestine is observable; and in consequence, this part has to coil itself in many loops. The various parts of the small intestine which we

have yet to distinguish differentiate later in a very simple way; these are the gall-intestine (*duodenum*), which is next to the stomach, the long empty intestine (*jejunum*) which succeeds, and the last section of the small intestine, the crooked intestine (*ileum*).

The two large glands which we have already named, the liver and the ventral salivary gland, grow out, as protuberances, from the gall-intestine, or duodenum. The liver first appears in the form of two small sacs, situated right and left just behind the stomach (Figs. 284, *f*, 285, *c*). In many low Vertebrates the two livers remain quite separate for a long time (in the *Myxinoidea* for life), and coalesce only imperfectly. In higher Vertebrates, on the other hand, the two livers coalesce more or less completely at an early period, and constitute one large organ. The intestinal-glandular layer, which lines the hollow, pouch-like rudiment of the liver, sends a number of branched processes into the investing intestinal-fibrous layer; as these solid processes (rows of gland-cells) again branch out, and as their branches coalesce, the peculiar netted structure of the developed liver is produced. The liver-cells, as the secreting organs which form the bile, all originate from the intestinal-glandular layer. The fibrous mass of connective tissue, which joins this great cellular network into a large compact organ, and which invests the whole, comes, on the other hand, from the intestinal-fibrous layer. From the latter originate also the great blood-vessels which traverse the entire liver, and the innumerable netted branches of which are interlaced with the network of the liver-cells. The gall-ducts, which traverse the entire liver, collecting the bile and discharging it into the intestine, originate as intercellular passages along

the axis of the solid cell-cords; they all discharge into two primitive main gall or biliary ducts, which originate from the base of the two original protuberances of the intestine. In Man, and in many other Vertebrates, the two ducts afterwards unite, and form one simple gall duct which discharges into the ascending portion of the large intestine. The gall bladder originates as a hollowing or dilatation of the right primitive liver duct. The rate of the growth of the liver is at first exceedingly rapid; in the human embryo, even in the second month, it attains such dimensions that during the third month it occupies by far the largest part of the body-cavity (Fig. 288). At first



FIG. 288.—Chest and abdominal viscera of a human embryo of twelve weeks, in nature (After Koelliker.) The head is omitted; the thoracic and abdominal walls removed. The greater part of the abdominal cavity is occupied by the liver, from an opening in the centre of which the large intestine (caecum, *c*), with the worm appendix, protrudes. Above the diaphragm the lungs are visible in the centre, with the small lung on the right and left.

the two halves are equally well developed; afterwards the left half lies considerably behind the right. In consequence of this asymmetrical development and alteration in the position of the stomach and other abdominal viscera, the whole of the liver is eventually forced over on to the right side. Although the growth of the liver is, afterwards, not so excessive as at the end of gestation, it is comparatively much larger in the embryo than in the adult. In the latter, its

a proportion to that of the whole body is as 1:36; in the former, as 1:18. The physiological significance of the liver during embryonic life—which is very great—depends especially on the part it plays in the formation of blood, and less on its secretion of bile.

From the gall-intestine, immediately behind the liver, grows another large intestinal gland, the ventral-salivary gland, or pancreas. This organ, which occurs only in Skulled Animals, also develops as a hollow sac-shaped protuberance of the intestinal wall. The intestinal-glandular layer of the latter sends out branching shoots, which afterwards become hollow. The ventral-salivary gland, just like the salivary glands of the mouth, develops into a large and very complex gland shaped like a bunch of grapes. The outlet of this gland (*ductus pancreaticus*), through which the pancreatic juice passes into the gall-intestine, seems to be at first simple and single; afterwards it is often double.

The last section of the intestinal tube, the terminal intestine or large intestine (*epigaster*), in mammalian embryos, is, at first, a very simple, short, and straight tube, opening posteriorly through the anus. In the lower Vertebrates it retains this form throughout life. In Mammals, on the other hand, it grows to a considerable size, coils, and differentiates into different sections, of which the foremost and longest is called the colon, the shorter and hinder the rectum. At the commencement of the former a valve (*valvula Bauhini*) forms, which divides the large intestine from the small intestine; behind appears a pouch-like protuberance, which grows larger and becomes the blind-intestine (*cæcum*) (Fig. 288, v). In plant-eating Mammals

this becomes very large, while in those which eat flesh remains very small, or is entirely aborted. In Man, as in most Apes, the beginning of the blind intestine at first becomes wide; its blind end remains very narrow, and afterwards appears only as a useless appendage of the former. This "vermal appendage" is interesting in teleology as a rudimentary organ. Its only importance to Man consists in the fact that now and then a raisin-stone or some other hard, indigestible particle of food becomes lodged in its narrow cavity, causing inflammation, suppuration, and, consequently, killing individuals otherwise perfectly healthy. In our plant-eating ancestors this rudimentary organ was larger, and was of physiological value.

Finally, we must mention another important appendage of the intestinal tube; this is the urinary bladder (*cystis*) with the urinary tube (*urethra*), which in development and in morphological character belong to the intestinal system. These urinary organs, which act as receptacles and excretory passages for the urine secreted by the kidneys, originate from the inner part of the allantois-stalk. The allantois develops, as a sac-like protuberance, from the anterior wall of the last section of the intestine (Fig. 286). In the Dipneusta and Amphibia, in which this blind sac first appears, it remains within the body-cavity (*cælon*) and acts entirely as a urinary bladder. In all Amniota, on the other hand, it protrudes considerably out of the body-cavity of the embryo, and forms the large embryonic "primitive urinary sac," which, in higher Mammals, forms the placenta. At birth this is lost; but the long allantois-stalk (*r*) remains, its upper portion forming the central n

band of the urinary vesicle (*ligamentum vesico-umbilicale medium*), a rudimentary organ which extends as a solid cord from the top of the urinary bladder to the navel. The lower part of the allantois-pedicle (the "*urachus*") remains hollow, and forms the urinary bladder. At first, in Man, as in the lower Vertebrates, this organ discharges into the last section of the posterior intestine, and there is, therefore, a true "cloaca," receiving both urine and excrement; but, among the Mammals, this cloaca is permanent only in the Cloacal Animals, or Monotremes, as in Birds, Reptiles, and Amphibia. In all other Mammals (*Marsupialia* and *Placentalia*) a transverse partition forms at a later period, and separates the urinary-sexual aperture in front from the anal aperture behind. (Cf. Chapter XXV.)

EXPLANATION OF PLATE I.—(FRONTISPIECE.)

DEVELOPMENT OF THE FACE.

The twelve figures in Plate I. represent the faces of four different Mammals in three distinct stages of individual evolution: M_I–M_{III} that of Man, B_I–B_{III} of the Bat, C_I–C_{III} of the Cat, S_I–S_{III} of the Sheep. The three different stages of evolution have been chosen to correspond as far as possible; they have been reduced to about the same size, and are seen from in front. In all the figures the letters indicate the same: *a*, eye; *v*, fore-brain; *m*, mid-brain; *s*, frontal process; *k*, nose-roof; *o*, upper jaw process (of the first gill-arch); *u*, lower jaw process (of the first gill-arch); *l*, second gill-arch; *d*, third gill-arch; *r*, fourth gill-arch; *g*, ear-fissure (remains of the front gill-opening); *z*, tongue. (Cf. Plates VI. and VII. Figs. 232–236, p. 243; also Figs. 123, 124, vol. i. p. 370.)

TABLE XXXVII.

SYSTEMATIC SURVEY OF THE MOST IMPORTANT PERIODS IN THE
PHYLOGENY OF THE HUMAN INTESTINAL SYSTEM.I. *First Period: Intestine of Gastræa* (Figs. 274–277; Plate V. Figs. 9, 10).

The whole intestinal system is a simple pouch (primitive intestine), the simple cavity of which has one orifice (the primitive mouth).

II. *Second Period: Intestine of the Scolecida* (Plate V Fig. 11).

The simple intestinal tube widens in the middle into the stomach, and acquires, at the end opposite to the primitive mouth, a second opening (primitive anus); as in the lower Worms.

III. *Third Period: Intestine of Chorda Animals* (Fig. 281; Plate V. Fig. 12).

The intestinal tube differentiates into two main sections—the respiratory intestine with gill-openings (gill-intestine) in front, the digestive intestine with stomach-cavity (stomach-intestine) behind; as in *Ascidia*.

IV. *Fourth Period : Intestine of Skull-less Animals (Acrania)*

(Fig. 282; Plate XI. Fig. 15).

The gill-streaks appear between the gill-openings of the respiratory intestine; a liver blind-sac grows from the stomach-pouch of the digestive intestine; as in the *Amphioxus*.

V. *Fifth Period : Intestine of Cyclostoma* (Plate XI. Fig. 16).

The thyroid gland develops from the ciliated groove on the base of the gills (hypobranchial groove). A compact liver-gland develops from the liver blind-sac.

VI. *Sixth Period : Intestine of Primitive Fishes* (p. 114).

Cartilaginous gill-arches appear between the gill-openings. The foremost of these form the lip-cartilages and the jaw-skeleton (upper and lower jaw). The swimming-bladder grows from the pharynx. The ventral-salivary gland appears near the liver, as in *Selachii*.

VII. *Seventh Period : Intestine of Dipneusta* (p. 118).

The swimming-bladder modifies into the lungs. The mouth-cavity becomes connected with the nose-cavity. The urinary bladder grows from the last section of the intestine, as in *Lepidosiren*.

VIII. *Eighth Period : Intestine of Amphibia* (p. 126).

The gill-openings close. The gills are lost. The larynx originates from the upper end of the trachea.

IX. *Ninth Period : Intestine of Monotremes* (p. 145).

The primitive mouth and nasal cavity is separated by the horizontal palate-roof into the lower mouth-cavity (food passage) and the upper nose-cavity (air passage); as in all Amnion Animals.

X. *Tenth Period : Intestine of Marsupials* (p. 149).

The existing cloaca is separated by a partition wall into an anterior urinary-sexual aperture and a posterior anal aperture.

XI. *Eleventh Period : Intestine of Catarrhine Apes* (p. 176).

All parts of the intestine, and especially the teeth-apparatus, acquire the characteristic development common to Man and Catarrhine Apes.

CHAPTER XXIV.

DEVELOPMENT OF THE VASCULAR SYSTEM.

Application of the Fundamental Law of Biogeny.—The Two Sides.—Heredity of Conservative Organs.—Adaptation of Progressive Organs.—Ontogeny and Comparative Anatomy complementary of each other.—New “Theories of Evolution” of His.—The “Envelope Theory” and the “Waste-rag Theory.”—Main Germ and Supplementary Germ.—Formative Yolk and Nutritive Yolk.—Phylogenetic Origin of the latter from the Primitive Intestine.—Origin of the Vascular System from the Vascular Layer, or Intestinal-fibrous Layer.—Phylogenetic Significance of the Ontogenetic Succession of the Organ-systems and Tissues.—Deviation from the Original Sequence; Ontogenetic Heterochronism.—Covering Tissue.—Connective Tissue.—Nerve-muscle Tissue.—Vascular Tissue.—Relative Age of the Vascular System.—First Commencement of the Latter; Cœloma.—Dorsal Vessel and Ventral Vessel of Worms.—Simple Heart of Ascidia.—Atrophy of the Heart in the Amphioxus.—Two-chambered Heart of the Cyclostoma.—Arterial Arches of the Selachii.—Double Auricle in Dipneusta and Amphibia.—Double Ventricle in Birds and Mammals.—Arterial Arches in Birds and Mammals.—Germ-history (Ontogeny) of the Human Heart.—Parallelism of the Tribal-history (Phylogeny).

“Morphological comparison of the adult conditions should naturally precede the study of the earliest conditions. Only in this way can the investigation of the history of development proceed in a definite direction; it is thus provided, as it were, to see each step in the formative process in its true relation with the condition which is finally to be reached. Treatment of the history of development without preparatory study is only too

likely to lead to groping in the dark; and it not unfrequently leads to the most unfortunate results—far inferior to those which might be established beyond question without any study of the history of development.”—ALEXANDER BRAUN (1872).

IN applying to Organogeny the fundamental law of Biogeny, we have already afforded some conception of the degree in which we may follow its guidance in the study of tribal history. The degree differs greatly in the different organ-systems; this is so, because the capacity for transmission on one side, and the capacity for modification on the other, vary greatly in the different organs. Some parts of the body cling tenaciously to the inherited germ-history; and, owing to heredity, accurately retain the mode of evolution inherited from primæval animal ancestors; other parts of the body, on the contrary, exhibit very small capacity for strict heredity, and have a great tendency to assume new kenogenetic forms by adaptation, and to modify the original Ontogeny. The former organs represent, in the many-celled community of the human organism, the constant or conservative; the latter, on the contrary, the changeable or progressive element of evolution. The mutual interaction of both elements determines the course of historical evolution.

Only to the conservative organs, in which Heredity preponderates over Adaptation, in the course of tribal evolution, can we directly apply the Ontogeny to the Phylogeny, and can infer, from the palingenetic modification of the germ-forms, the primæval metamorphosis of the tribal forms. In the progressive organs, on the contrary, in which Adaptation has acquired the ascendancy over Heredity, the original course of evolution has, usually, been so changed,

vitiated, and abbreviated, in the course of time, that we can gain but little certain information as to the tribal history from the kenogenetic phenomena of their germ history. Here, therefore, Comparative Anatomy must come to our help, and it often affords much more important and trustworthy disclosures as to Phylogeny than Ontogeny is able to impart. It is, therefore, most important, if the fundamental law of Biogeny is to be correctly and critically applied, to keep its two sides continually in view. The first half of this fundamental law of evolution enables us to use Phylogeny, as it shows us how to gain an approximate knowledge of the history of the tribe from that of the germ: the germ-form reproduces, by Heredity, the corresponding tribal form (Palingenesis). The other half of the law, however, limits this guiding principle, and calls attention to the foresight with which it must be employed; it shows us that the original reproduction of the Phylogeny in the Ontogeny has been in many ways altered, vitiated and abbreviated, in the course of millions of years. The germ-form has deviated, by Adaptation, from the corresponding tribal form (Kenogenesis); the greater this deviation, the more are we compelled to employ Comparative Anatomy in the study of Phylogeny.

Perhaps in no other system of organs of the human body is this so greatly the case as in the vascular system (circulatory, or circulatory apparatus), the development of which we will now examine. If we attempted to infer the original structural features of our older animal ancestors solely from the phenomena which the individual development of these organ-systems, in the embryo of Man and other high Vertebrates, exhibit, we should obtain wh

erroneous views. By many influential embryonic adaptations, among which the development of an extensive nutritive yolk must be regarded as the most important, the original course of development of the vascular system has been so altered, vitiated, and abbreviated, in the higher Vertebrates, that no, or very little, trace of many of the most important phylogenetic features are retained in the Ontogeny. Such explanation as is afforded by the latter would be entirely useless to us if Comparative Anatomy did not lend its aid, and afford us the clearest guidance in our search for tribal history.

Comparative Anatomy is, therefore, especially important in helping us to understand the vascular system, and, equally, the skeleton system, so that, without its guidance, it is unsafe to take a single step in this difficult field. Positive proof of this assertion can be gained by studying the complex vascular system as explained in the classical works of Johannes Müller, Heinrich Rathke, and Karl Gegenbaur. An equally strong negative proof of the assertion is afforded by the ontogenetic works of Wilhelm His, an embryologist of Leipsic, who has no conception of Comparative Anatomy, nor consequently, of Phylogeny. In 1868, this industrious but uncritical worker published certain comprehensive "Studies of the First Rudiment of the Vertebrate Body," which are among the most wonderful productions in the entire literature of Ontogeny. As the author hopes to attain a "mechanical" theory of development by means of a most minute description of the germ-history of the Chick alone, without the slightest reference to Comparative Anatomy and Phylogeny, he falls into errors which are unparalleled in the whole literature of

Biology, rich as this unfortunately is in errors. Only in magnificent germ-history of the Bombinator by Alex. Goette is incomprehensible nonsense and derision of a reasonable causal connection in evolution more nakedly forth. (Cf. vol. i. pp. 65, 66.) His announces, as the result of his investigations, "that a comparatively simple law of growth is the only essential in the first process of evolution. All formation, whether it consist in fissile layers, or in the formation of folds, or in complete articulation, results from this fundamental law." Unfortunately the author does not say in what this all-embracing "law of growth" really consists; just like other opponents of the theory of descent who substitute a great "law of evolution" without telling anything of its nature. From the study of the ontogenetic works of His, on the other hand, it becomes evident that he conceives form-constructing "Mother Nature" merely as a kind of clever dressmaker by cutting out the germ-layers in various ways, by bending, folding, pulling, and splitting them, this clever dressmaker easily brings into existence the various forms of animal species, by "development" (!). The bendings and foldings especially play the most important part. Not only the differentiation of head and trunk, of right and left central stem and periphery, but also the rudiment of limbs, as also the articulation of the brain, the sense-organs, the primitive vertebral column, the heart, and the ear and intestines, can be shown, with convincing necessity (!) to be the mechanical results of the first development of folds. How grotesque is the mode in which the dressmaker proceeds in forming the two pairs of limbs. Their first form is determined by the crossing of four folds bordering the b

like the four corners of a letter." Yet this wonderful "envelope theory" of the vertebrate limbs is surpassed by the "waste-rag theory" (Höllen-lappen Theorie) which His gives of the origin of the rudimentary organs: "Organs like the hypophysis and the thyroid gland) to which no physiological part has yet been assigned, are embryonic remnants, comparable to the clippings, which in the cutting of a dress cannot be entirely avoided, even by the most economical use of the material" (!). Nature, therefore, in cutting out, throws the superfluous rags of tissue into the waste heap. Had our skull-less ancestors of the Silurian age had any presentiment of such aberrations of intellect of their too speculative human descendants, they would certainly have preferred relinquishing possession of the hypobranchial groove on the gill-body, instead of transmitting it to the extant *Amphioxus*, and of leaving a remnant of it to us, in the equally unsightly as useless thyroid gland. (Cf. p. 336).

It will probably be thought that the ontogenetic "discoveries" of His, which appear in a doubly comical light in consequence of the accompanying display of mathematical calculations, can only have occasioned momentary amusement in critical scientific circles. Far from it! Immediately after their appearance, they were not only much praised as the beginning of a new "mechanical" era in Ontogeny, but they have even yet numerous admirers and adherents, who seek to spread the scientific errors of His as far as possible. On this account, I have felt myself obliged to point out emphatically the complete falsity of these views. The vascular system affords especial occasion for this; for among the most important advances which His

claims to have caused by his new conception of g history, is, according to him, his discovery that "the h and tissue of the connective substance" (that is to say, greatest part of the vascular system) "do not originate the two primary germ-layers, as do all the other or but from the elements of the white yelk." The h is designated as "supplementary yelk, or parablast," distinguish it from the "main-germ, or archiblast" germ-disc composed of the two primary germ-layers).

The whole of this artificial development theory of and above all the unnatural distinction between the su mentary and the main germ, collapses like a card l when the Anatomy and Ontogeny of the Amphioxus, invaluable lowest Vertebrate, is contemplated, which can elucidate the most difficult and darkest features in development of the higher Vertebrates, and thus a Man. The gastrula of the Amphioxus alone overtl the whole artificial theory; for this gastrula teach that all the various organs and tissues of complete l brates originally developed entirely from the two pr germ-layers. The developed Amphioxus, like all Vertebrates, has a differentiated vascular system a skeleton of "connective substance tissues" exte throughout its body, and yet there is in this case no plementary germ" from which these tissues can orig thus, contrasting with the other tissues.

The larvæ of the Amphioxus, arising from the or bell-gastrula (*archigastrula*), in its further develop throws the most important rays of light also upon the cult history of development of the vascular system. l first place, it answers the very important question,

have already frequently indicated, as to the origin of the four secondary germ-layers; it clearly shows that the intra-fibrous layer originates from the exoderm, the intestinal-fibrous layer, on the contrary, in an analogous manner, from the entoderm of the gastrula; the cavity thus caused between the two fibrous layers is the first rudiment of the body-cavity, or the coelom (Figs. 50, 51, vol. i. p. 236). As the Amphioxus larva thus shows that the fission of the layers is the same in the lowest Vertebrates as in the Worms, it at the same time represents the phylogenetic connection between the Worms and the higher Vertebrates. As, moreover, the primitive vascular stems in the Amphioxus originate in the intestinal wall, and in this, as in the embryos of all other Vertebrates, proceed from the intestinal-fibrous layer, proof is afforded us that the earlier embryologists were right in calling the latter the vascular layer. Finally, the Comparative Ontogeny of the different vertebrate classes further convinces us that the vascular layer is originally everywhere the same. The vascular system in Man, as in all Skulled Animals, forms a complex apparatus of cavities, which are filled with juices, or fluids, containing cells. The vessels play an important part in the nourishment of the body; some of them conduct the nutritive blood fluid round in the different parts of the body (blood-vessels); some collect the wasted juices and discharge them from the tissues (lymph-vessels). With the latter, the great "serous cavities" are also connected, especially the body-cavity, or coeloma. The heart, acting as a centre of motion for the regular circulation of the juices, is a strong muscular pouch, which contracts in regular pulsations, and is provided with valves, like those of a pump apparatus.

This constant and regular circulation of the blood makes the complex change of substance with the animals possible.

Important as is the vascular system in the most developed and differentiated animal body, it is not, as an apparatus as indispensable to animal life as is supposed. In the older theory of medicine the blood was regarded as the real source of life, and "humoral pathology" referred most diseases to "corrupt blood-mixture." Especially, the blood plays the most important part in the prevailing, obscure conception of Heredity. Just as "pure blood," etc., etc., are yet common phrases, so it is believed that the transmission, by Heredity, of morphological and physiological characters from the parent to the child "lies in the blood." That this conception is entirely false, is easily seen from the facts: neither in the act of procreation is the blood of the parent directly transmitted to the procreated germ, nor do the embryo acquire blood at an early period. As we have already seen, not only the separation of the four germ-layers, but also the beginning of the most important organs, takes place, in the embryos of all Vertebrates, before the rudiment of the vascular systems, of the blood and blood vessels, is formed. In accordance with this ontogenetic fact, we must, from a phylogenetic point of view, regard the vascular system as the most recent, the intestinal system on the contrary, as the oldest formation of the animal body. The origin of the vascular system is, at least, much more recent than that of the intestinal system. If the fundamental principle of Biogeny is rightly appreciated, it is possible, from the ontogenetic sequence, in which the various organs

animal body consecutively originate in the embryo, approximately to infer the phylogenetic sequence, in which these organs gradually developed, one after the other, in the ancestral line of animals. In the "Gastræa theory" I made the first attempt to establish the phylogenetic significance of the ontogenetic sequence of the organ-systems; but it must be remarked that this sequence is not always identical in the higher animal tribes. In Vertebrates, and therefore also in our own ancestral line, the organ-systems may be ranged according to age, in something like the following order: I. The skin-system (*A*) and the intestinal system (*B*). II. The nerve (*C*) and muscular systems (*D*). III. The kidney system (*E*). IV. The vascular system (*F*). V. The skeleton system (*G*). VI. The sexual system (*H*). (Cf. Table XXXIX., p. 367.)

In the first place, the gastrula proves that in all animals with the exception of the Primitive Animals (*Protozoa*),—therefore, in all Intestinal Animals (*Metazoa*),—two primary organ-systems originally arose simultaneously and first; these were the skin-system (skin-covering) and the intestinal system (stomach-pouch). The first is represented, in its earliest and simplest form, by the skin-layer or exoderm, the latter by the intestinal layer or entoderm of the Gastræa. As we can ascribe the same origin, and, therefore, also the same morphological significance, to these two primary germ-layers in all Intestinal Animals, from the simplest Sponge to Man, the homology of these two layers seems sufficient proof of the above assumption.

Immediately after the differentiation of the two primary germ-layers, an inner or outer skeleton develops in many lower animals (*e.g.*, in Sponges, Corals, and other Plant

Animals). In the ancestors of Vertebrates, the development of the skeleton did not take place till much later than in Chorda Animals (*Chordonia*). In them, after the digestive system and the intestinal system, two other organ-systems simultaneously arise; these are the nervous and muscular systems. The way in which these two organ-systems, which mutually condition each other, developed separately and independently, in reciprocal action and in opposition to each other, was first explained by H. H. Kleinenberg in his excellent monograph on the Hydra, a common fresh-water Polyp.¹⁹⁰ In this interesting animal, single cells of the skin-layer send fibre-shaped processes inward, which acquire the power of contraction, capacity, characteristic of the muscles, of contraction in a constant direction. The outer, roundish part of the derm cell remains sensitive and acts as the nervous element; the inner, fibre-shaped part of the same cell becomes contractile, and, incited to contraction by the former, acts as the muscular element (Fig. 293). These are neuro-muscular cells thus still unite in a single cell, of the first order the functions of two organ-systems. A step further; the inner, muscular half of the neuro-muscular cell (Fig. 293, *m*) acquires its own nucleus, and separates from the outer, nervous half (*n*), and both organ-systems have their independent element of form. The fissure between the muscular skin-fibrous layer from the nervous skin-layer in embryonic Worms confirms this important genetic process (Figs. 50, 51, vol. i. p. 236).

These four organ-systems, which have been mentioned, were already in existence, when an apparatus developed secondarily, in the human ancestral line, which,

ght, seems of subordinate significance, but which proves, its early appearance in the animal series and in the embryo, that it must be very ancient and, consequently, of great physiological and morphological value. This is the urinary apparatus, or kidney system, the organ-system which secretes and removes the useless fluids from the body. We have already seen how soon the primitive kidneys appear in the embryo of all Vertebrates, long before any trace of the heart is discoverable. Correspondingly, we also find a pair of simple primitive kidney ducts (the so-called excretory ducts or lymphatic vessels) almost universally diffused in the Worm tribe, which is so rich in forms. Even the lowest classes of Worms, which have as yet neither body-cavity nor vascular system, are furnished with these primitive kidneys (Fig. 280, *nc*, p. 327). It was only in the fourth place, after the kidney system, that the vascular system developed in our invertebrate ancestors; this is plainly shown in the Comparative Anatomy of Worms. The lower Worms (*Acoelomi*) possess no part of the vascular system, no body-cavity, no blood, no heart, and no vessels; this is the case, for example, in the comprehensive group of the Flat Worms (*Plathelminthes*), the Gliding Worms (*Turbellaria*), the Sucking Worms (*Trematoda*), and the Tape Worms. In the higher Worms, which are therefore called *Cœlomati*, a body-cavity (*cœloma*), filled with blood, first begins to form; and, side by side with this, special blood-vessels then also develop. These features have been transmitted from the *Cœlomati* to the four higher animal tribes.

These organ-systems are common to Vertebrates and to the three higher animal tribes, the Articulated Animals

(*Arthropoda*), the Soft-bodied Animals (*Mollusca*), and Star Animals (*Echinoderma*), and we may, therefore, infer that they have all acquired these, as a common inheritance from the Coelomati; but we now meet with a new apparatus of movement, the skeleton system, which, in form, is exclusively peculiar to Vertebrates. Only the first rudiment of this, the simple notochord, is found in *Ascidia*, which are the nearest invertebrate blood-relatives of Vertebrates. We infer from this, that the common ancestors of both, the Chorda Animals, did not branch off from the Worms till a comparatively late period. The notochord is, it is true, one of those organs which appear at a very early period in the vertebrate embryo; but this is clearly due to an ontogenetic heterochronism, to a delay in time in the germ-history, that is, a gradual arrangement in the original phylogenetic sequence, by embryonic adaptation. On Comparative Anatomy grounds it may safely be assumed, that the first of the skeleton system did not precede, but followed the kidney system and of the vascular system, and that Ontogeny appears to indicate the contrary.

Last of all the organ-systems, the sexual system developed, in the sixth place, in our ancestors; of course it must be understood that this was last, in the sense that the sexual apparatus acquired the independent form of a distinct organ-system subsequently to all the other organ-systems. The simplest form, that of reproductive cells, is certainly very ancient. Not only the lowest Worms and Plants propagate sexually, but this was also probably the common parent-form of all Metazoa, in the beginning; but in all these low animals, the reproductive cells

institute special sexual organs in a morphological sense; they are rather, as we shall soon see, simple component parts of other organs.

Like the organ-systems of the human body, the tissues, which compose these systems, are of different ages and of varying morphological value. As we were justified in drawing an inference as to the phylogenetic sequence in the case of the organ-systems, from the ontogenetic sequence in which they successively appear in the embryo, so are we justified in inferring the order in which the tissues originated during the course of tribal history, from the sequence of the stages in germ-history. The result of this is a phylogenetic classification (Table XXXVIII.) of the tissues of the human body, similar to that of the organs (Table XXXIX., p. 367).

The tissues of the human body, arising by division of labour, the separation and the connection of the component cells, may be distributed, with reference to their development, in the four following distinct groups:—1, covering-tissue (*epithelium*); 2, connective tissue (*connectivum*); 3, nerve and muscular tissue (*neuro-musculum*); and 4, vascular tissue (*vasalium*). Of these, in accordance with the Gastraëa theory, we must regard the covering-tissue as the oldest and most original form, as the actual primary or primitive tissue; the three other main forms must, on the other hand, be considered as secondary or derived forms, which developed at a later period from the covering-tissue; the connecting-tissue first, then the neuro-muscular, and lastly the vascular tissue.

The oldest and most original form of tissue is, undoubtedly, the covering-tissue (*epithelium*), the cells of

which are arranged in a simple strata-like way, and cover over the outer and inner surface of the body as a protective and secreting cover. This is proved by the simple fact that the formation of the tissues of the animal body begins with the formation of the gastrula, and that the gastrula itself consists solely of two simple epithelial strata, the skin-layer (Fig. 274, *e*), and of the intestinal-layer. Histologically, the two primary germ-layers are simple epithelia. When these, afterwards, separate into the secondary germ-layers, the skin-sensory layer becomes the outermost of the external coverings (dermal-epithelium), and the intestinal-glandular layer becomes the innermost of the internal coverings (gastral-epithelium). The tissue of the outer skin and of all its appendages, such as nails (Fig.




FIG. 289.—Tissue of the nails (flattened epithelium): *a-e*, cells of the upper strata; *f, g*, cells of the lower strata.

FIG. 290.—Tissue of the covering of the small intestine (coelomic epithelium): *a*, side view of three cells (with thicker, porous borders); *b*, surface view of four cells. (After Frey.)

hairs, skin-glands, etc., arise from the skin-sensory layer. (Cf. Table XXIX., p. 232.) The inner covering of the intestinal tube and of its intestinal glands originates, on the other hand, from the intestinal-glandular layer (Fig. 2

Connective tissue (*connectivum*) must be regarded as forming, in order of phylogenetic age, the second main group of tissues. This is morphologically characterized by the intercellular substance, which develops between the



FIG. 291.—Jelly-like tissue from the vitreous body of an embryo of four months (round cells as jelly-like intercellular substance).

FIG. 292.—Cartilage-tissue of the fibrous or netted cartilage of the ear-shell: *a*, cells; *b*, intercellular mass; *c*, fibres in the latter. (After Frey.)

cells, physiologically, by the double part which it plays, as connecting substance and as complementary substance between the other tissues, as an inner supporting substance and as a protective covering for the inner organs. Of the numerous forms and varieties of connective tissue, we regard the jelly-like tissue (Fig. 291; Fig. 6, vol. i. p. 126), the fatty tissue, and the chorda tissue as the earlier; the fibrous, cartilaginous (Fig. 292), and bone-tissue (Fig. 5, vol. i. p. 126) as the more recent formations. All these various forms of connective tissue are products of the middle germ-layer, or mesoderm; or, more accurately, of the two fibrous layers, of which the skin-fibrous layer is originally derived from the exoderm, the intestinal-fibrous layer from the entoderm.

The nerve-muscular tissue (*neuro-musculum*) is of much more recent origin than the connective tissue. If epithelial tissue represents a primary period in tribal history, and

connective tissue a secondary period, then we may characterize a third, much later period, by nerve-muscle tissue.




FIG. 293.




FIG. 295.




FIG. 294.

FIG. 293.—Nerve-muscle tissue. Three cells from *Hydra*: *n*, nervous; *m*, inner, muscular part of the cells. (After Kleinenberg.)

FIG. 294.—Nerve-tissue (from a spinal nerve knot): *a*, anterior root of the spinal nerve; *d, e*, fibrous nerve-stem; *f, g, h*, cells in ganglion (*f*, unipolar, *g, h*, bipolar cells); *k, l*, nerve fibres. (After Frey.)

FIG. 295.—Muscle-tissue. Three pieces of striped muscle fibre (interfibrillar fat-cells (*b*)). (After Frey.)

For while in the lowest Plant Animals the body is covered merely of covering tissue, and while in many

Phytes a middle layer of connective tissue develops between the two primary germ-layers, it is only in the highly developed Plant Animals that muscle and nerve is formed. As has already been said, the latter first appeared as a common nerve and muscle tissue (*neuroculum*, Fig. 293; cf. p. 358). It was only afterwards the muscle-tissue (Fig. 295) separated from the nerve-tissue (Fig. 294). The greater part of the nerve-tissue is derived from the skin-sensory layer, the greater part of the muscle-tissue from the skin-fibrous layer.

Vascular tissue (*vasalium*) must be regarded as forming



FIG. 296.—Vascular tissue (*vasalium*). A hair-vessel from the enteric: *a*, vascular cells; *b*, the kernels of these ("endothelium").

FIG. 297.—Red blood cells (corpuscles) of various Vertebrates (equally magnified): 1, Human; 2, Camel; 3, Pigeon; 4, *Proteus* (p. 129); 5, Water-mander (*Triton*); 6, Frog; 7, Fish (*Cobitis*); 8, Lamprey (*Petromyzon*). *a*, surface view; *b*, edge view. (After Wagner.)

TABLE XXXVIII.

Systematic Survey of the Sequence, according to Age, of the Hms
Tissue-groups.

(Phylogenetic Classification of Vertebrate Tissues.)

FIRST GROUP: PRIMARY TISSUES (*Epithelium*).

1. FIRST HISTOLOGICAL STAGE OF EVOLUTION.

I. Covering-tissue (*Epithelium*).

- | | | |
|---|---|---|
| I. A. Skin-covering tissue (<i>Epithelium dermale</i>).
Skin-layer, or Exoderm, of Gastrula (after-
wards skin-sensory layer) | { | 1. Outer skin (<i>Epi-dermis</i>)
2. Glands of outer skin
3. (Earliest site of origin of the germ-cells?) |
| I. B. Intestinal covering tissue (<i>Epithel. gastrale</i>).
Intestinal layer, or Entoderm, of Gastrula
(afterwards the intestinal-glandular layer) | { | 1. Real intestinal epithelium
2. Epithelium of the intestinal glands
3. (Earliest site of origin of egg-cells?) |

SECOND GROUP: SECONDARY TISSUES.

(All derived from the Covering-tissue, or *Epithelium*.)

2. SECOND HISTOLOGICAL STAGE OF EVOLUTION.

II. Connective-tissue (*Connectivum*).

- | | | |
|--|---|---|
| II. C. Filling-up tissue (<i>Tela conjunctiva</i>) (softer
[surrounding] connective tissue) | { | 1. Jelly-like tissue
2. Fatty tissue
3. Fibrous tissue |
| II. D. Supporting tissue (<i>Tela skeletaria</i>) (firmer
[supporting] connective tissue) | { | 4. Chorda tissue
5. Cartilaginous tissue
6. Bone-tissue |

3. THIRD HISTOLOGICAL STAGE OF EVOLUTION.

III. Nerve-muscle Tissue (*Neuro-musculum*).

- | | | | | |
|--|---|--|---|---|
| III. E. Nerve-tissue (<i>Tela nervea</i>). Original outer
portion of the nerve-
muscle cells of the
Exoderm | { | 1. Nerve-cells
(Ganglion-cells)
2. Nerve-fibres
(Nerve-tubes) | { | 1. a. Periphere nerve-cells (Red-cells of the sense-organs)
1. b. Central nerve-cells (mind-cells)
2. a. Sheath-less nerve-fibres (pink, medulla-less fibres)
2. b. Sheathed nerve-fibres (dark fibres with medulla) |
| III. F. Muscle-tissue (<i>Tela muscularis</i>). Original
inner portion of the
nerve-muscle cells of
the Exoderm | { | 1. One-celled muscle-fibres
2. Many-celled muscle-fibres | { | 1. a. Smooth contractile fibre-cells
1. b. Striped contractile fibre-cells
2. a. Smooth muscle-masses
2. b. Striped muscle-masses |

4. FOURTH HISTOLOGICAL STAGE OF EVOLUTION.

IV. Vascular Tissue (*Vasallum*).

- | | | | | |
|---|---|---|---|---|
| IV. G. Vascular lining tissue (<i>Tela vasalis</i>). Inner
wall-covering of the
Coelom system | { | 1. Coelarium
(Coelom - epithe-
lium)
2. Endothelium
(Vascular epithe-
lium) | { | 1. a. Exocoelarium (Parietal Coelom-epithelium) (and secondary site of origin of the sperm-cells?)
1. b. Endocoelarium (Visceral coelom-epithelium) (and secondary site of origin of the egg-cells?)
2. a. Endothelium of the lymph-vessels
2. b. Endothelium of the blood-vessels |
| IV. H. Lymph-tissue (<i>Tela lymphatica</i>). Liquid
contents of the Coelom
system | { | 1. Lymph (Colourless blood-cells and fluid intercellular substance)
2. Blood (Red blood-cells and fluid intercellular substance) | | |

TABLE XXXIX.

Systematic Survey of the Sequence, according to Age, of the Human Organ-systems.

(Phylogenetic Classification of Vertebrate Organs.)

(On the right are given the Ancestral Stages, in which the respective organs probably first appeared.)

1. FIRST STAGE IN THE EVOLUTION OF ORGANS.

I. Skin and Intestinal Systems.

The two Systems appear first, and simultaneously, in the Gastread ancestors.

A. Skin-system (<i>Systema dermale</i>)	{	A 1. Simple exoderm	Gastreaads
		A 2. Outer skin (Skin-sensory layer) and leather skin (Skin-fibrous layer)	Worms
		A 3. Outer skin, with hairs, glands, etc.	Mammals
B. Intestinal system (<i>Systema gastrale</i>)	{	B 1. Simple entoderm	Gastreaads
		B 2. Intestinal epithelium (Intestinal-glandular layer) and intestinal muscular skin (Intestinal-fibrous layer)	Worms
		B 3. Gill-intestine and stomach-intestine	Chorda-animals

2. SECOND STAGE IN THE EVOLUTION OF ORGANS.

II. Nerve and Muscle Systems.

The two Systems appear first, and simultaneously, in the Primitive Worm ancestors.

C. Nerve-system (<i>Systema nerveum</i>)	{	C 1. Upper throat ganglia	Primitive Worms
		C 2. Simple medullary tube	Chorda-animals
		C 3. Brain and spinal marrow	Monorhina
D. Muscle-system (<i>Systema musculare</i>)	{	D 1. Skin-muscle pouch	Primitive Worms
		D 2. Side muscles of the trunk	Acrania
		D 3. Trunk and limb muscles	Fishes

3. THIRD STAGE IN THE EVOLUTION OF ORGANS.

III. Kidney and Vascular Systems.

The two Systems first appear, one after the other, in the Soft-worm ancestors (*Scotecida*).

E. Kidney-system (<i>Systema renale</i>)	{	E 1. Primitive kidney canals	Scotecida
		E 2. Segmental canals	Acrania?
		E 3. Primitive kidneys	Monorhina
		E 4. Permanent kidneys	Protamnina
F. Vascular system (<i>Systema vasculare</i>)	{	F 1. Simple coelom	Scotecida
		F 2. Dorsal and ventral vessels	Worms
		F 3. Heart (part of the ventral vessel)	Chorda-animals
		F 4. Heart, with auricle and ventricle	Monorhina

4. FOURTH STAGE IN THE EVOLUTION OF ORGANS.

IV. Skeleton and Sexual Systems.

The two Systems first appear, one after the other, in the Chordonia-ancestors.

G. Skeleton-system (<i>Systema skeletare</i>)	{	G 1. Simple notochord	Chorda-animals
		G 2. Cartilaginous primitive skull	Monorhina
		G 3. Gill-arches, ribs, limbs	Selachii
		G 4. Limbs, with five digits	Amphibia
H. Sexual system (<i>Systema sexuales</i>)	{	H 1. Simple hermaphrodite glands	Chorda-animals
		H 2. Distinct testes and ovaries	Acrania
		H 3. Seed-duct and oviduct	Selachii
		H 4. Phallus (penis, clitoris)	Protamnina

the most recent group of tissues, that which originated last. Under this name are included those epithelial-like tissues which line the closed inner cavities of the body (the coelom, chest-cavity, ventral cavity, heart-cavity, blood-vessels, etc.) (Fig. 296). In addition to this vascular carpet (endothelium), the liquids containing cells, which fill these cavities (lymph, blood, serum, etc.), must be classed with this tissue (Fig. 297). All these tissues may be grouped as *vasalia*. His wrongly ascribed to them a quite different "parablastic" origin (from the nutritive yolk); they are, however, products of the intestinal-fibrous layer (and partly, perhaps, of the skin-fibrous layer). As the coeloma and the whole vascular system is of more recent phylogenetic origin its peculiar tissues must also be more recent.

This phylogenetic explanation of the ontogenetic succession of the tissues and of the organ systems arising from them, appears to me to be satisfactorily proved by Comparative Anatomy, and by the Gastræa theory. If it is correct, it discloses an interesting glimpse into the entire various age of the most important constituent parts of the body. The human skin and intestine are, according to the theory, many thousands of years older than the muscles and nerves; these again are much more ancient than kidneys and blood-vessels, and the latter, finally, are many thousands of years older than the skeleton and the sexual organs. The common view, that the vascular system is one of the most important and original organ-systems, is, therefore, erroneous; it is as false as the assumption of Aristotle that the heart is the first part to form in the incubated chick. On the contrary, all lower Intestinal Animals show plainly that the historic evolution of the vascular system did

till a comparatively late period. Not only all Plantals (Sponges, Corals, Hydropolyps, Medusæ), but also lower Worms (*Acoelomi*), are entirely destitute of a vascular system. In both groups, the fluid acquired by digestion is conveyed directly from the intestinal tube, through processes of this latter (the gastro-canals), into the distant parts of the body. It is only in the intermediate higher Worms that the vascular system first begins to appear, in consequence of the formation of a simple cavity (*coeloma*), or of a system of connected spaces, round the intestinal tube, in which cavities the nutritive fluid (blood) after passing through the intestinal wall, collects.

In the human ancestral line we meet with this first element of the vascular system in that group of Worms which we spoke of as Soft Worms (*Scolecida*; p. 85). Soft Worms, as we said, formed a series of intermediate stages between the lowest bloodless Primitive Worms (*helminthes*) and the Chorda-worms (*Chordonia*), which were already provided with a vascular system and a notochord. The vascular system must have begun, in the older *Scolecida*, with a very simple coelom, a "body-cavity," filled with blood, and which surrounded the intestinal tube. Its origin was probably due to the accumulation of nutritive fluid in a cleft between the intestinal-fibrous layer and the skin-fibrous layer. A vascular system in its simplest form is yet found in the Moss-polyps (*Bryozoa*) and the Wheel-animalcule (*Rotatoria*), and in other lower forms. The inner, visceral, part of the wall of the coelom is naturally formed by the intestinal-fibrous layer (*endoderm*), the outer, parietal, part by the skin-fibrous layer (*ectoderm*). The coelom fluid, collected between the two,

may contain detached cells (lymph-cells) from either layer.

A first advance in the development of this most primitive vascular system was accomplished by the formation of canals or blood-conducting tubes, which developed independently of the coeloma, in the intestinal wall, that is, in the intestinal-fibrous layer of the wall. These real vessels, in the stricter sense, appear in very different form in Worms of the intermediate and higher groups. Sometimes they are very simple, sometimes very complex. Two primordial "primitive vessels" must be regarded as representing that form, which probably formed the basis of the more complex vascular system of Vertebrates; these are a dorsal vessel, which passes from front to back along the middle line of the dorsal wall of the intestine, and a ventral vessel which passes, in the same direction, along the middle line of the ventral wall. Both at the front and at the back these two vessels are linked together by a loop encircling the intestines. The blood enclosed in these tubes is driven forward by the peristaltic contractions of the intestine.

The further development of this simplest rudimentary blood-vessel system is evident in the class of the higher Worms (*Annelida*), in which we find it in very many stages of development. In the first place, many transverse connections probably arose between the dorsal and ventral vessels, so as to encircle the intestine (Fig. 1). Other vessels then penetrated into the body-wall, and became branched, so as to conduct blood to this part. As in the ancestral Worms, which we have called Chordones, the front section of the intestine changed into a gill-body

lar loops, within the wall of this gill-body, which led from the ventral vessel to the dorsal vessel, became modified into respiratory gill-vessels. Even at the present time the organization of the remarkable Acorn-worm (*Acornoglossus*) exhibits a similar condition of gill-circulation (Fig. 186, p. 86).

A further important advance is exhibited, among extant Worms, in the Ascidia, which may be regarded as the nearest blood-relation to our primitive Chordonia ancestors. Here we find, for the first time, a real heart, that is, a *central organ of the circulation of the blood*, by the pulsating contractions of the muscular wall of which the blood is driven forward in the vascular tubes. The heart appears here in the simplest form, as a spindle-shaped pouch which passes at both ends into a main vessel (Fig. 188, c. p. 90; XI. Fig. 14, *hz*). The original position



Fig. 298.—Blood-vessel system of a Ringed Worm (*Polychaeta*); front section: *d*, dorsal vessel; *v*, ventral vessel; *c*, transverse connection between the two (enlarged like a heart). The arrows indicate the direction of blood current. (After Gegenbaur.)

The heart on the ventral side, behind the gill-body of the Ascidian, plainly shows that it originated in a local dilation of a section of the ventral vessel. The alternating direction of the movements of the blood, which has already been mentioned, is remarkable; the heart expels the blood alternately through the anterior and through the posterior end, a fact which is very suggestive, because in most Worms the blood

in the dorsal vessel moves from back to front, in Vertebrates, on the contrary, it flows in the opposite direction, from front to back. As the heart of the *Ascidia* constantly alternates between these two opposite directions it exhibits permanently, to a certain extent, the phylogenetic transition between the older direction of the dorsal current toward the front in Worms, and the newer direction of the same toward the rear in Vertebrates.

As in the more recent Chorda Animals, which give rise to the Vertebrate tribe, the newer direction becomes permanent, the two vessels which proceeded from two ends of the heart-pouch, acquired a constant difference. The front section of the ventral vessel, since it has steadily conducted the blood from the heart, consequently, as an artery; the hinder section of the ventral vessel, on the contrary, leads the blood, circulating in the body, back into the heart, and must, therefore, be called a vein. In reference to their relation to the different sections of the intestine, we may speak of the latter more accurately, as the intestinal vein, and of the former as the gill-artery. The blood contained in both vessels, when alone fills the heart also, is venous blood; that is, contains much carbonic acid. On the other hand, the blood which flows from the gills into the dorsal vessel is the arterial blood, furnished with oxygen; is arterial blood. The most distal branches of the arteries and veins pass into each other within the tissue, through a network of extremely fine neutral hair-vessels or capillaries (Fig. 296).

If we now turn from the *Ascidia* to the nearest form, the *Amphioxus*, we are immediately surprised by an apparent retrogression in the development of the va-

system. The *Amphioxus*, as has been stated, has no real heart; but the blood is circulated in its vascular system by the main vascular stems themselves, which contract and dilate along their whole length. (Cf. Fig. 151, vol. i. p. 420.) The dorsal vessel (aorta), situated over the intestine, absorbs the arterial blood from the gills and propels it through the body. The venous blood, in its return, collects in a ventral vessel (intestinal vein), situated under the intestine, and thus returns to the gills. Numerous vascular gill-arches, which accomplish respiration, and absorb oxygen from the water and emit carbonic acid, unite the ventral vessel with the dorsal vessel before. As, in *Ascidia*, that section of the ventral vessel which also forms the heart in Skulled Animals (*Craniota*), is already fully developed into a simple heart-pouch, we must regard the absence of the latter in the *Amphioxus* as the result of retrogression, as a reversion, in these Acrania, to the older form of vascular system, as it exists in *Scolecida* and many other Worms. We may assume that those Acrania which actually formed part of our ancestral line did not share this relapse, but rather inherited the one-chambered heart from the Chordonia and transmitted it directly to the older Skulled Animals (*Craniota*).

The Comparative Anatomy of Skulled Animals clearly exhibits the further phylogenetic development of the blood-vessel system. In the lowest stage of this group, in the *Cyclostoma* (p. 102), we first meet with a real lymph-vessel system, side by side with the blood-vessel system, a system of canals which collect the colourless fluid flowing from the tissues, and conduct it to the blood-current. Those lymph-vessels which absorb the milky, nutritive fluid, obtained

directly by digestion, from the intestinal wall, and conducted to the blood-current, are distinguishable as chyle-vessels or "milky juice vessels." While the chyle, or milky juice, in consequence of the great amount of fat globules which it contains, appears milk white, the real lymph is colourless. The chyle, as well as the lymph, contain the same colourless amœboid cells (Fig. 9, vol. i. p. 132), which are distributed in the blood as colourless blood-cells (corpuscles); the latter contains, in addition, the much greater quantity of red blood-cells (corpuscles), which gives the blood of Skulled Animals its red colour. The distinction, common to all Craniota, between lymph-vessels, chyle-vessels, and blood-vessels, is to be regarded as the result of a division of labour which took place between different portions of an original unitary, primitive blood-vessel system (or hæmolymph system).

The heart, the central organ of the circulation of the blood, which exists in all Craniota, also exhibits an advance in structure, even in the Cyclostoma. The simple spindle-shaped heart-pouch is separated into two divisions, or chambers, which are divided by two valves (Plate XI. Fig. 16, *hv*, *hk*). The posterior division, the fore chamber (*atrium*, *hv*), absorbs the venous blood from the veins of the body, and discharges it into the anterior division, the chamber, or main chamber (*ventriculus*, *hk*). From here it is propelled by the gill-artery stem (the foremost section of the ventral vessel) into the gills.

In Primitive Fishes (*Selachii*), an arterial stalk (*bulbus arteriosus*), separated by valves, originates, as a distinct section, from the foremost end of the ventricle. It forms the enlarged, hindmost end of the gill-artery stem (Fig.

abr). From each side of this, from five to seven gill-arteries proceed; these rise between the gill-openings (*s*) the gill-arches, encircle the throat, and combine above to a common aorta-stem, the continuation of which, going backward above the intestine, corresponds to the dorsal vessel of Worms. As the arched arteries distribute themselves in a respiratory capillary net over the gill-arches, they thus contain venous blood in their lower part (as arterial gill-arches), and arterial blood in their upper part (as aorta-arches). The points at which separate aorta-arches unite, which occur on the right and left sides, are called aorta-roots. Of an originally greater number of aorta-arches, only five pairs are retained, and from these (Fig. 300), in all higher Vertebrates, the most important parts of the arterial system develop.



FIG. 299.—Head of an embryonic Fish, with the rudiment and the blood-vessel system; seen from the left side: *dc*, Cuvierian duct (point of union of the front and hind main veins); *sv*, venous sinus (enlarged anterior portion of the Cuvierian duct); *a*, auricle; *v*, main chamber; *s*, gill-openings (between the arterial arches); *ad*, dorsal vessel; *c*, head-artery (carotis); *n*, nose-groove. (After Gegenbaur.)

The appearance of the lungs, connected with the respiration of air, which first occurs in the Dipneusta, is most important in the further development of the arterial

system. In *Dipneusta*, the auricle of the heart separates into two halves by the formation of an incomplete partition. Only the right auricle now absorbs the venous blood from the body-veins. The left auricle, on the other hand, absorbs the arterial blood of the lung-veins; both auricles then discharge in common into the simple ventricle, in which the two kinds of blood mingle, and are then propelled through the arterial stalk into the arterial arches. From these latter spring the lung-arteries (Fig. 301, *p*); they convey a part of the mixed blood into the lungs, while the remainder is driven through the aorta into the body.

From the *Dipneusta* upward, we trace a progressive development of the vascular system, which finally results with the loss of gill respiration, to a complete separation of the two parts of the double circulatory system. In *Amphibia*, the partition between the two auricles becomes complete. In their young form, these yet retain gill respiration and the circulatory system as in *Fishes*, as the heart contains only venous blood; at a later period, the lungs, with their vessels, are developed also, and the main chamber of the heart then contains mixed blood. In *Amphibia*, *Amphibia* and *Reptiles*, the main chamber and the arterial stalk belonging to it begin to separate, by the formation of a longitudinal partition, into two halves, and this partition becomes complete in the higher reptiles on the one side, and in the parent-form of *Mammals* on the other. The right auricle of the heart alone now contains venous blood, the left contains only arterial, as in all *Birds* and *Mammals*. The right auricle receives venous blood from the body-veins, and the right ventricle propels this through the lung-arteries into the lungs; from there it returns as arterial blood through the arterial stalk into the arterial arches.

the lung-veins to the left auricle, and is driven through the left ventricle into the body-arteries. Between the lung-arteries and lung-veins is situated the capillary system of the lesser, or lung-circulation; between the body-arteries and the body-veins lies the capillary system of the greater, or body-circulation. Only in the two highest Vertebrate



FIG. 300.—The five arterial arches of Skulled Animals (1-5) in their original form: *a*, arterial stalk; *a''*, main stem of the aorta; *c*, head-artery (carotis, anterior continuation of the aorta-roots). (After Rathke.)


FIG. 301.—The five arterial arches of Birds; the light portions of the rudiment disappear; only the dark parts are permanent. Letters as in Fig. 300: *s*, arteries of the clavicula (sub-clavian); *p*, lung-artery; *p'*, branches of the same. (After Rathke.)

FIG. 302.—The five arterial arches of Mammals. Letters as in Fig. 301: *v*, vertebral artery; *b*, Botalli's duct (open in the embryo, afterwards closed). (After Rathke.)

classes, in Birds and Mammals, is this complete separation of the two courses of the circulation perfect. Moreover, this separation has taken place in the two classes independently of each other, as is shown by the unequal development of the aortas. In Birds, which are descended from Reptiles,

the right half of the fourth arterial arch has become permanent arterial arch (*arcus aortæ*, Fig. 301). (On the other hand, the latter has developed from the left half of the same arch (Fig. 302) in Mammals, which are descended from the Protamnia.

On comparing the arterial system in the various classes of the Skulled Animals (*Craniota*) in its matured condition it appears in very various forms, and yet it develops all, from the same primitive form. This development takes place in man exactly as in other Mammals; especially the modification of the five arterial arches precisely the same in both cases (Figs. 303–306). At first, only a single



FIGS. 303–306.—Metamorphosis of the five arterial arches in the embryo (diagram after Rathke): *ta*, arterial stalk; 1, 2, 3, 4, 5, the arches from the first to the fifth pair; *ad*, main stem of the aorta. In Fig. 303, three of the arterial arches are given; in Fig. 304, the whole five (those indicated by dots are not yet developed); in Fig. 305, the first two have again disappeared; in Fig. 306, the permanent arches are represented. The dotted parts disappear. *s*, Sub-clavian artery; *v*, vertebral artery; *ax*, axillary artery; *c*, carotid artery (*c'*, inner carotis); *p*, pulmonary artery (lung-artery).

arches develop, and these lie on the inner surface of the first pair of gill-arches (Figs. 147–150, vol. i. pp. 39–40, Fig. 303). A second and a third pair of arches then

ind the first, and these are situated on the inner surface of the second and third gill-arches. At length, a fourth and fifth pair appear behind the third (Fig. 304); but while the latter are developing, the first two are again disappearing by growing together (Fig. 305). The permanent main series develop only from the three posterior arterial arches (3, 4, 5, in Fig. 304), the lung-arteries from the last (Fig. 306). (Cf. with this Fig. 302.)

The human heart also (Fig. 314) develops exactly like that of other Mammals. We have already considered the first principles of its germ-history (vol. i. pp. 392-395, Figs. 143-147), which essentially corresponds with its Phylogeny.¹⁹¹ We saw that the very first rudiment of the heart is a spindle-shaped thickening of the intestinal-fibrous layer in the ventral wall of the head-intestine (Fig. 143, *df*). This spindle-shaped invagination then becomes hollow, forms a simple pouch, and separates from the place at which it originated, so that it then lies freely in the cardiac cavity (Figs. 145, 146). This pouch bends into the form of an S (Fig. 144, *c*), and, at the same time, turns spirally on an imaginary axis, so that the posterior part lies on the dorsal surface of the anterior part. The combined yolk-veins open into its posterior extremity; from the anterior extremity proceed the arterial arches (Fig. 150, vol. i. p. 398).

This first rudiment of the human heart, which encloses a very simple cavity, corresponds to the heart of the Ascidians, and must be regarded as a reproduction of the heart of the Chordonia; it now, however, separates into two, and then into three parts, thus exhibiting for a very brief period the heart-structure of the Cyclostoma and of Fishes. The spiral turn and curve of the heart increases, and, simultaneously,

two shallow transverse indentations of the circumference appear, which externally mark the three sections (Fig. 308). The anterior section, which is turned toward




FIG. 307.—Heart of an embryonic Rabbit, from behind : *a*, fore-brain; *b*, auriculæ; *c*, auricle (atrium); *d*, ventricle; *e*, artery-stalk; *f*, three pairs of arterial arches. (After Bischoff.)

FIG. 308.—Heart of the same embryo (Fig. 307), from the yolk-veins; *a*, auricle; *ca*, auricular canal; *l*, left ventricle; *v*, ventricle; *ta*, artery-stalk. (After Bischoff.)

FIG. 309.—Heart and head of an embryonic Dog, from the front; *a*, fore-brain; *b*, eyes; *c*, mid-brain; *d*, primitive lower jaw; *e*, upper jaw; *f*, gill-arches; *g*, right auricle; *h*, left auricle; *i*, left ventricle; *k*, right ventricle. (After Bischoff.)

FIG. 310.—Heart of the same embryo, from behind : *a*, entrance of yolk-veins; *b*, left auricular process; *c*, right auricular process; *e*, auricular canal; *f*, left ventricle; *g*, right ventricle; *h*, aorta. (After Bischoff.)

ventral side, and from which the aortal arches reproduce the arterial stalk (*bulbus arteriosus* Selachii). The central section is the rudiment of a chamber, or ventricle (*ventriculus*); and the anterior section, the one turned toward the dorsal side, into which the yolk-veins open, is the rudiment of a simple

atrium). The latter, like the simple auricle of the heart of the Fish, forms a pair of lateral protuberances, the heart s, or auricular appendages (*auriculæ*, Fig. 307, *b*); and hence the indentation between the auricle and ventricle is called the auricular canal (*canalis auricularis*, Fig. 308, *ca*). The heart of the human embryo is now a complete Fish heart.

Corresponding exactly with the Phylogeny of the human heart (Table XLI.), its Ontogeny exhibits a gradual transition from the Fish heart through the Amphibian heart to the Mammalian heart. The most important step in this advance is the formation of a longitudinal partition, imperfect at first, afterwards complete, by which all the three sections of the heart are separated into a right (venous) and a left (arterial) half. (Cf. Figs. 309-314.) The auricle (*atrium*) is thus divided into a right and a left auricle, each of which acquires its respective auricular process; the body-veins discharge into the right auricle (ascending and descending *vena cavae*, Fig. 311, *c*, Fig. 313, *c*); the left auricle receives the lung-veins. Similarly, a superficial "inter-ventricular furrow" (*sulcus interventricularis*, Fig. 312, *s*) appears at an early period on the main chamber of the heart, the external expression of the internal partition, by the formation of which the ventricle is divided into two chambers, a right (venous) and a left (arterial) ventricle. Finally, a longitudinal partition forms, in a similar way, in the third section of the primitive heart, which so much resembles that of a Fish, in the arterial stalk, which is also externally indicated by a longitudinal furrow (Fig. 312, *af*). This separates the cavity of the artery-stalk into two lateral halves; the main lung artery, which opens into the



FIG. 311.



FIG. 312.



FIG. 313.




FIG. 314.

FIG. 311.—Heart of a human embryo of four weeks; 1, from the front; 2, from the back; 3, open, and with the upper half of the auricle removed. *a'*, left auricular process; *a''*, right auricular process; *v'*, left ventricle; *v''*, right ventricle; *ao*, artery-stalk; *c*, upper hollow vein (*vena cava*) on the right; *cs*, left; *s*, rudiment of the partition, between the chambers. (After Koelliker.)

FIG. 312.—Heart of a human embryo of six weeks, from the front. *r*, right ventricle; *t*, left ventricle; *s*, furrow between the two ventricles; *ta*, artery-stalk; *af*, furrow on its surface; *a*, at the right and left are the two large auricular processes of the heart. (After Ecker.)

FIG. 313.—Heart of a human embryo of eight weeks, from the front. *a'*, left auricular process; *a''*, right auricular process; *v'*, left ventricle; *v''*, right ventricle; *cd'*, right upper *vena cava*; *cs*, left upper *vena cava*; *ci*, lower *vena cava*. (After Koelliker.)

FIG. 314.—Heart of human adult, perfectly developed, from the front in its natural position: *a*, right auricular process (below it, the right ventricle); *b*, left auricular process (below it, the left ventricle); *C*, upper *vena cava*; *V*, lung-veins; *P*, lung-artery; *d*, Botalli's duct; *A*, aorta. (After Me

right ventricle, and the aorta-trunk, which opens into the left ventricle. Not until all these partitions are completed is the lesser, or lung-circulation, entirely distinct from

water, or body-circulation; the right half of the heart is the centre of motion for the former, the left half for the latter. (Cf. Table XLI.)

In the human embryo, and in all other Amniota, the heart originally lies far forward on the lower side of the head, as in Fishes it remains permanently near the throat. Afterwards, with the advancing development of the neck and chest, the heart continually moves further back, until at last it is situated in the lower part of the breast between the lungs. At first its position is symmetrical, in the central plane of the body, so that its longitudinal axis corresponds with that of the body (Plate IV, Fig. 8). In most mammals it retains this symmetrical position permanently; but in the Apes the axis begins to incline obliquely, and to move the apex of the heart to the left side. This inclination is carried furthest in the Man-like Apes; in the Chimpanzee, Gorilla, and Orang, which also resemble Man in this oblique position of the heart.

The germ-history of all other parts of the vascular system, like that of the heart, point out many and valuable facts regarding the history of our descent. But as an accurate knowledge of the complex arrangement of the entire vascular system of Man and other Vertebrates is required, in order to follow the matter sufficiently far to make it intelligible, we cannot here enter into any further detail.¹⁹² Moreover, many important features in the Ontogeny of the vascular system, especially in regard to the derivation of its various parts from the secondary germ-layers, are as yet very obscure and doubtful. This is true, for example, of the question as to the origin of the coelom-epithelium—that is, of the cell-layer coating the body-cavity. Probably there is an important phylogenetic

distinction between the exocoelar, or the parietal coelom-epithelium, which originates from the skin-fibrous layer, and the endocoelar, or the visceral coelom-epithelium, which is derived from the intestinal-fibrous layer. The former is, perhaps, connected with the male germ-epithelium (the rudiment of the testes), the latter with the female germ-epithelium (the rudiment of the ovary). (Cf. Chapter XII.)

TABLE XL.

SYSTEMATIC SURVEY OF THE MOST IMPORTANT PERIODS IN THE PHYLOGENESIS OF THE HUMAN VASCULAR SYSTEM.

I. *First Period: Vascular System of the earlier Scolecida.*

Between the skin-covering and the intestinal wall is formed a sub-body-cavity (*cœloma*), or a perienteric cavity (as in the extant *Bryozoa* and other *Cœlomati*).

II. *Second Period: Vascular System of the more recent Scolecida.*

The first real blood-vessels form in the intestinal wall (in the intestinal fibrous layer), a dorsal vessel in the central line of the dorsal side of the intestinal tube, and a ventral vessel in the central line of its ventral side. The two vessels are connected by several circular vessels, encircling the intestine.

III. *Third Period: Vascular System of the earlier Chordonia.*

By the modification of the anterior half of the intestine into a pharynx, the anterior section of the ventral vessel becomes a gill-artery, and the anterior section of the dorsal vessel a gill-vein; between them a gill capillary network develops.

IV. *Fourth Period: Vascular System of the more recent Chordonia.*

The portion of the ventral vessel, lying immediately behind the pharynx, enlarges to a simple heart-pouch (Ascidian).

V. *Fifth Period: Vascular System of the Acrania.*

The ventral vessel (intestinal vein) forms, round the developing liver, the first rudiment of a vena portæ system.

VI. *Sixth Period: Vascular System of the Cyclostomi.*

The single-chambered heart divides into two chambers; a posterior tricle, and an anterior auricle. The lymph-vessel system develops side by side with the blood-vessel system.

VII. *Seventh Period: Vascular System of the Primitive Fishes, or Selachii.*

From the anterior section of the main chamber of the heart arises an artery-stalk or trunk, from which five (?) pairs of arterial arches proceed.

VIII. *Eighth Period: Vascular System of the Mud-fishes.*

From the last (fifth) pair of arterial arches the lung-arteries develop, in the Dipneusta.

IX. *Ninth Period: Vascular System of Amphibia.*

The gill-arches gradually disappear with the gills. The right and left aortal arches remain.

X. *Tenth Period: Vascular System of Mammals.*

The separation of the greater from the lesser circulation is complete. The right aortal arch unites with Botalli's duct.

TABLE XLI.

SYSTEMATIC SURVEY OF THE MOST IMPORTANT PERIODS IN THE PHYLOGENY OF THE HUMAN HEART.

I. *First Period: Heart of Chordonia.*

The heart forms a simple spindle-shaped enlargement of the ventral vessel, with an alternating blood-current (as in Ascidia).

II. *Second Period: Heart of Acrania.*

The heart is like that of *Chordonia*, but the blood-current acquires a constant direction, passing only from back to front. (Retrograded in *Amphioxus*.)

III. *Third Period: Heart of Cyclostoma.*

The heart divides into two chambers, a posterior auricle (*atrium*) and an anterior ventricle (*ventriculus*).

IV. *Fourth Period: Heart of Primitive Fishes.*

From the anterior section of the ventricle is differentiated an aortic stalk (*bulbus arteriosus*), as in all Selachii.

V. *Fifth Period: Heart of the Mud-fishes.*

The auricle divides, by an imperfect and interrupted partition, into a right and a left half, as in Dipneusta.

VI. *Sixth Period: Heart of Amphibia.*

The partition between the right and left auricles becomes complete in the higher Amphibia.

VII. *Seventh Period: Heart of Protamnia.*

The main chamber of the heart divides, by an incomplete partition, into a right and a left half, as in Reptiles.

VIII. *Eighth Period: Heart of Monotremata.*

The partition between the right and left ventricles becomes complete in all Mammals.

IX. *Ninth Period: Heart of Marsupials.*

The valves between the auricles and ventricles (atrio-ventricular), together with the connecting filaments and papillary muscles below them, are differentiated from the muscular masses of Monotremes.

X. *Tenth Period: Heart of Apes.*

The main axis of the heart, lying in the central line of the body, becomes oblique, so that the apex is turned to the left, as in Apes and Man.

TABLE XLII.

Comparative Survey of those Primitive Organs which must probably be regarded as homologous in Worms, Articulated Animals, Soft-bodied Animals, and Vertebrates.¹⁹³

Worms (Annelides).	Articulated Animals (Arthropoda).	Soft-bodied Animals (Mollusca).	Vertebrates (Vertebrata).
I. Products of the Differentiation of the Skin-sensory Layer.			
skin (dermis) upper throat- ganglia) sensory organs (nerve-vessels, sensory organs)	1. Chitinous skin (Hypodermis) 2. Brain (upper throat- ganglia) 3. Shell-glands of the Crustacean (trachea of the Tracheata?)	1. Outer skin (Epidermis) 2. Brain (upper throat- ganglia) 3. Rudimentary kid- neys (Primitive kidneys)	1. Outer skin (Epidermis) 2. Medullary tube (an- terior part) 3. Primitive kidney- ducts (Protoneph- roses) and seg- mental organs
II. Products of the Differentiation of the Skin-fibrous Layer.			
leather-skin (Corium) together with the muscular layer (together with the muscular layer of the skin?) 5. Side trunk-muscles	4. Leather-skin (Rudiment) 5. Trunk-muscles 6. Exocoelomic innermost cell-layer of the body-wall (also male germ-plate?)	4. Leather-skin (Corium) (together with the muscles of the skin?) 5. Inner trunk-muscles 6. Exocoelomic parietal epithelium of the coelom (also male germ-plate?)	4. Leather-skin (Corium) (together with the muscular layer of the skin?) 5. Side trunk-muscles 6. Exocoelomic parietal epithelium of the coelom (also male germ-plate?)
I. Products of the Differentiation of the Intestinal-fibrous Layer.			
body-cavity (Coeloma) endocoelomic outer- most cell-layer of the intestinal wall (together with the female germ-plate?) 9. Heart	7. Body-cavity (Coeloma) 8. Endocoelomic outer- most cell-layer of the intestinal wall (together with the female germ-plate?) 9. Heart	7. Body-cavity (Coeloma) 8. Endocoelomic visceral epithelium of the coelom (together with the female germ-plate?) 9. Chamber of the heart (and main artery)	7. Pleuro-peritoneal cavity 8. Endocoelomic visceral epithelium of the coelom (together with the female germ-plate?) 9. Aorta (primordial)
10. — 11. Intestinal wall (ex- cept the epithe- lium)	10. — 11. Intestinal wall (ex- cept the epithe- lium)	10. — 11. Intestinal wall (ex- cept the epithe- lium)	10. Heart (and gill- artery) 11. Intestinal wall (ex- cept the epithe- lium)
Products of the Differentiation of the Intestinal-glandular Layer.			
intestinal epithe- lium	12. Intestinal epithe- lium	12. Intestinal epithe- lium	12. Intestinal epithe- lium

CHAPTER XXV.

DEVELOPMENT OF THE URINARY AND SEX

Importance of Reproduction.—Growth.—Simplest Form of Reproduction: Division and the Formation of Buds (Gemmation).—Forms of Sexual Reproduction: Amalgamation of Male and Female Cells; the Male Sperm-cell and the Female Egg-cell.—Source of Love.—Original Hermaphroditism; Later Separation of Sexes (Gonochorism).—Original Development of Male and Female Sexual Cells from the Two Primary Germ-layers.—Development of the Male and Female Entoderm.—Development of the Testes.—Passage of the Sexual Cells into the Coelom.—Hermaphroditism of the Embryonic Epithelium, or Sexual Plate.—Classification of Sexual Ducts.—Egg-duct and Seed-duct.—Development of the Primitive Kidney Ducts.—Excretory Organs of the "Side Canals" of Ringed Worms (*Annelida*).—Side Canals of the Primitive Kidneys of the *Mysinioides*.—Primitive Kidneys of Animals (*Craniota*).—Development of the Paired Kidneys in *Amniota*.—Development of the Urinary Bladder.—Allantois.—Differentiation of the Primary and Secondary Kidneys.

the phenomena observed, and thus finds principles from which many experiences can be derived. This is more than mere experience; it is, — to speak, philosophical experience."—JOHANNES MÜLLER (1840).

we judge of the importance of the organ-systems of the animal body according to the number and variety of phenomena which they present, and according to the physiological interest connected with them, we must recognize as one of the most important and interesting organic systems, the one to the development of which we now, finally, turn; the system of the reproductive organs. Just as nutrition is the first and most important condition of self-preservation of the organic individual, so by reproduction alone is the preservation of the kind or species effected, or, rather, the preservation of the long series of generations, which in their genealogical connection form the sum of the organic tribe, or phylum. No organic individual enjoys an eternal life. To each is granted but a short span of time for his individual evolution, a brief, fleeting moment in the long millions of years of the earth's organic history.

Reproduction in connection with Heredity has, therefore, long been regarded as, after nutrition, the most important fundamental function of the organism, and it is customary to make this a primary distinction between living bodies and lifeless or inorganic bodies. But this distinction is in reality not so deep and thorough as it at first appears, and as is generally assumed. For, if the nature of the phenomena of reproduction is closely considered, it is soon seen that it may be reduced to a more general quality, that of growth, which belongs to inorganic, as well as to organic bodies. Reproduction is a nutrition

and a growth of the organism beyond the individual which, therefore, raises a part of the organism to the level of a whole (vol. i. p. 159). This is most clearly seen in observing the reproduction of the simplest and lowest organisms, especially of the Monera (p. 46) and of the unicellular Amœba (p. 48). In these, the simple individual possesses only the form-value of a single plastid. As soon as by continued nutrition and simple growth, this has reached a certain size, it does not exceed that size, but falls by simple division, into two similar halves. Each of the two halves thenceforth leads an independent life, and grows, till, having reached the same limit of growth, it again divides. At each of these simple self-divisions new central points of attraction for the particles of the body are formed, as foundations of the two new individuals.¹⁹⁴

In many other Primitive Animals (*Protozoa*), the reproduction is accomplished, not by division, but by the formation of buds (gemination). In this case, the growth which prepares the way for reproduction, is not total (as in the case of division), but partial. Hence in the case of gemination, the product of local growth, which, as soon as it forms a new individual, can be distinguished, as a new individual, from the parent-organism from which it originates. The latter is older and larger than the former. In the case of division, on the contrary, the two products are of equal age and of equal form-value. From the differentiated forms of asexual reproduction, connected with gemination, are, thirdly, the formation of germ-cells, and, fourthly, the formation of germ-cells. The latter, however, brings us directly to sexual reproduction, for

The opposed differentiation of the two sexes is the condition. In my *Generelle Morphologie* (vol. ii. pp. 32-71), and in my "Natural History of Creation" (vol. i. p. 183), I have fully discussed the connection of these various forms of reproduction.

None of the earliest ancestors of Man and of the higher animals were capable of the higher function of sexual reproduction, but multiplied only in an asexual manner, by division or gemmation, by the formation of germ-buds, or of germ-cells, as is still the case with most Primæval Animals or Protozoa. It was not until a later period in the organic history of the earth, that sexual difference of the two sexes could arise; and this took place at first in the simplest manner by the severance of two cells which amalgamated from the community of the many-celled organism. We may say that, in this case, growth, which is the condition necessary to reproduction, was attained by the union of two full-grown cells into a single cell which then exceeded its proper size ("copulation" or conjugation"). At first, the two united cells may have been entirely alike. Soon, however, by natural selection, a contrast must have arisen between them. For it must have been very advantageous to the newly-created individual in the struggle for existence, to have inherited various qualities from the two parent-cells. The complete development of this progressive contrast between the two producing cells, led to sexual differentiation. One cell became a female egg-cell, the other, a male seed or sperm cell.

The simplest form of sexual reproduction among existing animals, is exhibited in Gastræads and the lower Sponges, especially the Chalk Sponges, and, also, in the simplest

Hydroid Polyps. In the *Haliphysema* (Fig. 315) and the *Olynthus* the whole body is a simple intestinal pouch which is only essentially distinguished from the stomach by the fact that it is adherent by the end opposite the mouth. The thin wall of the pouch consists only of the two primary germ-layers. As soon as it is sexually mature, the single cells of the wall become female egg-cells, or become male sperm-cells, or seed-cells; the former are very large, as they form a considerable number of yolk granules in their protoplasm (Fig. 181, e); the latter, on the contrary, by continued division, become very small, and modify into movable "pin-shaped" spermatozoa (Fig. 182, f, vol. i. p. 173). Both kinds of cells sever themselves from their birthplace, the primary germ-layers, fall either into the surrounding water or into the intestinal cavity, and unite by amalgamation. This is the very important process of the fertilization of the egg-cell by the sperm-cell (Fig. 18, vol. i. p. 175.)

These simplest processes of sexual reproduction are exhibited at the present time in the lowest Plant Animals, especially in the Chalk Sponges and Hydroid Polyps, and among us of several extremely important and significant processes. In the first place, we learn, that for sexual reproduction in its simplest form, nothing more is required than the blending or amalgamation of two differing cells, a female egg-cell and a male sperm-cell, or seed-cell. All the circumstances, and all the other extremely complex phenomena, accompanying the act of sexual reproduction in higher animals, are of a subordinate and secondary character, and have only attached themselves secondarily to the simplest primary process of copulation or fertilization.

arisen by differentiation. But, now, if we consider that an extraordinarily important part is everywhere played by the relation of the two sexes in organic nature, in the vegetable kingdom, as in animal and human life; how the reciprocal repulsion and attraction of the sexes, which gives the impetus of the most varied and remarkable processes, is, in fact, one of the most important mechanical causes of the highest differentiation in life;—if we consider this, we cannot over-estimate this meaning of “love” to its primitive origin, to the power of attraction between two differing cells. Everywhere throughout animated nature



g. 315.—Longitudinal section through a hysema (*Gastrotricha*). The egg-cells (*e*) are between the epithelial cells of the entoderm (*g*), and lie freely in the primitive intestinal cavity. *m*, mouth-opening; *h*, exoderm.

greatest results proceed from this most insignificant cause. It is only necessary to think of the part played in nature by the flowers, the reproductive organ of flowering plants; or of the multitude of wonderful phenomena produced by sexual selection in animal life; or, finally, of the important influence exerted by love on human life: the coalescence of two cells is everywhere the single, original, all-pervading motive; everywhere this apparently trivial pro-

cess exerts the greatest influence on the development in the most varied circumstances. We may, indeed, assert that no other organic process can be, even remotely, compared to this in extent and intensity of differentiating effect. It is not the Semitic myth of Eve, who seduced Adam into knowledge, and is not the old Greek legend of Paris and Helen, and are not very many other famous fictions, but the poetical expression of the immeasurable influence of love, in connection with "sexual selection,"⁸⁶ has exercised ever since the differentiation of the two sexes, on the progress of the world's history? All other passions that issue from the human breast are in their combined effects far less powerful than love, which inflames the senses and fogs the understanding. On the one hand, we gratefully glorify in it as the source of the most splendid creations of art; the noblest productions of poetry, of plastic art and of music; we reverence in it the most powerful factor in the development of civilization, the basis of family life, and, consequently, the development of the state. On the other hand, we see in it the devouring flame which drives the unfortunate to ruin, and which has caused more misery, vice, and crime than all the other evils of the human race taken together. So wonderful is love, and so immeasurably important its influence on mental life, on the most varied functions of the medullary tube, that in this point, more than in any other, "supernatural" causation seems to mock every scientific explanation. And yet, notwithstanding all this, the comparative history of evolution leads us back very clearly and indubitably to the oldest and simplest source of the elective affinity of two differing cells: the sperm-cell and the egg-cell.

Just as the lowest Plant Animals exhibit this most simple origin of the complex phenomena of reproduction, in the second place, they reveal the highly important fact, that the earliest and most primitive sexual relation is hermaphroditism, and that the separation of the sexes originated from this only secondarily (by division of labour). Hermaphroditism is prevalent in lower animals of the most different groups; in these, each single individual, when sexually mature, each person, contains male and female sexual cells, and is, therefore, capable of self-fertilization and self-reproduction. Thus, not only in the lowest Plant Animals just mentioned (the *Gastræads*, Chalk-sponges, and many Hydroid Polyps) do we find egg-cells and sperm-cells united in one and the same person; but many Worms (for example, the *Ascidians*, Earth Worms and Leeches), many Snails (the common garden Snail), and many other invertebrate animals are also hermaphrodite. All the earlier invertebrate ancestors of man, from the *Gastræada* up to the *Chordonia*, must also have been hermaphrodite. So, probably, were also the earliest Skulled Animals (Figs. 52-56, *e, h*, vol. i. p. 256). One extremely weighty piece of evidence of this is afforded by the remarkable fact, that even in Vertebrates, in Man as well as other Vertebrates, the original rudiment of the sexual organs is hermaphrodite. The separation of the sexes (*Gonochoresis*), the assignment of the two kinds of sexual cells to different individuals, originated from hermaphroditism only in the farther course of tribal history. At first, male and female individuals differed only in the possession of the two kinds of cells, but in other respects were exactly alike, as is now the case in the *Amphioxus* and the *Cyclostoma*.

Not until a later period, by the law of sexual selection brilliantly elucidated by Darwin, were developed the so-called "secondary sexual characters," that is, those differences in the male and female sexes which are exhibited not in the sexual organs themselves, but in other parts of the body (for example, the beard of the man, the breasts of the woman).³⁶

The third important fact, taught us by the lower Animals, refers to the earliest origin of the two kinds of sexual cells. For, as in *Gastrea*, and in many *Sponges* and *Hydroids*, in which we meet with the simplest rudiments of sexual differentiation, the whole body consists throughout life only of the two primary germ-layers, the two kinds of sexual cells can, therefore, only have originated from one of the two primary germ-layers. This simple discovery is of extreme importance, because the question of the origin of the egg-cells as well as of the sperm-cells in the higher animals—and especially in *Vertebrates*—presents unusual difficulties. In these animals it usually appears as if the sexual cells developed, not from one of the two primary, but from one of the four secondary germ-layers. If, as most authors assume, they do originate from the middle-layer, or *mesoderm*, the fact is due to an ontogenetic heterotopism, to a displacement in position. (Cf. vol. i, p. 100.) Unless the unjustifiable and paradoxical assumption that the sexual cells are of entirely different origin in the higher and in the lower animals, is accepted, we are compelled to derive them originally (phylogenetically), in the form of the latter, from one of the two primary germ-layers. It can then be assumed that these cells of the skin-layer developed from the intestinal layer, which must be regarded as the

anitors of the sperm-cells and of the egg-cells, with-
out, during the separation of the skin-fibrous layer from
the skin-sensory layer, or of the intestinal-fibrous layer
from the intestinal-glandular layer, into the body-cavity
(*coeloma*), which was in process of formation; and that
they thus acquired the internal position between the two
germ-layers, which appears as their original position,
even when the sexual cells first become distinct in the vertebrate
embryo. Otherwise, we should be obliged to accept the
probable polyphyletic hypothesis, that the origin of the
egg-cells and sperm-cells is different in the higher and in
the lower animals, that their origin in the former is inde-
pendent of that in the latter.

If we, accordingly, derive the two kinds of sexual cells
from the two primary germ-layers in man as in all other
animals, the farther question arises: Did the female egg-
cells and the male sperm-cells develop from both primary
germ-layers, or from one only? and, in the latter case, from
which of the two? This important and interesting question
is one of the most difficult and obscure problems in the
history of evolution, and, up to the present moment, no full
and clear solution has been attained. On the contrary,
the most opposite answers are given to it even yet by
naturalists of note. Among the various possible solutions
only two have been generally considered. It has been
supposed that both kinds of sexual cells originally de-
veloped from the same primary germ-layer, either from the
skin-layer or the intestinal layer; but almost as many and
as able observers have accepted the one as the origin as
the other. Quite recently the Belgian naturalist, Eduard
van Beneden, has asserted, on the contrary, that the egg-cells

originate from the intestinal layer, the sperm-cells in skin-layer.¹⁹⁶ In *Gastræads*, *Sponges*, and *Hydra* this appears really to be the case. The development of sexual differences, which is so rich in results, accordingly, have commenced even during the differentiation of the two primary germ-layers in the simplest and most primitive *Plant Animals*; the exoderm would be the male germ-layer, the entoderm, the female. If this discovery of Van Beneden is established and proves to be a universal law, Biology will gain a most pregnant advance; for not only would the contradictory empiric explanations be answered, but a new path would be opened for philosophic reflection on the most important of biogenetic processes.

If we now trace the Phylogeny of the sexual differentiation in our earliest Metazoic ancestors further, as it is indicated at the present time, in the Comparative Anatomy and Ontogeny of the lowest Worms and *Plant Animals*, we note, as the first advance, the accumulation of the sexual cells of both sexes into definite groups. While in *Sponges* and the lowest *Hydra-Polyps* single scattered cells separate the cell-layers of the two primary germ-layers, and isolated and free sexual cells, in the higher *Plant Animals* and Worms we find these same cells associated and collected into groups of aggregate cells, which are, in the most forward, called "sexual glands," or "germ-glands" (*gonads*). It is only now that we can speak of sexual organs in a morphological sense. The female germ-glands which, in their simplest form constitute a mass of numerous egg-cells, are the ovaries (*ovaria*, or *oophores*) (211, c, p. 198). The male germ-glands, which in their primitive form also consist merely of a mass of sperm-

the testes (*testiculi*, or *orchides*; Fig. 211, *h*). We find ovaries and testes in this earliest and simplest shape only in many Worms (*Annelida*) and Plant Animals, also in the lowest Vertebrates, in the Skull-less Animals (*Crania*). In the anatomy of the *Amphioxus* we found the ovaries of the female and the testes of the male consisting of twenty to thirty elliptic or roundly four-cornered simple cells, of small size, attached to the inside of the gill-cavity on each side of the intestine. (Cf. vol. i. p. 425.)

Only a single pair of germ-glands, lying far down in the interior of the body-cavity (Fig. 316, *g*), exist in all Skulled Animals (*Craniota*). The first traces of these appear in the coelom-epithelium. Probably, in this case also, the male germ-cells originate from the skin-layer, the female egg-cells, on the contrary, from the intestinal layer. The earliest traces are visible in the embryo at the point where the skin-fibrous layer and the intestinal-fibrous layer meet in the middle plate (mesentery-plate) (Fig. 318, *mp*, p. 408). At this very important point in the coelom-wall, where the endocoelar (or visceral coelom-epithelium) merges into the exocoelar (or parietal coelom-epithelium), in the embryo of Man and the other Skulled Animals a small aggregation of cells becomes visible, at a very early period, and this, according to Waldeyer,¹⁹⁶ we may call the "germ-epithelium," or (corresponding with the other plate-shaped rudiments of organs) the sexual plate (Fig. 316, *g*; Plate IV. Fig. 5, *k*). The cells of this germ-plate, or sexual plate (*lamella sexualis*) are essentially distinguished by their cylindrical form and by their chemical constitution from the other cells of the coelom; they are of quite different significance from the flat cells of the "serous coelom-epithelium" which line the

remainder of the body-cavity (*coeloma*). Of these are the true coelom-cells—those which invest the intestine and the mesentery (“*endocoelar*”) originate from




FIG. 316.—Transverse section through the pelvic region and limbs of an embryo Chick in the fourth day of incubation, enlarged 40 times: *h*, horn-plate; *w*, medullary tube; *n*, canal of the medullary tube; *u*, primitive kidneys; *x*, notochord; *e*, hind limbs; *b*, allantois in ventral wall; *t*, aorta; *v*, cardinal veins; *a*, intestine; *d*, glandular layer; *f*, intestinal-fibrous layer; *g*, germ-epithelium; *m*, muscles; *c*, body-cavity, or Coelom. (After Waldeyer.)

intestinal-fibrous layer (in Fig. 5, Plate IV., coloured red) those which line the inner surface of the external wall of the abdomen (“*exocoelar*”) are, on the contrary, the products of the skin-fibrous layer (coloured blue in Fig. 5, Plate IV.) but the sexual cells which make their appearance at this boundary line between the two forms of coelom-cell

ch insert themselves, to a certain extent, between the
ocœlar and the exocœlar, there forming the germ-
e, cannot be referred either to the intestinal-fibrous
r or to the skin-fibrous layer, but directly to the two
ary germ-layers; for there are important grounds for
posing that even the first rudiment of the sexual plate is,
ably, hermaphroditic, and that this "sexual epithelium"
ble, in Man and all other Vertebrates, between the exo-
ar and the endocœlar) represents a primæval and simple
naphrodite gland. (Cf. vol. i. p. 256, Figs. 52-56, *e, h.*)
e inner half of this, in contact with the intestinal-fibrous
er, which is derived from the intestinal-glandular layer,
uld be the rudiment of the ovary; its outer half, in
tact with the skin-fibrous layer, which originates from
intestinal-glandular layer, would be the rudiment of the
tes. This is, of course, only conjectural.

We ought, accordingly, to distinguish two different
ual plates or germ-epithelia; the female sexual plate, a
oduct of the intestinal layer, which gives rise to the
ary-epithelium—the mother cells of the ova ("ovary-
te"); and the male sexual plate, lying externally over the
mer, and which is a product of the skin-layer, from which
ginates the testes-epithelium—the mother cells of the
erm-threads ("testes-plate"); but even the first recog-
able rudiments of the two sexual plates appear, indeed,
intimately associated in the human embryo and in those
the higher Vertebrates, that hitherto they have been re-
ded as a single, undifferentiated, common rudiment of an
gan; and it is still possible that the two kinds of sexual
nds arise by secondary differentiation from a common
liment.

Though we must recognize the formation of the kinds of sexual cells, and in their union at fertilization the one essential act of sexual reproduction, yet, in the majority of animals, other organs exist which also part in the act of fertilization. The most important of these secondary sexual organs are the exit-ducts which serve to conduct the mature sexual cells out of the body, and, next to these, the copulative organs, which transfer the fertilizing sperm from the male person to the female in which the eggs are situated. These latter organs exist only in the higher animals of various tribes, and are more widely distributed than the exit-ducts. Even these, however, are only of secondary formation, and are wanting in many animals of the lower groups. In these, as in the higher, the mature sexual cells are simply ejected from the body. In some cases they pass out directly through the skin-covering (as in the Hydra and many of the Infusorians and Protozoa); in other cases, they enter the stomach and are ejected through the mouth-opening (in Gas Sponges, and other Hydroid Polypes and Coral Animals); in yet other cases, they enter the body-cavity and pass out through a special aperture in the ventral wall (*porus genitalis*). The latter is the case in many of the higher Invertebrates and even in a few lower Vertebrates (Cyclostomes and a few Fishes). These indicate the earliest condition of this matter as it was in our ancestors. On the other hand, in all higher, and most lower Vertebrates (and in most, higher Invertebrates) special tube-shaped ducts from the sexual cells, or sexual ducts (*gonoducts*), are present in both sexes. In the female these convey the egg-cells out from the ovaries, and hence they have

egg-ducts (*oviductus*, or *tubæ fallopiaë*). In the sex these tubes convey the sperm-cells from the testes, hence they are called sperm-ducts (*spermaductus*, or *deferentia*).

The original, genetic condition of these two outlets is exactly the same in Man as in all higher Vertebrates, but in most Invertebrates it is entirely different; for in the latter the sexual ducts develop directly from the sexual glands, or from the external skin, or from the intestinal canal, in Vertebrates an organ-system is employed for the conveyance of the sexual products; one which originally had a very different significance and function—the urinary system, or urinary organs. The original, primary function of these organs is simply to eliminate useless matter from the body in a liquid form. The liquid product of this function is called the urine, and is discharged either directly through the external skin, or through the last section of the intestine. The tube-shaped "urinary ducts" only secondarily absorb the sexual products also and convey them out; thus become "urogenital ducts" (*ductus urogenitales*). This remarkable secondary combination of the urinary and sexual organs into a common "urogenital apparatus," or "urogenital system," is highly characteristic of the higher Vertebrates. In the lowest of these it is, however, wanting, but on the other hand, it is found in the higher Ringed Animals (*Annelida*). To estimate this rightly, we must first look at the comparative economy of the urinary organs in the whole.

The kidney system or urinary system (*systema urogenitum*) is one of the earliest and most important organs in the differentiated animal body, as has already

been incidentally mentioned. (Cf. Chapter XVII.) It is found almost universally distributed, not only in the higher animal tribes, but even in the more primitive Worms. Among the latter it even occurs in the lowest and imperfect known Worms—the Flat Worms (*Plathelminthes*) (Fig. 184, *nc*, p. 80). Although these acelomate Worms have no body-cavity, no blood, no vascular system, they always have a kidney system. It consists of a pair of simple or of branched canals, lined by a layer of cells, which absorb useless juices from the tissues and discharge them through an external skin-opening (Fig. 184, *nm*). Not only the free-living Gliding Worms (*Turbellaria*), but also the parasitic Sucking Worms (*Trematoda*), and even the still more degraded Tape Worms, which, in consequence of their parasitic habit of life, have lost their intestine, are all provided with these “kidney canals” or nephridia. Usually these canals in the Worms are excretory organs, and in former times they used to be regarded as water-vessels. Phylogenetically they must be regarded as highly-developed pouch-like skin-glands resembling the sweat-glands of Mammals, and, like these, developed from the skin-sensory layer. (Cf. Fig. 210, *n*, p. 198, and Fig. 202.)


While in these lowest unsegmented Worms only a pair of kidney ducts is present, in the higher segmented Worms these ducts exist in greater numbers. In Ringed Worms (*Annelida*), in which the body is composed of a great number of segments, or metamera, a pair of primitive kidneys (hence known as segmental organs) exists in each separate segment. In this case, the canals are very simple tubes, which, on account of

iled or looped form, are called "coiled canals." To the primary, external aperture in the outer skin, originally one present, a secondary, internal aperture into the body-cavity (*coeloma*) is now added. This opening is provided with vibratory cilia, and is thus enabled to absorb the secretional juices from the body-cavity and to discharge them from the body. Now in these Worms also the sexual cells, which develop in the simplest form upon the inner surface of the abdominal wall, pass, when mature, into the coelom, are drawn into the internal, funnel-shaped ciliated openings of the kidney canals, and are carried out of the body with the urine. Thus the urine-forming "coiled canals," or "primitive kidneys," serve, in the female Ringed Worms, as "oviducts," and, in the male, as "sperm-ducts."

It would of course be most interesting to know the condition, on this point, of the *Amphioxus*, which, standing midway between Worms and Vertebrates, affords us so much valuable information. Unfortunately this animal, for the present, affords no solution of this matter. At present we know nothing certainly as to the relation between the urinary and the sexual organs of the *Amphioxus*. Some zoologists assert that this animal has no kidneys; others regard the two long "side canals" as atrophied primitive kidney ducts (Fig. 152, *S*, vol. i. p. 423); yet others consider certain glandular epidermis-swelling on the inner surface of the gill-cavity to be rudimentary kidneys. Most probably, a great reversion has affected the original primitive kidney canals in the *Amphioxus*, amounting perhaps to their entire phylogenetic loss.

Very interesting inferences may be drawn from the Vertebrates of the next stage—the *Monorhina*, or *Cyclos-*

toma. Although both orders of this class—the Myx as well as the Petromyzontes—possess developed, secreting kidneys, these organs do not in this case carry away the sexual cells. These cells pass direct the germ-glands into the coelom, and are discharged t a posterior aperture in the abdomen. The condition primitive kidneys in these is, however, very interesti



throws light on the complex structure of the higher Vert. In the first place, in the noides (*Bdellostoma*) we find tube, the primitive kidney (*protureter*, Fig. 317, *a*), o side. This opens internal the coelom through a ciliated shaped aperture (as in Worms); it opens externally an opening in the outer sk great number of small hor tubes ("segmental canals," or

FIG. 317.—*A*. Portion of kidney lostoma: *a*, primitive kidney duct (*protureter*); *b*, segmental canals, or urine canals (*tubuli uriniferi*); *c*, vesicles (*capsulae Malphigianae*).—*B*. Portion of the same, much enlarged: *c*, vesicle, with the *glomerulus*; *d*, afferent artery; *e*, retrograde artery. (After J Müller.)

tive urine tubes) open on its inner side. Each o terminates in a blind, vesicular capsule (*c*) enclo

lot of blood-vessels (*glomerulus*, an arterial net, Fig. 17, B, c). Afferent arterial branches (*vasa afferentia*) convey arterial blood into the coiled branches of the "*glomerulus*" (d), and efferent arterial branches (*vasa efferentia*) again carry it out of the *glomerulus* (e).

In Primitive Fishes (*Selachii*) also there is a longitudinal series of segmental canals, which open outwardly in the primitive kidney ducts. The segmental canals (a pair in each metamer on of the central part of the body) open, in this case, freely into the body-cavity, through a ciliated funnel (as in Ringed Worms, or Annelids). A part of this organ forms a compact primitive kidney, while the rest is employed in the formation of the sexual organs.

The primitive kidney in the embryo of Man and in that of all other Skulled Animals (*Craniota*) is first formed in the same simple shape which persists throughout life in Myxinoides, and partly in *Selachii*. We found this primitive organ in the human embryo at that early period just succeeding the separation in the skin-sensory layer, of the medullary tube from the horn-plate, and the differentiation, in the skin-fibrous layer, of the notochord, the primitive vertebral plate, and the skin-muscle plate. As the first rudiment of the primordial kidneys, a long thin, thread-like string of cells, which is soon hollowed out into a canal, appears in this case, on each side, immediately below the horn-plate; this extends in a straight line from front to back, and is plainly seen in the cross section of the embryo (Fig. 318) in its original position in the space between the horn-plate (*h*), the primitive vertebræ (*uw*), and the skin-muscle plate (*hpl*). The first origin of this primitive kidney duct is still a matter of dispute, some ontogenists

referring it to the horn-plate, others to the primitive vertebral plate, and yet others to the skin-muscle plate. Probably its earliest (phylogenetic) origin is to be found in the skin-sensory layer; but it very soon quits its superficial position, and passes inward, between the primitive vertebral plates and the side plates, and finally lies upon the inner surface of the body-cavity. (Cf. Figs. 66-69, *u*, vol. i. p. 277. and Figs. 95-98, p. 319; also Plate IV. Figs. 3-6, *u*.) While the primitive kidney duct is thus making its way inward, on its inner and under side appear a large number of small horizontal tubes (Fig. 319, *a*), exactly corresponding to the segmental canals of the Myxinoides (Fig. 317, *b*). Like the latter, these are, probably, originally protuberances of the primitive kidney ducts (Fig. 316, *u*). At the blind, inner end of each of the primitive urinary tubes an arterial glomerulus is formed, which grows into this blind end from within, forming a "vascular coil." The glomerulus to a certain extent expands the bladder-like blind end of the small urinary tubes. As the primitive urinary tubes



FIG. 318.—Transverse section through the embryo of a Chick, on the second day of incubation: *h*, horn-plate; *m*, medullary tube; *uv*, primitive kidney duct; *ch*, notochord; *uv*, primitive vertebral cord; *spl*, skin-fibrous layer; *df*, intestinal-fibrous layer; *mp*, mesentery-plate, or middle plate (point of attachment of the two fibrous layers); *sp*, body-cavity (*coeloma*); *ao*, primitive aorta; *dd*, intestinal-glandular layer. (After Kölliker.)

position, passes inward, between the primitive vertebral plates and the side plates, and finally lies upon the inner surface of the body-cavity. (Cf. Figs. 66-69, *u*, vol. i. p. 277. and Figs. 95-98, p. 319; also Plate IV. Figs. 3-6, *u*.) While the primitive kidney duct is thus making its way inward, on its inner and under side appear a large number of small horizontal tubes (Fig. 319, *a*), exactly corresponding to the segmental canals of the Myxinoides (Fig. 317, *b*). Like the latter, these are, probably, originally protuberances of the primitive kidney ducts (Fig. 316, *u*). At the blind, inner end of each of the primitive urinary tubes an arterial glomerulus is formed, which grows into this blind end from within, forming a "vascular coil." The glomerulus to a certain extent expands the bladder-like blind end of the small urinary tubes. As the primitive urinary tubes

which are, at first, very short, grow longer and broader, each of the two primitive kidneys assumes the form of a uni-pinnate leaf (Fig. 320). The urinary tubes (*u*) repre-




FIG. 319.—Rudimentary primitive kidney of embryonic Dog. The posterior portion of the body of the embryo is seen from the ventral side, covered by the intestinal layer of the yolk-sac, which has been torn away, and thrown back in front in order to show the primitive kidney ducts with the primitive kidney tubes (*a*): *b*, primitive vertebræ; *c*, dorsal medulla; *d*, passage into the pelvic intestinal cavity. (After Bischoff.)

FIG. 320.—Primitive kidney of a human embryo: *u*, the urine-tubes of the primitive kidney; *w*, Wolffian duct; *w'*, upper end of the latter (Morgagni's hydatid); *m*, Müllerian duct; *m'*, upper end of the latter (Fallopian hydatid); *g*, hermaphrodite gland. (After Kobelt.)

sent the tissue and the primitive kidney duct (*w*) the mid-rib. On the inner margin of the primitive kidney the rudiment of the hermaphrodite sexual gland already

appears as a body of considerable size. The posterior end of the primitive kidney duct opens into the lower extremity of the last section of the rectum, so that this organ becomes a cloaca. But this opening of the primitive kidney duct into the intestinal canal must be regarded, phylogenetically, as a secondary condition. Originally, as is indicated clearly in the Cyclostoma, they issued through the external abdominal skin, quite independently of the intestinal canal, thus proving their early phylogenetic origin from the horn-plaques as outer skin glands.

While in the Myxinoidea the primitive kidneys permanently retain this simple form, as they do partially in Primitive Fishes (*Selachii*), in all other Craniota it appears only temporally in the embryo, as the ontogenetic reproduction of the primordial phylogenetic condition. In these Skulled Animals the primitive kidney, by vigorous growth, increases in length, and by the increase in number and the coiling of the urinary tubes, very soon assumes the form of a large compact gland, of oblong, oval, or spindle-shaped form, which extends longitudinally through the greater part of the body-cavity (*coeloma*) of the embryo (Figs. 123, *m*, 124, *m*, vol. i. p. 370). In this case, it lies near the middle line, directly under the primitive vertebral column, and extends from the region of the heart to the cloaca. The right and left primitive kidneys lie parallel and close together, being separated only by the mesentery, that narrow, thin lamella which connects the central intestine with the lower surface of the primitive vertebral column. The excretory duct of each primitive kidney, the protureter, traverses the lower and outer side of the gland in a posterior direction, and opens into the cloaca, close to the root of the allantois; at

later period, it opens into the allantois itself (Fig. 136, *o*, *pl. i.* p. 381).

The primitive kidney (primordial kidney) in the embryo of Amniota was formerly called the "Wolffian body," also the "Okenian body." In all cases it acts for a time as a true kidney, draining and secreting the useless fluids of the embryonic body, and discharging them into the cloaca and then into the allantois. The "primitive urine" collects in the latter organ, and hence the allantois in the embryo of man and of the other Amniota acts as a real urinary bladder, or "primitive urinary sac;" yet it is in no way genetically connected with the primitive kidneys, but is rather, as we have already seen, a pouch-like protuberance of the anterior wall of the terminal intestine (Fig. 135, *u*, *vol. i.* p. 380). The allantois is, therefore, a product of the intestinal layer, while the primitive kidneys are a product of the skin-layer. Phylogenetically we must conceive that the allantois originated as a pouch-shaped protuberance of the cloacal wall resulting from the distension caused by the collection in the cloaca of the primitive urine secreted by the primordial kidneys. It is, originally, a blind sac belonging to the rectum (Plate V. Fig. 15, *hb*). The true urinary bladder of Vertebrates, evidently, first appeared in Dipneusta (in the *Lepidosiren*), and was thence transmitted, first to the Amphibia, and then to the Amniota. In the embryo of the latter it protrudes far out of the yet unclosed abdominal wall. Many Fishes, indeed, also possess a so-called urinary bladder. But this is merely a local distension in the lower section of the primitive kidney ducts, and hence, both in origin and in constitution, is essentially distinct from the true urinary bladder. The two structures

are only physiologically comparable; they are, then, analogous, as having the same function; morphologically, however, they are not to be compared, or are not homologous.¹⁸⁸ The false urinary bladder in Fishes is a duct of the primitive kidney duct, therefore of the inner layer; the true urinary bladder in Dipneusta, Amphibia, and Amniota is, on the contrary, a blind-sac (terminal intestine, and hence a product of the outer layer).

In all low Skulled Animals (*Craniota*), without exception (in *Cyclostoma*, Fishes, *Dipneusta*, and *Amphibia*) the urinary organs remain in an inferior stage of development in so far as the primitive kidneys (*protonephra*), much modified, here act permanently as urine-secreting glands. In the three higher vertebrate classes, including the term Amnion Animals, on the contrary, this is not only for a short period during early embryonic life permanent, or secondary kidneys (*renes*, or *metanephra*), which are peculiar to these three classes, are very highly developed. These originate, not (as was long believed on the authority of Remak) as entirely new, independent glands of the intestinal tube, but from the posterior end of the primitive kidney duct (*protonephros*). From the point near where it opens into the cloaca, a simple pouch-like secondary kidney duct—grows out, and this increases considerably in length forwards; from the blind, upper anterior portion of this the permanent kidney originates precisely as the primitive kidney originates from the primitive kidney duct. The secondary kidney duct gives rise to a number of small blind tubes—the secondary urinary tubes—and the blind capsule-shaped ends of

occupied by vascular coils (*glomeruli*). The further growth of these tubes results in the compact secondary kidney, which, in Man and most higher Mammals, acquires the well-known bean-like form; in the lower Mammalia, Birds and in Reptiles, on the other hand, it is separated into several lobes. The lower, or posterior part of the permanent kidney duct retains the form of a simple canal, widens, and thus forms the permanent urine duct (*ureter*). At first this canal, yet united with the last section of the primitive kidney duct, discharges into the cloaca; at a later period, it separates from the primitive kidney duct, and yet later from the rectum, and then it discharges into the permanent urinary bladder (*vesica urinaria*). The latter originates from the posterior, or lower part of the stalk of the allantois (*urachus*), which widens and becomes spindle-shaped before opening into the cloaca. The anterior, or upper part of the allantois-stalk, which passes in the abdominal wall of the embryo to the navel, afterwards disappears, a useless cord-shaped remnant alone remaining as a rudimentary organ: this is the single urinary-bladder navel-cord (*ligamentum vesico-umbilicale medium*). On the right and left of this, in the adult Man, there are two other rudimentary organs: the lateral urinary-bladder navel-cords (*ligamenta vesico-umbilicalia lateralia*). These are the obsolete cord-like remnant of the former navel-arteries (*arteriæ umbilicales*, vol. i. p. 400; Fig. 326, a).

Although in Man, as in all other Amnion Animals, the primitive kidneys are thus very early displaced by the secondary kidneys, and although the latter alone afterwards act as urinary organs, the former are not, however, altogether discarded. Indeed, the primitive kidney ducts acquire

a high physiological significance, as they modify the excretory ducts of the sexual glands. In all Amphibia and Gnathostomi—therefore in all Vertebrates from Fish to Man—at a very early period, a second similar duct appears in the embryo at the side of each primitive kidney. This canal is commonly called, after its discoverer, Johannes Müller, “Müller's duct” (*ductus Mülleri*). The earlier, primitive kidney duct is distinguished as the “Wolffian duct” (*ductus Wolffii*). The actual origin of Müller's duct is still undetermined; Comparative Anatomy and Ontogeny seem, however, to indicate that it probably arises by differentiation from the Wolffian duct. It is, probably, most correct to say, that the original (primary) primitive kidney duct breaks up by differentiation (or fission) into two secondary, similar ducts; these are the Wolffian




FIG. 321.—Primitive kidneys and rudiments of the sexual organs of Amphibia (Frog larvæ); A, earlier, B, later condition. C, of man (embryo of Ox): u, primitive kidneys; k, sexual glands (rudiments of testes and ovaries). The primary primitive kidney duct (ug) in A separates (in B and C) into the two secondary primitive kidney ducts, the Müllerian duct (m) and the Wolffian duct (ug'), which unite behind the genital cord (g); l, groin-cord of the primitive kidneys. (After Gegenbaur.)

Müllerian ducts. The latter (Fig. 320, *w*) lies immediately inside the former (Fig. 320 *m*). Both open posteriorly into the cloaca.

obscure and uncertain as to the origin of the Müllerian ducts, their later fate is clear and definite. In Double-nostrilled (*Amphioxus*) and Jaw-mouthed (*Stomatopoda*) animals, from the Wolffian duct becomes the oviduct, and the Müllerian duct becomes the oviduct. In each of these is per-



Fig. 322, 323.—Urinary and sexual systems of an Amphibian (Water-Newt, *Amphioxus*). Fig. 322 (A), female; Fig. 323 (B), male: *r*, primitive kidney; *ov*, ovary; *od*, egg-duct and oviduct, both formed from the Müllerian duct; *u*, primitive urinary duct, in man, also as seed-duct—opening below into Wolff's duct; *ms*, ovary-mesentery (mesenteries). (After Gegenbaur.)

the other entirely disappears, or leaves only a rudimentary organ. In the male sex, in the two Wolffian ducts become sperm-ducts, certain parts of the Müllerian duct are often found, which I call "Rathke's canals" (Fig. 323, *c*). In the female there, on the contrary, the two Müllerian ducts

become oviducts, traces of the Wolffian ducts remain, and are known as "Gartner's canals."



FIGS. 324-326.—Urinary and sexual organs of an embryonic *Ox*. 1. 324, of female embryo of $1\frac{1}{2}$ inch in length; Fig. 325, of male embryo of $2\frac{1}{2}$ inches in length; Fig. 326, of female embryo of $2\frac{1}{2}$ inches in length; *wg*, Wolff's duct; *m*, Müller's duct; *m'*, upper end of the latter (opened at *t*); *i*, lower thickened end of the same (rudiment of uterus); *g*, genital cord; *h*, testes (*h'*, lower, *h''*, upper testis-cord); *o*, ovary; *o'*, lower ovary-cord; *i*, groin-cord of the primitive kidney; *d*, diaphragm-cord of the primitive kidney; *n*, permanent kidneys (between these the S-shaped urine-duct; between the two the rectum); *v*, urinary bladder; *a*, navel-artery. (After Kölliker.)

The most interesting facts in reference to this remarkable development of the primitive kidney ducts and their union with the sexual glands are exhibited in Amphibia (Figs. 321-323). The first rudiment of the primitive kidney ducts and their differentiation into the Müllerian

ian ducts is identical in both sexes, as is the case in embryos of Mammals (Fig. 321, *C*, Fig. 324). In the Amphibia the Müllerian duct on each side develops a large ovary (Fig. 322, *od*), while the Wolffian duct acts permanently as a urinary duct (*u*). In the male, on the contrary, the Müllerian duct persists only as a rudimentary organ, without functional significance, as Rathke's canal (Fig. 323, *c*); the Wolffian duct serves, in this case also, as a urinary duct, but also as a sperm or seed duct, the seminal duct (*se*) from the testes (*t*) entering the upper part of the primitive kidneys, and there uniting with the urinary canals. In Mammals these conditions, persistent in Amphibia, are early traversed by the embryo in an early period of its development (Fig. 321, *C*). The primitive kidneys, which in non-amnionate Vertebrates persist throughout life as the excretory organ, are superseded by the secondary kidneys. The actual primitive kidneys disappear almost entirely in the embryo at an early period, leaving but small rudiments. In the male Mammal the supplementary testis (*epidymis*) develops from the upper part of the primitive kidney; in the female the same part gives rise to a useless rudimentary organ, the supplementary ovary (*parovarium*). In the female Mammal the Müllerian ducts undergo very considerable changes. The actual ovaries develop only from the upper part; the lower part widens out into a spindle-shaped pouch, with a thick, fleshy wall, within which the fertilized egg develops into the embryo. This pouch is the uterus (*uterus*). At first the two uteri are perfectly separate, and open on each side of the urine-bladder (*vu*) into the cloaca, as is yet permanently the case in the lowest living Mammals, the Beaked Animals (*Ornithostoma*);

II. 2 E

but even in Pouched Animals (*Marsupialia*) a form exists between the two Müllerian ducts, and in some Animals they coalesce below with the rudimentary




FIG. 327.—Female sexual organs of a Beaked Animal (*Ornithorhynchus*, Figs. 195, 196): *o*, ovaries; *t*, oviduct; *u*, uterus; *u.g.*, urinary sexual cavity (*sinus urogenitalis*); the two parts of the uterus open into this at *u'*: *cl*, cloaca. (After Gegenbaur.)

ducts, forming with them "sexual cord" (*funiculalis*). But the original tendency of the two parts of the uterus, and of the two canals which proceed out to the lower extremities, persists in the lower Placental Animals, and in the higher members of the group, these organs coalesce to form one single

The process of coalescence advances steadily from behind (from behind) upwards (from behind) upwards). While in many Animals (*Rodentia*, e.g., *Squirrels*) two separate uteri open into the vagina canal which is already become simple, in other Gnawing Animals (e.g., *Beasts of Prey*, *Whales*, and *Hoofed Animals*) the lower halves of the two uteri are already coalescent, and their upper halves (the so-called horns, "*cornua*") remain distinct ("*uterus bicornis*"). In *Bats* and *Semi-bats* the upper horns are very short, while the unified lower part becomes longer. Finally, in *Apes*, as in *Man*, the coalescence of the two parts is complete, one simple pear-shaped pouch alone remaining, and into this the oviducts open on each side.

In the male Mammal also, a similar coalescence of the portion of the Müllerian and Wolffian ducts takes

In this case also, these ducts form a single "sexual" (Fig. 325, *g*), which likewise opens into the original primary sexual cavity (*sinus urogenitalis*), which develops into the lower part of the urinary bladder (*v*). While, however, in the male Mammal the Wolffian ducts develop into the permanent sperm-ducts, only very slight traces of Müllerian ducts remain as rudimentary organs. The most remarkable of these is the "male uterus" (*uterus masculinus*), which originates from the lowest, coalescent portion of the Müllerian ducts, and which is homologous to the female uterus. It forms a small flask-shaped sac, entirely without physiological significance, which opens into the urinary tubes between the two sperm-ducts into the prostatic lobes (*vesicula prostatica*).

The internal sexual organs in Mammals undergo very peculiar modifications in point of position. At first the germ-glands, in both sexes, lie deep down in the ventral region, on the inner side of the primitive kidneys (Figs. 320, 321, *k*), attached to the vertebral column by a short suspensory artery (in the male, the *mesorchium*; in the female, the *paramesarium*). It is only, however, in Monotremes that this ventral position of the germ-glands is (as in lower Vertebrates) permanent. In all other Mammals (Marsupials as well as Placentals) these glands quit their place of origin and take their way more or less downward (or towards the anterior extremity), following the course of a cord which extends from the primitive kidney to the groin region of the abdominal wall. This is the groin-cord of the primitive type; in the male, the "Hunterian guiding-cord" (*gubern-*

naculum testis) (Fig. 328, *M*, *gh*); in the female, the round uterus-cord (Fig. 328, *F*, *r*). In the latter the ovaries migrate more or less in the direction of the small pelvis.


FIG. 328, *M*.FIG. 328, *F*.

FIG. 328.—Original position of the sexual glands in the abdominal cavity of the human embryo (of three months). Fig. 328, *M*, male (natural size); *h*, testis; *gh*, the conducting-cord of the testis; *wg*, seed-duct; *b*, urine bladder; *uh*, lower hollow vein (*vena cava*); *nn*, supplementary kidneys. Fig. 328 *F*, female (somewhat enlarged): *r*, round uterus (below this the urine-bladder, above it the ovary); *r'*, kidney; *s*, supplementary kidney; *c*, blind-intestine (*caecum*); *o*, small intestine; *om*, stomach (between the two is the stomach); *l*, spleen. (After Kölliker.)

even enter this. In the male the testis quits the abdominal cavity altogether, passing through the groin-canal, enters a sac-shaped, distended fold of the external skin covering. The coalescence of the right and left folds ("sexual folds") gives rise to the testis-sac (*scrotum*). Various Mammals exhibit the various stages of this migration. In the Elephant and in Whales the testes descend very little, and lie below the kidneys. In many Gnawing Animals (*Rodentia*) and Beasts of Prey (*Carnaria*) they enter the groin-canal. In most higher Mammals they descend down through this into the testis-sac; usually the wall

The groin-canal coalesce. When, however, this remains open, the testes are able to descend periodically (in the rutting season) into the testis-sac, returning again into the abdominal cavity (e.g., in Pouched Animals or *Marsupialia*, Gnawing Animals, Bats, etc.).

Another peculiarity of Mammals is the formation of the external sexual organs which, as copulative organs, serve to carry the fertilizing sperm from the male into the female organism in the act of copulation. Organs of this sort are altogether wanting in most lower Vertebrates. In those which are aquatic (e.g., *Acrania*, *Cyclostoma*, and most Fishes) the eggs and sperm are simply discharged into the water, and their coming together is the result of some lucky accident which in this way brings about impregnation. On the other hand, in many Fishes and *Amphibia* which bring forth their young alive, there is a direct transfer of the sperm from the male to the female organism; and this is the case in all *Amniota* (Reptiles, Birds, and Mammals). In these animals the urinary and genital organs always open originally into the lower part of the rectum, which thus forms a "cloaca" (p. 345); but among Mammals the cloaca is permanent only in the Beaked Animals (*Ornithostoma*), which have, on this account, been called Cloacal Animals (*Monotrema*, Fig. 327, *cl*). In all other Mammals a lateral partition wall develops in the cloaca (in the human embryo about the middle of the third month), by which the latter is separated into two cavities. The urinary sexual canal passes into the anterior cavity (*sinus urogenitalis*), and it is through this cavity alone that the urinary and sexual products are discharged, while the "anal cavity," which lies behind it, serves merely to eject the excrement through the

anus. Even before the appearance of this partition, in Pouched Animals (*Marsupialia*) and Placental Animals, a conical papilla—the sexual protuberance (*phallus*, Fig A, e, B, e)—rises on the anterior part of the circumference




FIG. 329.—External sexual organs of the human embryo: A, male germ (in the eighth week; twice the natural size; with cloaca); B, male germ (in the ninth week; twice the natural size; anus distinct from urogenital opening); C, female germ in the eleventh week; D, female germ in the fourteenth week; e, sexual protuberance (*phallus*); f, sexual folds; hl, sexual folds; r, Raphe (point of union of the penis and clitoris); a, anus; ug, urinary sexual opening; n, navel-cord; s, tail. (After Cf. Table XLIV., p. 431.)

the cloaca-opening. The apex of this is swollen into a bulb (the "acorn," *glans*). On the under side appears a deep fold (the *sulcus genitalis*, f), and on each side of the latter fold, or sexual fold (hl). The phallus is especially thick at the base of the "sexual sense," and over it are distributed the

s (*nervi pudendi*) which are especially concerned in inducing the sexual sensations (p. 238). In the male the penis develops into the masculine "*penis*" (Fig. 329, *D, e*); in the female it becomes the much smaller "*clitoris*" (Fig. 330, *C, e*); only in some Apes (*Ateles*) does this become unusually large. The "fore-skin" (*præputium*), in both sexes, also develops as a skin-fold from the anterior part of the circumference of the phallus. In the male sexual furrow the lower side of the phallus receives the urogenital opening, and, as a continuation of the latter, modifies, by the invagination of its two parallel edges, into a closed canal—the male urinary tube (*urethra*). In the female this occurs in a few instances (in some Semi-apes, Gnawing Animals, *Proboscidea*, and Moles); as a rule the sexual furrow remains open, and its edges are developed into the *labia minora*. The *labia majora* of the female develops from the two lateral skin-folds which appear on each side of the sexual furrow. In the male these last folds coalesce, forming a closed sac, the testis-sac (*scrotum*). Occasionally this invagination does not take place, and the sexual furrow also sometimes remains open (*hypospadiæ*). In these cases the external male genitalia resemble the female, and this phenomenon has often been mistaken for hermaphroditism (pseudo-hermaphroditism).¹⁹⁷

From this and other cases of false "hermaphroditism," much less frequent cases of "true hermaphroditism" are distinguished. This exists only when the essential organs of reproduction, both kinds of germ-glands, are united in one individual. Either an ovary is then developed on the right, and a testis on the left (or *vice versa*); or testes and ovaries are developed on both sides, one more, the other less

perfectly. As we have already seen that the original rudiment of the sexual organs is really hermaphroditic in all Vertebrates, and that the separation of the sexes is due to a one-sided development of this hermaphroditic rudiment, these remarkable cases offer no theoretical difficulties. They very seldom, however, occur in Man and higher Vertebrates. On the other hand, we find original hermaphroditism constant in some lower Vertebrates, in some Fishes of the Perch kind (*Serranus*), and in Amphibia (*Bombinator* and in Toads). In these cases the male has usually a rudimentary ovary at the upper extremity of the testis; on the other hand, the female sometimes a rudimentary testis, without function. It also occurs occasionally in Carp and some other Fish. We have already seen how the original hermaphroditism is maintained in the excretory ducts, in Amphibia.

In the germ-history of the human urinary and sexual organs, the outlines of the history of human descent have been faithfully maintained up to the present time. We trace their development in the human embryo step by step in the same gradations as are exhibited, one after another, in the comparison of the urogenitals in Acrania, Cyclostomes, Fishes, Amphibians, and then further, in the series of Mammals, in Cloacal Animals (*Monotremes*), Marsupial Animals (*Marsupialia*), and the various Placental Animals (Cf. Table XLIII.) All the structural peculiarities of the urogenitals, distinguishing Mammals from other Vertebrates, are also present in Man; and in all special characteristics the latter resembles the Apes, and especially the Anthropoid Apes. As evidence that the special peculiarities of Mammals have been transmitted to Man, I will first

riefly notice the similar manner in which the eggs are formed in the ovary. In all Mammals, the mature eggs are contained in peculiar vesicles, which, after their discoverer, Regner De Graaf (1677), are called the Graafian follicles. These were formerly regarded as the actual eggs, which were, however, discovered by Baer *within* the Graafian follicles (vol. i. p. 55). Each follicle (Fig. 330, *C*) consists of a round, fibrous capsule, which contains fluid and is coated by several layers of cells. At one point this cellular layer has a knob-like enlargement (*C, b*), and, there, surrounds the real egg (*C, a*). The mammalian ovary is, originally, a very simple oblong little body (Fig. 320, *g*), formed only of connective tissue and blood-vessels, and surrounded by a cell-layer (the epithelium of the ovary, or the female germ-epithelium). From this epithelium, cords of cells grow inward, into the connective tissue or "stroma" of the ovary (Fig. 330, *A, b*). Single cells of these cords increase in size and become egg-cells (primitive eggs, *A, c*); but the greater number of the cells remain small and form an enveloping and nutritive cellular layer (the follicle-epithelium) round each egg.

In Mammals the follicle-epithelium is at first one-layered (Fig. 330, *B, 1*), afterwards many-layered (*B, 2*). In all other Vertebrates, the egg-cell is, indeed, enclosed in a permanent covering of small cells, an egg-follicle; but only in Mammals does fluid accumulate between the growing follicle-cells, and thus extends the follicle into a round bladder of considerable size, on the inner wall of which the egg lies excentrically. In this point, as in his whole Morphology, Man unmistakably indicates his descent from Mammals.



FIG. 330, A.

FIG. 330, B.




FIG. 330, C.

FIG. 330.—Development of human ovules within the female ovary.—A. Vertical section through the ovary of a new-born female: *a*, epithelium of ovary; *b*, rudiment of an egg-cord; *c*, young eggs in the epithelium; *d*, longer egg-cord with the follicles; *e*, group of young follicles; *f*, single young follicle; *g*, blood-vessels in the connective tissue (*stroma*) of the ovary. In the cords the young primitive eggs can be distinguished from surrounding cells of the follicle by their relatively large size. (After Waldeyer).—330, B. Two young follicles isolated; in 1, the cells of the follicle form but a single layer around the young primitive egg; in 2, they form a double layer; in 3, they begin to form the primary chorion (*a*), or the zona pellucida (vol. i. p. 135).—330, C. A mature human Graafian follicle: *a*, the mature egg; *b*, the surrounding follicle-cells; *c*, the epithelial cells of the follicle; *d*, the fibrous membrane of the follicle; *e*, its outer surface.

The entire natural history of the human sexual organs is one of the branches of Anthropology which affords the strongest proofs of the origin of the human race from the animal kingdom. Each man, on knowing the pertinent facts, and without prejudice, judging these comparatively, can but be convinced that he is descended from lower Vertebrates. The general, and the more minute structure, the activity and the individual evolution of the sexual organs, is exactly the same in Man as in Apes. This is as true of the male as of the female, of the internal as of the external genitalia. The differences in this matter between Man and the most man-like Apes are far less than the differences between the various forms of Apes. As, however, all Apes are undoubtedly from a common origin, this fact alone proves, with absolute certainty, the descent of Man from Apes.

TABLE XLIII.

SYSTEMATIC SURVEY OF THE MOST IMPORTANT PERIODS IN THE PHYLUM OF THE URINARY AND SEXUAL ORGANS OF MAN.¹²

XLIII. A. First main division: the sexual organs (G) and the urinary organs (U) are distinct. (The sexual or genital system (G) and the excretory or urinary system act independently of each other.)

I. First Period: *Genitals and Kidneys of Gastrozoa.*

G. Single, scattered cells of the entoderm change into egg-cells; single scattered cells of the exoderm into sperm-cells.

U. Special urinary organs are as yet wholly wanting. Secretion performed by the cells of the exoderm.

II. Second Period: *Genitals and Kidneys of Primitive Worms.*

G. The egg-cells of the entoderm gather into groups (ovary-plates); do the sperm-cells of the exoderm (testis-plates).

U. A pair of simple pouch-like skin-glands (products of the skin-sensory layer) develop into extremely simple kidney-canals (excretory organs of Flat-worms, *Platelmintes*).

III. Third Period: *Genitals and Kidneys of Scolecida.*

G. After the differentiation of the four secondary germ-layers is complete, the egg-cells pass from the skin-sensory layer into the skin-fibrous layer; the sperm-cells also pass from the intestinal-glandular layer into the intestinal-fibrous layer.

U. After the formation of the coelom is completed, the blind inner ends of the two kidney-canals (or "primitive kidney ducts") open into the body cavity (*coeloma*).

IV. Fourth Period: *Genitals and Kidneys of Chordonia.*

G. The groups of egg-cells (ovarial plates) and the groups of sperm-cells (testes-plates) meet at the boundary between the endocoele (the visceral intestinal-fibrous layer of the coelom-epithelium) and the exocoele (the parietal skin-fibrous layer of the coelom-epithelium), so as to form hermaphrodite glands.

U. The primitive kidney ducts differentiate into an excretory and glandular part.

V. Fifth Period: *Genitals and Kidneys of Acrania.*

G. The sexes become distinct. In the female, only the ovary is developed; in the male, only the testes.

U. The primitive kidney ducts remain simple (atrophied in *Amphioxus*).

VI. *Sixth Period : Genitals and Kidneys of Cyclostoma.*

G. The sexual glands (numerous in *Acrania*) coalesce into a pair.

U. The primitive kidney ducts send out lateral branches which acquire scalar coils (*glomeruli*) (the semi-pinnate primitive kidneys of *Bdellostoma*).

LIII. B. Second main division: the genital organs (G) and the urinary organs (U) become united. (The sexual system and the urinary system are united in the "urogenital system.")

VII. *Seventh Period : Urogenitals of Primitive Fishes (Selachii).*

The primary primitive kidney duct differentiates on each side, forming two secondary canals; the Wolffian duct, which develops into the seed-duct, and the Müllerian duct, which develops into the oviduct. Both genital ducts originally open behind the anus (*Proselachii*).

VIII. *Eighth Period : Urogenitals of Dipneusta.*

A cloaca is formed by the union of the urogenital opening and the cavity of the anus. The single urinary bladder grows out from the anterior wall of the rectum (*Lepidosiren*).

IX. *Ninth Period : Urogenitals of Amphibia.*

From the uppermost part of the primitive kidney which is in process of atrophy, proceeds, in the male sex, the supplementary testis; in the female sex, the supplementary ovary. The Wolffian duct yet acts, in both sexes, as urinary canal, and, in the male, also as the seed-duct. The Müllerian duct acts in the female sex as oviduct; in the male it is a rudimentary organ (Rathke's duct).

X. *Tenth Period : Urogenitals of Protamnia.*

The atrophied primitive kidney is replaced by the permanent secondary kidney as the urinary organ. The urinary bladder grows out from the ventral orifice of the embryo and forms the allantois. From the anterior wall of the cloaca grows the sexual protuberance (*phallus*), which, in the male, develops to the penis, in the female, to the clitoris.

XI. *Eleventh Period : Urogenitals of Monotremes.*

The lower end of the oviduct enlarges on each side to a muscular uterus.

XII. *Twelfth Period : Urogenitals of Marsupialia.*

The cloaca is separated by a partition into an anterior urogenital opening (*pertura urogenitalis*) and a posterior anal opening (*anus*). From the

lower part of the uterus the vagina-canal passes out on each side. Ovaries and testes begin to move downward from their place of formation.

XIII. Thirteenth Period: Urogenitals of Semi-apes.

The lower parts of the Müllerian and the Wolffian ducts coalesce to form a sexual cord. The coalescence of the two uteri at the lower part gives rise to the *uterus bicornis*. A part of the allantois becomes the placenta.

XIV. Fourteenth Period: Urogenitals of Apes.

The two uteri coalesce throughout their entire length, forming a pear-shaped uterus, as in Man.

TABLE XLIV.

ic Survey of the Homologies of the Sexual Organs in the two Sexes
of Mammals.

CLIV. A. HOMOLOGIES OF THE INTERNAL SEXUAL ORGANS.

<i>non Rudiments of the nal Sexual Organs.</i>	<i>M. Internal Male Parts.</i>	<i>F. Internal Female Parts.</i>
rm-gland (testes-plate embryo, product of the yer?)	1. Testis (<i>Testis</i> , or <i>Orchis</i>)	1. Rudimentary testis dis- appears,—remains in some Amphibia
erm-gland (ovary-plate, t of the intestinal)	2. (Rudimentary ovary, dis- appears,—remains in some Amphibia)	2. Ovary (<i>Ovarium</i> , or <i>Oophoron</i>)
duct (lateral primitive duct)	3. Seed-duct (<i>Spermaductus</i>)	3. Gartnerian duct (rudi- mentary canal)
an duct	4 a. Rathke's duct (rudi- mentary canal in Amphibia)	4 a. Oviduct (<i>Oviductus</i> , or <i>Tuba Fallopia</i>)
s <i>Mülleri</i> , central pri- kidney duct)	4 b. <i>Hydatis Morgagni</i>	4 b. <i>Hydatis Fallopia</i>
part of the Müllerian	4 c. <i>Uterus masculinus</i> (<i>Vesicula prostatica</i>)	4 c. Uterus, sheath (<i>vagina</i>)
of the primitive kidney	5. Supplementary testes (<i>Epididymis</i>)	5. Supplementary ovary (<i>Parovarium</i>)
nephron, <i>corpus Wolf-</i>	6. Hunterian guiding-cord (<i>Gubernaculum Hun-</i> <i>teri</i>)	6. Round uterus-cord (<i>Ligamentum uteri rotundum</i>)
ament of the primitive	7. Testis-mesentery (<i>Mesorchium</i>)	7. Ovary-mesentery (<i>Mesocarium</i>)
<i>mentum protonephroin-</i> (e)		
mesentery		
<i>terium sexuale</i>)		

CLIV. B. HOMOLOGIES OF THE EXTERNAL SEXUAL ORGANS.

<i>non Rudiments of the al Sexual Organs.</i>	<i>M. External Male Parts.</i>	<i>F. External Female Parts.</i>
protuberance (<i>us</i>)	8. Penis	8. Clitoris
in <i>utium</i>)	9. Male fore-skin (<i>Praputium penis</i>)	9. Female fore-skin (<i>Praputium clitoridis</i>)
fold (<i>genitales</i>)	10. Testes-sac (<i>Scrotum</i>)	10. <i>Labia pudendi majores</i>
between the two sexual	11. Seam of the testis-sac (<i>Raphe scroti</i>)	11. Female Vulva
edges (edges of the furrow)	12. Edges of the sexual furrow coalesce	12. <i>Labia pudendi minora</i>
tal canal (<i>urepentalis</i>)	13. Ureters (<i>Urethra</i>)	13. Antechamber of the vagina (<i>Vestibulum vagina</i>)
ar appendages of the tal canal	14. Cowper's glands	14. Bartholi's glands

CHAPTER XXVI.

RESULTS OF ANTHROPOGENY.

Review of the Germ-history as given.—Its Explanation by the Fundamental Law of Biogeny.—Its Causal Relation to the History of the Rudimentary Organs of Man.—Dysteleology, or the Doctrine of Purposelessness.—Inheritances from Apes.—Man's Place in the System of the Animal Kingdom.—Man as a Vertebrate and a Mammal.—Special Tribal Relation of Men and Apes.—Evidences regarding the Ape Question.—The Catarrhina and the Platyrrhina.—The Divine Origin of Man.—Adam and Eve.—History of the Evolution of the Human Mind.—Important Mental Differences within a Single Class of Animals.—Mammalian Mind and the Insect Mind.—Mind in the Ant and the Scale-louse (*Coccus*).—Mind in Man and in Ape.—The Organ of Activity: the Central Nervous System.—The Ontogeny and Phylogeny of the Mind.—The Monistic and Dualistic Theories of the Mind.—Heredity of the Mind.—Bearing of the Fundamental Law of Biogeny on Psychology.—Influence of Anthropogeny on the History of the Monistic Philosophy and the Defeat of the Dualistic.—Natural Science and Spiritual Science.—Conception of the World reformed by Anthropogeny.

"The Theory of Descent is a general inductive law which results with absolute necessity from the comparative synthesis of all the phenomena of organic nature, and especially from the threefold parallel of phylogenetic, ontogenetic, and systematic evolution. The doctrine that man has developed from lower Vertebrates, and immediately from genuine Apes, is a special deductive conclusion, which results with absolute necessity from the general inductive law of the Theory of Descent. This view of "

place in nature,' cannot, we believe, be made too prominent. If the Theory

Descent is correct as a whole, then the theory that man has developed from lower Vertebrates is simply an unavoidable deductive conclusion from that general inductive law. Hence, all farther discoveries which may in future enrich our knowledge of the phyletic development of man, can only be confirmative of special points of that deduction, which rests on the broadest inductive basis."—*Generelle Morphologie* (1866).

As we have now traversed the wonderful territory of the history of human development, and learned its most important parts, it seems appropriate that, at the close of our travels, we should look back on the road behind us, and, on the other hand, glance forward along the further path of knowledge into which our road will lead in future. We started from the simplest facts of the history of man's individual development; ontogenetic facts which can, at any moment, be shown and established by microscopic or anatomic research. The first and most important of these ontogenetic facts is, that every man, like every other animal, is at the commencement of his individual existence, a simple cell. This egg-cell exhibits precisely the same structure and mode of origin as that of any other Mammal. From this cell proceeds, by repeated division, a many-celled body, the mulberry-germ (*morula*); this changes into a cup-germ (*gastrula*), and this, again, into an intestinal germ-vesicle (*gastrocystis*). The two distinct cell-strata which compose its wall are the two primary germ-layers; the skin-layer (*exoderma*) and the intestinal layer (*entoderma*). This double-layered germ-form is the ontogenetic reproduction of that extremely important phylogenetic parent-form of all Intestinal Animals, to which we have given the name *Gastræa*.

As the human germ, like that of other Intestinal Animals,

passes through this gastrula-form, we are enabled to trace its phylogenetic origin back to the Gastraea. By tracing the germ-history of the two-layered germ still farther, we find that, by fission, four secondary layers are produced from the two original germ-layers. These have exactly the same constitution and genetic significance in Man as in all Vertebrates. From the skin-sensory layer develop the outer skin (*epidermis*) and the central nervous system; probably, the kidney system. The skin-fibrous layer develops into the leather-skin (*corium*) and the organs of motion (skeleton and muscle systems). From the intestinal-layer originates the vascular system and the fleshy part of the intestine. The intestinal-glandular layer, finally, gives rise to only the epithelium, or the inner cellular layer of the intestinal-mucous membrane and of the intestinal glands.

The manner in which these various organic systems develop from the four secondary germ-layers, is, from the very first, exactly the same in Man as in all other Vertebrates. The germ-history of each separate organ affords a proof that the human embryo takes exactly the same direction in its differentiation and formation, which, in Man, occurs only in the other Vertebrates. With the great animal tribe we then traced, step by step, and after stage, the farther development which takes place in the entire body as well as in all its several parts. This development takes place in the human embryo in the same manner as in the other Mammals. Finally, we saw, that even in this class the various stages of phylogenetic development, which determine the natural classification of Mammals, correspond throughout to the various stages of ontogenetic development through which the human embryo passes.

Further course of its development. We were thus enabled to determine the place of Man more definitely in the system of this class, and accordingly to establish the nature of his relation to the various mammalian orders.

The course of reasoning which we adopted in explaining these ontogenetic facts, was simply the logical carrying out of the fundamental law of Biogeny. In so doing we have constantly tried to carry out the significant distinction between palingenetic and kenogenetic phenomena. Palingenesis, or "the history of inheritance," alone enabled us to draw direct conclusions from observed germ-forms as to the tribal forms transmitted by heredity. On the other hand, these conclusions were more or less endangered, wherever Kenogenesis, or "vitiating evolution," was introduced by new adaptations. The whole understanding of the history of individual evolution depends on the recognition of this most important relation. We stand here on the border-line which sharply divides the new from the old method of scientific investigation, the new from the old conception of the world. All the results of recent morphological research drive us with irresistible force to the recognition of this fundamental principle of Biogeny, and of its far-reaching consequences. These are, it is true, irreconcilable with the customary mythological ideas of the world, and with the powerful prejudices engrafted into us in early youth by theosophic instruction; but, without this fundamental law of Biogeny, without the distinction between Palingenesis and Kenogenesis, and without the Theory of Descent, upon which these are based, we are entirely unable to understand the facts of organic development; without these, we cannot afford the faintest explanation of any part of this great and

wonderful world of phenomena. But, if we recognize the causal relation between the development of the germ and that of the tribe, if we recognize the true causal connection of Ontogeny and Phylogeny, which is expressed in that law, then the wonderful phenomena of Ontogeny explain themselves most simply; then the facts of germ-development appear but the necessary mechanical effects of the development of the tribe, conditioned by the laws of Heredity and Adaptation. The inter-operation of these laws among the everywhere-active influences of the struggle for existence—or, as we may simply say with Darwin, Natural Selection—is amply sufficient to explain to us the entire process of germ-history by the history of the tribe. Darwin's chief merit lies in the fact, that by the discovery of the interaction of the phenomena of Heredity and Adaptation, he prepared the way for a correct, logical understanding of the history of Evolution.

Among the numerous and important evidences that we have found for the truth of this view of our development history, I will only call attention here once more to the peculiarly valuable records of creation afforded by Dysteleology, or the doctrine of purposelessness, the science dealing with rudimentary organs. It is impossible to emphasize too often and too strongly the high morphological importance of those remarkable parts of the body, which are physiologically, completely worthless and useless. In every system of organs we find, in Man and in all higher Vertebrates, some of these worthless primæval heirlooms, which have been inherited from our lower vertebrate ancestors. Thus, first, we find on the outer surface of the body a scanty rudimentary covering of hair, which is thicker only on the

head, in the armpits, and on some other parts of the body. The short hairs on the greater part of the surface of our bodies are entirely useless, are without any physiological significance; they are the last scanty remains of the much more fully developed hairy covering of our Ape ancestors (p. 208). The sense-organs exhibit a series of the most remarkable rudimentary parts. As we have seen, the whole external shell of the ear, with its cartilages, muscles, and membranes, is, in Man, a useless appendage, destitute of the physiological importance that was formerly, erroneously, attributed to it. It is the atrophied remnant of the pointed, freely-moving, and much more highly developed mammalian ear, the muscles of which we retain, although we can no longer use them (p. 271). Again, we found, at the inner corner of the human eye, the remarkable little crescent-shaped fold, which is of no use to us, and is of interest only as being the last vestige of the nictitating membrane; of that third inner eyelid which is still of great physiological importance in Sharks and many Amnion Animals (p. 259). Numerous and interesting dysteleological proofs are also afforded by the apparatus of motion, both by the bony and the muscular systems. I will only cite the free, projecting tail of the human embryo, and the rudimentary caudal vertebræ developed in the latter, together with the pertinent muscles; this whole organ is entirely useless to Man, but is of great interest as the atrophied remnant of the long tail of our earlier Ape ancestors, which was composed of numerous vertebræ and muscles (p. 283). From these same ancestors we have also inherited various bone-processes and muscles, which were of great use to them in their climbing life among the trees, but with us have fallen out of use. At various points under the

skin we also have entirely unused skin-muscles; vestiges of the largely developed skin-muscles of our lower mammalian ancestors. It was the function of this "panniculus carnosus" to contract and wrinkle the skin, as we may see any day done by horses to drive away flies. We possess an active remnant of this great skin-muscle in the muscle of the forehead, by means of which we wrinkle the forehead and draw up the eyebrows; but we are not able to move at will another considerable remnant of this great skin-muscle of the neck (*platysma myoides*).

As in these animal organ-systems of our body, so also in the vegetative apparatus, we meet with many rudimentary organs, most of which we have incidentally noticed. I will only cite the remarkable thyroid gland (*thyreoidea*), the rudiment of the crop and the remnant of the ciliated gill (hypobranchial groove) present in Chordonia, *Ascidia*, and *Arcrania*, on the lower part of the gill-body (pp. 336, 337); also the vermiform process of the blind-intestine (*cæcum*) (p. 344). In the vascular system we find many rudiments, the vestiges of disused vessels which were formerly active blood-channels; such, for instance, are the "ductus Botalli," between the lung-artery and the aorta, the "ductus venosus Arantii," between the *vena portæ* and *vena cava*, and many others. The numerous rudimentary organs of the urinary and sexual systems (pp. 345-347) are especially interesting. Most of these are developed in one sex and rudimentary in the other. Thus, in the male the seed-ducts form from the Wolffian ducts, of which only traces remaining in the female are the Gartnerian canals. On the other hand, from the Müllerian ducts in the female are developed the oviducts and the uterus; wh

In the male, only the lower extremities of these ducts remain, forming the useless male uterus (*vesicula prostatica*). In the nipples and mammary glands, the male possesses other rudimentary organs which, as a rule, are functional only in the female (p. 204).

A closer anatomical examination of the human body would bring to our notice a number of other rudimentary organs, all of which can be explained only by the Theory of Descent. They are among the most important evidences for the truth of the mechanical theory of nature, and among the most overwhelming proofs against the prevailing teleological ideas of creation. If, in accordance with this latter view, Man and every other organism had been designed for his life-purpose from the beginning, and had been called into existence by an act of creation, the existence of these rudimentary organs would be an incomprehensible enigma; it would be impossible to understand why the Creator should have laid this useless burden on his creatures in their life-journey, so arduous at the best. On the other hand, by means of the Theory of Descent we can explain their existence in the most simple way, and say: The rudimentary organs are parts of the body, which, in the course of centuries, have gradually fallen out of use; organs which performed definite functions in our animal ancestors, but which, in us, have lost their physiological importance. They have become useless in consequence of our adaptation to new circumstances, but yet are transmitted from generation to generation by heredity, and have only slowly atrophied.

Like these rudimentary organs, so also all the other organs of our body have been transmitted to us from Mammals, and, immediately, from our Ape ancestors. The

human body includes no single organ which is not inherited from Apes ; but, by means of our fundamental law of Biogeny, we can trace the origin of our several systems of organs yet further down to various lower ancestral grades. Thus, for instance, we can say that we have inherited the earliest organs of our body, the outer skin (*epidermis*) and the intestinal canal, from the Gastræans, the nervous and muscular systems from the lower Worms (*Archelminthes*), the vascular system, body-cavity (*coelom*), and blood from Soft Worms (*Scolecida*), the notochord and the gill-intestine from Chorda Animals, the differentiated organs of sense from the Cyclostoma, the limbs and the Müllerian ducts from Primitive Fishes (*Selachii*), and the external reproductive organs from Primitive Mammals (*Præmammalia*). When we stated the "law of the ontogenetic connection of systematically allied forms," and determined the relative age of the organs, we saw how we could draw such phylogenetic conclusions as these from the ontogenetic succession of the organ-systems (vol. i. p. 390 ; ii. 357).

† By the help of this important law and of Comparative Anatomy, we were also enabled to determine definitely "man's place in nature," or, as we may say, to assign man his position in the system of the animal kingdom. It is now usual, in the more recent zoological systems, to distribute the whole animal kingdom into the seven tribes or *phyla*, which are again sub-divided, in round numbers, into about forty classes ; and these classes into about two hundred orders. According to his whole organization Man is undoubtedly, primarily, a member of but a single tribe, that of Vertebrates ; secondly, he is a member of but a single class, that of Mammals ; and, thirdly, a member

but a single order, that of the Apes. All the characteristic peculiarities, distinguishing Vertebrates from the other tribes, distinguishing Mammals from the other forty classes, and distinguishing Apes from the remaining two hundred orders of the animal kingdom, are also present in Man. Turn and twist as we may, we cannot escape this anatomical and systematic fact. Quite recently this very fact has led to the liveliest discussion, and has occasioned, especially, many disputes about the specific anatomical relationship of Man to Apes. The most astounding views on this "ape question," or "pithecoïd theory," have been uttered. It will therefore be well to examine it closely once more at this point, and to separate the essential from the non-essential in it.

We will start from the undisputed fact, that Man, at all events,—whether his special blood-relationship to Apes is acknowledged or denied,—is a genuine Mammal, is a Placental Mammal. This fundamental truth can be so easily proved at any moment by investigations in Comparative Anatomy, that it has been unanimously acknowledged since the separation of the Placental from the lower Mammals (Pouched Animals, or *Marsupialia*, and Beaked Animals, or *Ornithostoma*). But, from this, every logical adherent of the doctrine of development at once draws the conclusion, that man is descended from one and the same common parent-form, together with all other Placental Animals, from the progenitor of the *Placentalia*, just as, further, we must necessarily suppose a common mammalian ancestral form of all the various Mammals (*Placentalia*), Pouched Animals, and Cloacal Animals (*Monotremata*); but by this the great, all-agitating main question of man's place in nature is

conclusively settled, whether we ascribe to Man a nearer or a more remote relationship to Apes. No matter what Man is, in a phylogenetic sense, a member of the Ape order (or, if it is preferred, of the Primate order) or not, in any case, his direct blood-relationship to all other Mammals, and especially to the Placental Mammals, is established. It may be that the inter-relations of the various Mammals are quite different from those now hypothetically assumed, but, in any case, the common descent of Man and all other Mammals from a common parent-form is indisputable. This primæval, long since extinct parent-form, which probably developed during the Triassic Period, was the monotreme ancestral form of all Mammals.

If this fundamental and extremely significant principle is borne in mind, the "ape question" will appear to us in a wholly different light from that in which it is usually presented. A little reflection will bring conviction that this question has not the importance that has of late been attributed to it; for the origin of the human race from a series of various mammalian ancestors, and the historical development of the latter from an earlier series of lower vertebrate ancestors, remains indubitably established, in whatever matter whether the genuine "Apes" are regarded as the nearest animal ancestors of the human race or not. But it having become habitual to lay the principal weight on the entire question of the origin of man on this view of "descent from Apes," I find myself compelled to return once more to it here, and to recall those facts in Comparative Anatomy and Ontogeny, which conclusively settle this "ape question."

The shortest way to the goal is the one taken by

Huxley in his celebrated work, which we have so often cited, on the "Evidences as to Man's Place in Nature,"—the way afforded by Comparative Anatomy and Ontogeny. We have to compare objectively all the several organs of Man with the same organs in the higher Apes, and then to ascertain whether the differences between the former and latter are greater than the corresponding differences between the higher and lower Apes. The indubitable and indisputable result of this comparative anatomical investigation which was conducted with the greatest candour and accuracy, was the important law, which, in honour of its discoverer, we have named *Huxley's Law*; namely, that the physical differences between the organization of Man and that of the most highly developed Apes known to us, are much smaller than the corresponding differences between the higher and lower Apes. We might even define this law more exactly by excluding entirely the Platyrrhina or American Apes as being more remote relatives, and limiting our comparison to the narrower circle of relatives, the Catarrhina, or Apes of the Old World. Even within this small group of Mammals, we found the differences of structure between the higher and lower Narrow-nosed Apes, for example between the Gorilla and the Baboon, much greater than the differences between these Man-like Apes and Man. When, in addition, we now turn to Ontogeny, and when we find there, according to our "law or the ontogenetic connection of systematically related forms, that the embryos of Man and of the Man-like Apes, are identical for a longer period than the embryos of the highest and of the lowest Apes, we are certainly obliged to bring ourselves, whether with a good or a bad grace, to acknowledge our origin from

the Ape order. From the facts exhibited by Comparative Anatomy, we can undoubtedly form in imagination an approximate image of the structure of our ancestors of the older Tertiary Period; we may fill out the details as we will, yet this image will be a genuine Ape, and not a Catarrhine. For Man has all the physical characters distinguishing the Catarrhina from the Platyrrhina. Accordingly, in the mammalian pedigree, we must derive the human race directly from the Catarrhine group, and trace the origin of Man to the Old World. For the entire range of the Catarrhine Apes has, as yet, been confined to the Old World, just as the group of the Platyrrhine Apes has been limited to the New. Only the earliest root-form, that from which both groups sprang, was common to them; but it originated from the Semi-apes of the Old World.

Therefore, although it is thus indubitably established as the result of our objective scientific inquiry, that the human race is directly descended from the Apes of the Old World, yet we will once more state emphatically that this is a fact of no great importance to the main question of the origin of Man, as is generally supposed. For if we entirely ignore the fact or thrust it aside, it will not affect at all that the zoological facts of Comparative Anatomy and the history of development have testified concerning the placental character of Man. These facts prove the common descent of Man and the other Mammals. It is evident also, that the main question cannot be at least evaded or set aside by the statement: "Man is, in fact, a Mammal; but he branched off from the others at the root of the class, and has no nearer relationship to any other extant Mammal." At all events, the relat

idently more or less close if we comparatively examine the relation of the Mammalian class to the remaining forty per cent of the animal kingdom. All Mammals, including Man, are, at least, of common origin, and it is equally certain that their common parent-forms gradually developed through a long series of lower Vertebrates.

Feeling, evidently, rather than understanding, induces many people to combat the theory of their "descent from Apes." It is simply because the organism of the Ape appears a caricature of Man, a distorted likeness of ourselves in a very unattractive form, because the customary æsthetic pride and self-glorification of Man are touched by this in so sensitive a point, that most men shrink from recognizing their descent from Apes. It seems much pleasanter to be descended from a more highly developed, divine being, and hence, as is well known, human vanity has, from the earliest times, flattered itself by assuming the original parent of the race from gods or demi-gods. The church, with that sophistical distortion of ideas of which she is so great an adept, has managed to extol this ridiculous assumption as Christian humility; and those people who shudder with haughty horror every suggestion of descent from lower animals, and consider themselves children of God, those very people are exceedingly fond of boasting of their childlike humility of spirit. In most of the sermons delivered against the progress of the doctrine of evolution, human vanity and conceit play throughout a prominent part; and, although we have inherited this characteristic weakness from Apes, yet we must confess to have developed it to a degree of perfection which completely overthrows the unprejudiced judgment of the

"sound understanding of man." We ridicule the follies occasioned by the pride of ancestry among nobility, from the splendid Middle Ages down to our time, and yet no small portion of this groundless of nobility lurks in a great majority of men. Just as people prefer to trace their pedigree from a decayed or, if possible, from a celebrated prince, rather than an unknown, humble peasant, so they prefer seeing the genitor of the human race in an Adam degraded by the Fall rather than in an Ape capable of higher development and progress. It is a matter of taste, and such genealogical preferences do not, therefore, admit of discussion. I must confess that, personally, I am as proud of my paternal grandfather, who was simply a Silesian peasant, as of my maternal grandfather, who raised himself from the position of a Rhenish lawyer to the highest posts in the government of a state. And it is also much more to my individual pride to be the more highly developed descendant of a primate Ape ancestor, who, in the struggle for existence, developed progressively from lower Mammals, as the still lower Vertebrates, than the degraded descendant of an Adam, god-like, but debased by the Fall, who was created from a clod of earth, and of an Eve, created from the rib of Adam. As regards this celebrated "rib," I must here expressly add as a supplement to the history of the development of the skeleton, that the number of ribs is the same in man and in woman. In the latter as well as in the former the ribs originate from the skin-fibrous layer, and are regarded phylogenetically as lower or ventral vertebræ (1).

Now I certainly hear some one say: "That may be right and correct as far as the human body is concerned

the facts presented, it is certainly no longer to be doubted that this has actually developed gradually, step by step, from the long ancestral series of Vertebrates; but it is otherwise with the 'spirit of man,' with the human mind, which cannot possibly have developed in a similar manner from the mind of lower Vertebrates." Let us see if the known facts of Comparative Anatomy, Physiology, and Psychology can meet this grave objection. We shall best find firm ground from which to start in this matter by comparatively examining the minds of the different Vertebrates. Side by side within the various classes, orders, genera, and species of Vertebrates, we find so great a variety of vertebral intellects, that, at first sight, one can scarcely believe it possible that they can all be derived from the mind of a common "Primitive Vertebrate." First, there is the Lancelet, which has no brain at all, but only a simple notochord tube, the entire mental capacity remaining at the very lowest grade occurring among Vertebrates. The Cephalopods, also, standing just above, exhibit a hardly higher mental life, though they have a brain. Passing on to Fishes, we find their intelligence, as is well known, also at a very low point. Not until from these we ascend to the Amphibia, is any essential progress in mental development observable. This is much greater in Mammals, although, even here, in the Beaked Animals (*Ornithostoma*), and the lower higher class, the stupid Pouched Animals (*Marsupials*), the entire mental activity is still of a very low order; but we pass on from these to Placental Animals, within this uniform group we find such numerous and important steps in differentiation and improvement, that the mental differences between the most stupid Placental Animals (for

instance, Sloths and Armadillos) and the most intelligent animals of the same group (for instance, Dogs and Horses) seem much more considerable than the intellectual differences between those lowest Placentals and the Reptile Animals, or even the lower Vertebrates. Those differences are, at any rate, much more considerable than the differences in the intellectual life of dogs, apes, and men. yet all these animals are allied members of a single class.

This fact is shown to a yet more surprising degree by the Comparative Psychology of another class of animals, the Insects. It is well known that many Insects exhibit mental capacity approximately as highly developed as that possessed by Man only of the vertebrate group. It is not to speak of the celebrated organized communities and social arrangements of Bees and Ants; every one knows that very remarkable social arrangements occur among these, such as occur in the higher mammals, and nowhere else in the animal kingdom. I will allude to the civil organization and government of Monarchical bees and Republican ants, to their division into various orders: the queen, the drone nobility, the workers, the nurses, soldiers, and so on. Among the most remarkable phenomena in this extremely interesting life, is certainly the cattle-keeping of certain Ants, which tend plant-lice for the sake of their milk and regularly collect their honey-juice. Even more remarkable is the slave-holding of the large red Ants, which steal the eggs of the small black species and rear them to slaves. It has long been known that all these civil and social arrangements of the Ants were originated by the system

-operation of numerous citizens, understanding each other. Numerous observations have placed the astoundingly high intellectual development of these little Articulated Animals beyond all doubt. With this let us compare, as Darwin has done, the intellectual capacity of many lower, and, specially, of many parasitic, Insects. There, for example, are the Scale Insects (*Coccus*) which, when mature, consist of an entirely immovable shield-shaped body attached to the leaves of plants. Their feet are atrophied. Their mouths are embedded into the tissue of the plant, the juices of which they suck. The whole mental activity of this motionless female parasite consists in the enjoyment it derives from sucking these juices and from sexual intercourse with the unattached male. The same is true of the maggot-like female of the Twisted-wings (*Strepsiptera*), which spends its whole life, wingless and footless, as a motionless parasite in the body of the wasp. There can be no suspicion of any higher mental activity there. If these brutish parasites are compared with the mentally active and sensible ants, it will certainly be admitted, that the psychical differences between the two are much greater than those between the highest and lowest Mammals, between Beaked Animals (*Ornithostoma*), Pouched Animals (*Marsupialia*), and Armadillos on the one hand, and Dogs, Apes, and Men on the other. And yet all those insects belong, without question, to the single class of Arthropoda, just as all these Mammals undoubtedly belong to the single class of Vertebrates; and just as every logical adherent of the doctrine of evolution must assume a common parent-form for all those Insects, so also he must necessarily assert a common descent for all these Mammals.

Turning now from observing the comparative capacity of the various animals to the question as to the organs of these functions, we receive the answer, that in the higher animals they are invariably connected with certain groups of cells, those cells which compose the central nervous system. All naturalists, without exception, maintain that the central nervous system is the organ of the mental life of animals, and this assertion is at any time capable of experimental proof. If the central nervous system is wholly or partially destroyed, the "mind," or the psychological activity of the animal, is wholly or partially annihilated at the same time. We must, therefore, next inquire what is the character of the mental organ in man. The undeniable answer to this question has already been given. Man's mental organ is, in its whole structure and origin, the same as that of all other Vertebrates. It originates as a simple medullary tube from the outer skin of the embryo, from the skin-sensory layer, or the first of the secondary germ-layers. In the course of its gradual development it passes through the same stages of progression in the human embryo as in that of all other Vertebrates, and as these latter have undoubtedly a common origin, so must also the brain and spinal cord be of the same origin in all.

Physiological observation and experiment teaches, moreover, that the relation of the "mind" to its organ, the brain and spinal marrow, is exactly the same in Man as in all other Mammals. The former can in no case act without the latter; the one is connected with the other, as is muscular movement with muscle. Therefore, the mind can develop only in connection with its organ. Adherents of the Theory of Descent, who concede the causal connection

On Ontogeny and Phylogeny, are now compelled to make the following propositions: The mind, or "psyche," has developed together with, and as the function of the medullary tube, and just as even now the brain and spinal cord develop in each human individual from the medullary tube, so the human "mind," or the mental faculties of the entire human race, has developed gradually, step by step, from the mind of lower Vertebrates. Just as now in every individual of the human race the full and complex structure of the brain develops step by step from exactly the same rudiment, from the very simple brain-bladders, as in all other Skulled Animals (*Craniota*), so the human mind has gradually developed in the course of millions of years from the mind of the lowest Skulled Animals; and as now the brain of every embryo differentiates according to the special type of Ape-brain, so also the human psyche has historically differentiated from the Ape-mind.

The monistic idea will, of course, be indignantly rebuffed by most people, who accept the contrary dualistic view which denies the inseparable connection of the brain and mind, and regards "body and mind" as entirely separate and distinct; but how shall we reconcile this long accepted view with the facts taught by the theory of evolution? The dualistic view is, at least, as completely opposed to Ontogeny as to Phylogeny. If we agree with the majority of men, that the mind is a self-existent, independent being, which has originally nothing to do with the body, but only dwells in it for a time, and gives expression to its emotions through the brain, as a piano-player through his instrument, then we must

suppose a period in the human germ-history, at which the mind enters the body, enters the brain; and we must also suppose a moment at death, at which it leaves the body, and further, as every man inherits certain individual mental qualities from each parent, we must suppose the portions of the mind of each were transferred to the germ at the time of its procreation. A little piece of the father's mind accompanied the sperm-cell, a little piece of the mother's mind remained with the egg-cell. This dualistic view entirely fails to explain the phenomena of evolution. We all know that the new-born child has no consciousness, no knowledge of itself and of the objective world. Whoever has children of his own, and follows their mental development candidly, cannot possibly deny that processes of biological evolution are at work there. Just as all other functions of the body develop in connection with their organs, so does the mind develop in connection with the brain. And this gradual development of the child's mind is such a wonderful and beautiful phenomenon, that every mother and every father with eyes to see takes unwearied delight in observing it. The text-books of Psychology alone are ignorant of any such development, and we are almost forced to the conclusion that their authors themselves never had any children. The human mind, as it is represented in the great majority of psychological works, is only the one-sided mind of a learned philosopher, who, indeed, knows many books, but nothing of the process of evolution, and does not suspect that even his own mind has developed.

These same dualistic philosophers must, of course, if they are consistent, also assume that there was a moment

the Phylogeny of the human mind at which this mind first entered the vertebrate body of man. Accordingly, at the time when the human body developed from the body of the Anthropoid Ape (thus, probably, in the latter part of the Tertiary Period), a specific human mind-element—or, as it is usually expressed, a “divine spark”—must have suddenly entered or been breathed into the brain of the Anthropoid Ape, and there have associated itself with the already existing Ape-mind. I need not point out the theoretic difficulties involved in this conception. I will only remark that even this “divine spark,” by which the mind of Man is said to be distinguished from that of all other animals, must itself be a thing capable of evolution, and has actually developed progressively in the course of human history. This “divine spark” is usually understood to be “reason,” and is ascribed to man as a mental function distinguishing him from all “irrational animals.” Comparative Psychology, however, teaches that this frontier-post between man and beast is altogether untenable.¹⁹⁸ We must either take the idea of reason in its broader sense, in which case it belongs to the higher Mammals (the Ape, Dog, Elephant, Horse), as much as to the majority of men; or we must conceive it in its narrower sense, and then it is lacking in the majority of men, as well as in most animals. On the whole, that which Goethe's Mephistopheles said of his time, is true of Man's reason to-day:

“He might have kept himself more right
Hadst Thou ne'er shewn to him a glimpse of heaven's light.
He calls it Reason, but Thou seest
Its use but makes him beastlier than the beast.”

If, therefore, we must abandon this generally preferred.

and, in many respects, very pleasant dualistic theory of mind, as being wholly untenable, because irreconcilable with genetic facts, then the opposite monistic view remains to us, according to which the human mind, like that of any other animal, is a function of the central nervous system, with which it has developed in inseparable connection. Ontogenetically, we see this in every child; phylogenetically, we must assert it in accordance with the fundamental law of Biogeny. In every human embryo the medullary tube develops from the skin-sensory line and from the anterior part of that tube the five brain ventricles of Skulled Animals (*Craniota*), and from that the mammalian brain (at first with the characteristics of the lower, then with those of the higher Mammals). Just as this entire ontogenetic process is but a short reduction, occasioned by Heredity, of the same process in the Phylogeny of Vertebrates, so also the wonderful mental activity of the human race has gradually developed, by step, in the course of many thousands of years, from the less perfect mental activity of the lower Vertebrates. The evolution of the mind in each child is only a reproduction of that long phylogenetic process.

The extraordinary and important bearing of Antropogeny on Philosophy, in the light of the fundamental principle of Biogeny, now becomes apparent. The speculative philosophers who will take possession of the facts of Antropogeny and explain them phylogenetically (according to the law), will introduce a greater advance in the history of Philosophy than has been made by the greatest thinkers of all previous centuries. Undoubtedly every clear and logical thinker must draw from the facts of Comparative Anat-

and Ontogeny which have been brought forward, a mass of suggestive thoughts and reflections which cannot fail of their effect on the further development of the philosophical study of the universe. Neither can it be doubted that these facts, if properly weighed, and judged without prejudice, will lead to the decisive victory of that philosophical tendency, which we distinguish, briefly, as monistic or mechanical, in distinction from the dualistic or teleological, on which most philosophical systems of ancient, mediæval, and modern times are based. This mechanical, or monistic philosophy, asserts that everywhere the phenomena of human life, as well as those of external nature, are under the control of fixed and unalterable laws, that there is everywhere a necessary causal connection between phenomena, and that, accordingly, the whole knowable universe forms one undivided whole, a "*monon*." It further asserts, that all phenomena are produced by mechanical causes (*causæ efficientes*), not by pre-arranged, purposive causes (*causæ finales*). Hence there is no such thing as "free-will" in the usual sense. On the contrary, in the light of this monistic conception of nature, even those phenomena which we have been accustomed to regard as most free and independent, the expressions of the human will, appear as subject to fixed laws as any other natural phenomenon. Indeed, each unprejudiced and searching test applied to the action of our "free-will" shows that the latter is never really free, but is always determined by previous causal conditions, which are eventually referable either to Heredity or to Adaptation. Accordingly, we cannot assent to the popular distinction between nature and spirit. Spirit exists everywhere in nature, and we know of no spirit out-

side of nature. Hence, also, the usual distinction between natural science and mental science is entirely untrue. Every real science is at the same time both a natural and a mental science. Man is not above nature, but in nature.

The opponents of the doctrine of evolution are very fond of branding the monistic philosophy grounded upon "materialism," by confusing *philosophical* materialism with the wholly different and censurable *moral* materialism. Strictly, however, our "monism" might, as accurately as materialism, be called spiritualism as materialism. Real materialistic philosophy asserts, that the vital phenomena of motion, like all other phenomena of motion, are effects or products of matter. The other, opposite or spiritualistic philosophy, asserts, on the contrary, that matter is the product of motive force, and that all natural forms are produced by free forces entirely independent of the matter itself. Thus, according to the materialistic conception of the universe, matter, or substance, precedes motion, or active force. According to the spiritualistic conception of the universe, on the contrary, active force precedes matter. Both views are dualistic, and hold them both to be equally false. A contrast between the two views is presented in the *monistic* philosophy, which neither little believe in force without matter, as in matter without force. It is only necessary to reflect on this for a moment from a strictly scientific standpoint, to find that on a closer examination it is impossible clearly to represent motion without the other. As Goethe says, "Matter cannot exist and act without spirit; neither can spirit exist without matter."¹²

The "spirit" and "mind" of man are but forces

are inseparably connected with the material substance of our bodies. Just as the motive force of our flesh is involved in the muscular form-element, so is the thinking force of our spirit involved in the form-element of the brain. Our spiritual forces are as much functions of this part of the body, as every force is a function of a material body. We know of no matter which does not possess force, and, conversely, of no forces that are not connected with matter. When the forces manifest themselves in the phenomena of motion, they are called active forces; if, on the other hand, the forces are in a state of rest, or of equilibrium, they are called latent forces.¹⁹⁰ This is as true of inorganic natural substances as of organic. The magnet attracting iron-filings, powder exploding, steam driving the locomotive, are active inorganic substances; they work by active force just as does the sensitive mimosa, when it folds its leaves at a touch,—as does the *Amphioxus*, when it buries itself in the sand,—as does man, when he thinks. Only in these latter cases the combination of the different forces, appearing as phenomena of motion, are much more complex and much less easily recognized than in the former cases.

Anthropogeny has led us to the conclusion that even in the entire history of the evolution of man, in the history of the germ, as well as in that of the tribe, no other active forces have been at work, than in the rest of organic and inorganic nature. All the forces at work there can be reduced at last to *growth*—to that fundamental function of evolution by which the forms of inorganic, as well as of organic bodies, originate. Growth, again, itself rests on the attraction and repulsion of like and unlike particles.¹⁹¹ It has given rise to Man and to Ape, to Palm and Alga, to

crystal and water. Hence the evolution of man has taken place according to the same "eternal, immutable law," as has the evolution of any other natural body.

It is true that the prejudices that stand in the way of the general recognition of this "Natural Anthropogeny" are even yet intensely powerful; otherwise the ancient strife between the various philosophical systems would already have been decided in favour of "Monism." But it can be foreseen with certainty that a more general acquaintance with genetic facts, will gradually destroy those prejudices and bring about the victory of the natural idea of "Man's Place in Nature." The fear is often expressed in opposition to this view that it will cause a retrogression in the intellectual and moral development of man; but, on the contrary, I cannot withhold my conviction, that the very reverse will be true, that by it the progressive development of the human spirit will be advanced in an unusual degree. At all events, I hope and trust that I have, in these chapters, afforded convincing proof that the only way to attain a true scientific knowledge of the human organism, is by employing the method which we must acknowledge to be alone correct and successful in the study of organic nature,—by following the course of the History of Evolution.²⁰⁰

NOTES.

REMARKS AND REFERENCES TO LITERATURE.

1 (vol. i. p. 2). Anthropogeny (Greek) = History of the Evolution of Man; from Anthropos (*ἄνθρωπος*) = man, and genea (*γενεά*) = Evolution history. There is no especial Greek word for "the history of evolution;" in its place is used either *γενεά* (= descent), or *γονεία* (= generation). If goneia is preferred to genea, the word must be written Anthropogony. The word "*Anthropogony*," used first by Josephus, means, however, only "the generation of man." Genesis (*γένεσις*) means "origination, or evolution;" therefore *Anthropogenesis* = "the evolution of man."

2 (i. 3). Embryo (Greek) = germ (*ἔμβρυον*). Really τὸ ἐντὸς τῆς γαστρὸς βρύον (Eust.), i.e. the unborn germ in the mother's body (Latin *fœtus*, or, better, *fetus*). In accordance with this original sense, the term embryo should only be applied to those young organisms which are still enclosed in the egg-coverings. (Cf. "*Generelle Morphologie*," vol. ii. p. 20.) Inaccurately, however, various free-moving young forms of low animals (*larvæ*) are often spoken of as embryos. Embryonic life ends at birth.

3 (i. 5). Embryology (Greek) = Germ-science, from embryo (*ἔμβρυον*) = germ, and logos (*λόγος*) = science. Even now the whole history of the evolution of the individual is erroneously called "embryology." For corresponding with the term "embryo" (see note 2), by "embryology," or "embryogony," should only be understood "the history of the evolution of the

individual within the egg-coverings." As soon as the organism has left there, it is no longer a real "embryo." The later changes of this form the subject of the science of Metamorphoses, or *Metamorphology*.

4 (i. 5). Ontogeny (Greek) = "germ-history," or "the history of the evolution of the individual;" from *ὄντα*=individuals, and *γενεά* (γενεά)=history of evolution. (Cf. note 1.) Ontogeny, as the "history of the evolution of the individual," embraces both Embryology and Metamorphology (note 3).—"Generelle Morphologie," vol. ii. p. 30.

5 (i. 5). Phylogeny (Greek) = tribal history, or "the palæontological history of evolution;" from *phylon* (φύλον)=tribe and *γενεά* (γενεά)=history of evolution. The phylon includes all organisms connected by blood, which are descended from common typical parent-form. Phylogeny includes Palæontology and Genealogy.—"Generelle Morphologie," vol. ii. p. 305.

6 (i. 6). Biogeny (Greek) = the history of the evolution of organisms or of living natural bodies in the widest sense (Genea tu biu.) βίος=life.

7 (i. 6). The fundamental law of Biogeny. Cf. my "General History of the Evolution of Organisms" ("Generelle Morphologie," 1866, vol. ii.), p. 300 (Essays on the causal connection of biogeny and phyletic evolution); also the "Monograph of C Sponges" ("Monographie der Kalkschwämme," 1872, vol. i. 4) also my "Natural History of Creation."

8 (i. 10). Palingenesis (Greek) = original evolution, or palingenesia (παλιγγενεσία) = new-birth, renewal of the former course of evolution. Therefore, Palingeny = inherited history (from *παλιν*=reproduced, and *γενεά*=history of evolution).

9 (i. 10). Kenogenesis (Greek) = modified evolution, or kenos (κενός)=strange, meaningless; and *γενεά* (γενεά)=history of evolution. The modifications introduced into Palingeny by Kenogenesis are vitiations, strange, meaningless additions to the original, true course of evolution. Kenogeny = vitiated history.

10 (i. 12). Latin definition of the fundamental law

my: "Ontogenesis summarium vel recapitulatio est physeos, tanto integrius, quanto hereditate palingenesis concur, tanto minus integrum, quanto adaptatione kenogenesis vincitur." Cf. my "Aims and Methods of Recent History of Evolution" ("Ziele und Wege der Hentigen Entwicklungsgeschichte," p. 77. Jena, 1876).

(i. 17). Mechanical and purposive causes. Mechanical philosophy assumes that throughout nature, in organic as in inorganic processes, only non-purposive, mechanical, regularly-working causes exist (*causæ efficientes*, *mechanism*, *regularity*). On the other hand, vitalistic natural philosophy holds that the latter are at work only in inorganic processes, while in certain other, purposive, special causes are at work, such as purposive causes, working for a definite end (*causæ finales*, *Vitalism*, *Teleology*). (Cf. "Generelle Morphologie," p. 94.)

(i. 17). Monism and Dualism. Unitary philosophy, or Monism, is neither extremely materialistic nor extremely spiritualistic, but resembles rather a union and combination of these two principles, in that it conceives all nature as one whole everywhere recognizes any but mechanical causes. Binary philosophy, on the other hand, or Dualism, regards nature and matter and force, inorganic and organic nature as distinct dependent existences. (Cf. vol. ii. p. 456.)

(i. 20). Morphology and Physiology. Morphology (as the science of forms) and Physiology (as the science of the functions of organisms) are indeed connected, but co-ordinate sciences, independent of each other. The two together constitute the "science of organisms." Each has its peculiar methods and aids. (Cf. "Generelle Morphologie," vol. i. pp. 1-10.)

(i. 24). Morphogeny and Physiogeny. Biogeny, or the study of the evolution of organisms, up to the present time has been almost exclusively Morphogeny. Just as this first step is the way to a true knowledge of organic forms, so will Physiogeny afterwards make a true recognition of functions

possible, by discovering their historic evolution. Its promises to be most fruitful. Cf. "Aims and Methods of Recent History of Evolution" ("Ziele und Wege der Historie der Entwicklungsgeschichte," pp. 92-98. Jena, 1876).

15 (i. 27). Aristotle. Five books on the generation and evolution of animals (*περὶ ζῴων γενέσεως*).

16 (i. 28). Parthenogenesis. On "virginal generation or the "immaculate conception" of Invertebrates, especially Articulated Animals (*Crustacea, Insecta*, etc.), see Seidlitz, "Remarks on Parthenogenesis among Arthropoda" ("Beiträge zur Parthenogenesis der Arthropoden." Leipzig, 1871). Seidlitz, "Parthenogenesis and its Relation to other Forms of Generation in the Animal Kingdom" ("Die Parthenogenesis und ihr Verhältniss zu den übrigen Zeugungs-Arten im Thierreich." Leipzig, 1872).

17 (i. 34). The Preformation-theory. This theory is in Germany, usually called "Evolutions-theorie," in distinction to the "Epigenesis-theorie." As, however, in England, France, Italy, the latter is, on the contrary, usually called the theory of evolution, evolution and epigenesis being used as synonymous terms, it appears better to call the former "the theory of preformation." Recently Kölliker has called his "theory of homogeneous generation" "Evolutionism" (note 47). Cf. p. xxx.

18 (i. 37). Alfred Kirchhoff, "Caspar Friedrich Wolff: His Life and Teaching in the Science of Organic Evolution," "Jenaische Zeitschrift für Naturwissenschaft," 1868, vi. p. 193.

19 (i. 43). Part of the writings left by Wolff have not been published. His most important works are the dissertation for the degree of doctor, *Theoria generationis* (1759), and a model treatise "de formatione intestinorum" (on the formation of the intestinal canal).—"Nov. Comment. Acad. Sc. Petro-polit." xii. 1768; xiii. 1769. Translated into German by M. Halle, 1812.

20 (i. 51) Christian Pander, "*Historia metamorphoseos*,"

um incubatum prioribus quinque diebus subit." Vicebergi, 17. (Dissertatio inauguralis.) "Contributions toward the story of the evolution of the chick within the egg." ("Beiträge zur Entwicklungsgeschichte des Hühnchens im Eie." Würzburg, 1817.)

21 (i. 52). Karl Ernst Baer, "On the Evolution of Animals. Observations and Reflections" ("Ueber Entwicklungsgeschichte der Thiere. Beobachtung und Reflexion." 2 vols. Königsberg, 1827-1837). In addition to this chief work, cf. "Story of the Life and Writings of Dr. Karl Ernst Baer, told by himself" ("Nachrichten über Leben und Schriften des Dr. Karl Ernst Baer, mitgetheilt von ihm selbst." Petersburg, 1865).

22 (i. 60). Albert Kölliker. His "History of the Evolution of Man and the Higher Animals" ("Entwicklungsgeschichte des Menschen und der höheren Thiere"). The 2nd (corrected) edition, 1876, contains (pp. 28-40) a catalogue of ontogenetic literature. On the newer contributions to this, cf. the "Jahresberichte über die Leistungen und Fortschritte der Medicin" (Berlin), by Virchow and Hirsch (the "History of Evolution," by Waldeyer); also the "Jahresberichte über die Fortschritte der Anatomie und Physiologie," by Hofmann and Schwalbe (Leipzig); the "History of Evolution," by R. Hertwig and Nitsche. Most of Kowalevsky's researches are contained in the "Memoires de l'Académie impériale de St. Petersburg" (from the year 1866). Others are published in Max Schultze's "Archiv für mikroskopische Anatomie," and in other periodicals.

23 (i. 60). Theodor Schwann, "Microscopic Researches into the Identity in Structure and Growth of Plants and Animals" ("Mikroskopische Untersuchungen über die Uebereinstimmung in der Structur und Wachsthum der Thiere und Pflanzen." Berlin, 1839).

24 (i. 69). Ernst Haeckel, the Gastræa Theory, phylogenetic classification of the animal kingdom and homology of the germ-layers.—"Jenaische Zeitschrift für Naturwissenschaft," vol. viii. 1874, pp. 1-56.

25 (i. 75). Ernst Haeckel, "The History of Creation." London, 1876.

26 (i. 81). Fritz Schultze, "Kant and Darwin." A contribution to the history of the science of evolution. Jena, 1871.

27 (i. 81). Immanuel Kant, "Critique of Teleological Reason" ("Kritik der teleologischen Urtheilskraft"). 1790. §§ 64 and § 79. Cf. also my "History of Creation," vol. i. p. 103.

28 (i. 83). Jean Lamarck, "Philosophie Zoologique, ou Exposition des Considérations relatives à l'histoire naturelle des animaux," etc. 2 Tomes. Paris, 1809. Nouvelle édition, revue et précédée d'une introduction biographique par Charles Martin. Paris, 1873.

29 (i. 88). Wolfgang Goethe on Morphology (zur Morphologie). The formation and re-formation of organic bodies. On Goethe's morphological studies, cf. Oscar Schmidt ("Goethe's Verhältniss zu den organischen Naturwissenschaften." Jena, 1853). Rudolph Virchow, "Goethe as a Naturalist" (Berlin, 1861). Helmholtz, "On Goethe's Natural Scientific Works" (Brunswick, 1865).

30 (i. 96). Charles Darwin. His chief work is "On the Origin of Species by means of Natural Selection" (1859).

31 (i. 99). Darwin and Wallace. The general outlines of the theory of selection were discovered independently by Darwin and Wallace. It does not, however, follow that the services of the latter in furthering the science of evolution are at all comparable with those of the former. As many opponents of Darwin, especially the English Jesuit Mivart, have recently endeavoured to exalt Wallace at the expense of Darwin, and depreciate the latter, I take this opportunity of expressly asserting that Darwin's services are very far the greater.

32 (i. 101). Thomas Huxley. In addition to the works mentioned in the text, the following popular works are especially to be recommended: "On Our Knowledge of the Causes and Phenomena in Organic Nature," and the "Elementary Physiology" (1871).

33 (i. 101). Gustav Jaeger, "Zoological Letters" ("Zoologische Briefe." Vienna, 1876), and the "Text-book of General Zoology" ("Lehrbuch der Allgemeinen Zoologie." Stuttgart, 1875).

34 (i. 101). Friedrich Rolle, "Man, his Descent and Morality presented in the light of the Darwinian Theory, and on the basis of Recent Geological Discoveries" ("Der Mensch, seine Abstammung und Gesittung im Lichte der Darwin'schen Lehre," &c.). Frankfort, 1866.

35 (i. 102). Ernst Haeckel, "Generelle Morphologie der Organismen." General outlines of the science of organic forms, mechanically shown in accordance with the theory of descent as formed by Charles Darwin. Vol. i., "General Anatomy;" Vol. ii., "General History of Evolution." Berlin, 1866.

36 (i. 103). Charles Darwin, "The Descent of Man, and Selection in Relation to Sex." 2 vols. London, 1871.

37 (i. 108). Karl Gegenbaur, "Outlines of Comparative Anatomy" ("Grundzüge der vergleichenden Anatomie." Leipzig, 1st ed., 1870). "Elements of Comparative Anatomy" ("Grundriss der vergleichenden Anatomie." 3rd (improved) edition, 1874).

38 (i. 114). Migration-theory. Moritz Wagner, "The Darwinian Theory and the Law of Migration of Organisms" ("Die Darwin'sche Theorie und das Migrations-gesetz der Organismen." Leipzig, 1868). August Weismann, "On the Influence of Isolation in the Formation of Species" ("Ueber den Einfluss der Isolirung auf die Artenbildung." Leipzig, 1871).

39 (i. 116). Carus Sterne, "Evolution and Dissolution" ("Werden und Vergehen"). A popular history of the evolution of Nature as a whole. Berlin, 1876. Agassiz a "founder" of natural science. "Gegenwart." Berlin, 1876.

40 (i. 117). Ernst Haeckel, "The Chalk-sponges" ("Die Kalkschwämme; Calcispongien oder Grantien." Berlin, 1872). A monograph and an attempted solution of the problem of the origin of species. Vol. i., "Biology of Chalk-sponges;" vol. ii., "Classification of Chalk-sponges;" vol. iii., "Atlas of Chalk-sponges" (with 60 plates).

41 (i. 124). On the Individuality of Cells and recent reforms of the cell-theory, cf. my "Individualitätslehre," or "Tectologie" ("Generelle Morphologie," vol. i. pp. 239-274). Rudolf Virchow, "Cellular Pathologie." 4th edition. Berlin, 1871.

42 (i. 130). "The Plastid-theory and the Cell-theory" "Jenaische Zeitschrift für Naturwissenschaft," 1870, vol. i. 492.

43 (i. 138). Gegenbaur, "On the Structure and Evolution of Vertebrate Eggs with Partial Yolk-cleavage."—"Archiv f. d. n. Phys." 1861, p. 491.

44 (i. 153). Ernst Haeckel, "On Division of Labour in Nature and Human Life," in the collection of Lectures by Virchow, Holtzendorf, 1869. Sect. 78. 2nd edition.

45 (i. 160). Monogony (*Generatio neutralis*). On the various forms of asexual reproduction (Schizogony, Sporogony, etc.). "Generelle Morphologie," vol. ii. pp. 36-58.

46 (i. 160). Amphigony (*Generatio sexualis*). On the various forms of sexual reproduction (Hermaphroditism, Gonochorism, etc.), see "Generelle Morphologie," vol. ii. pp. 58-69.

47 (i. 168). Fitful evolution and gradual evolution. The theory of fitful evolution has recently been developed especially by Kölliker, who, under the title of heterogeneous genealogies, opposes it to gradual evolution as maintained by us ("Zeitschrift f. Wissensch. Zool.," vol. xiv. 1864, p. 181, and "Alcyonaria," pp. 384-415). This theory is distinguished by assuming *unknown causes* for the "fitful evolution of species," as so-called "great law of evolution" (an empty word indeed!). On the contrary, we see, with Darwin, in the facts of Heredity and Adaptation sufficient known (partly inner, partly external) physiological causes, which explain the gradual evolution of species under the influence of the struggle for existence.

48 (i. 170). Immaculate Conception never occurs in any vertebrate tribe. On the other hand, parthenogenesis frequently occurs among Articulated Animals (*Arthropoda*) (note 16).

49 (i. 171). Fertilization of Flowers by insects. (See Darwin on "The various contrivances by which British and Foreign Orchids are fertilized by Insects." Hermann Müller, "The Fertilization of Flowers by Insects, and the corresponding adaptations of both" ("Die Befruchtung der Blumen durch Insecten und die gegenseitigen Anpassungen Beider").

tion to our knowledge of causal connection in organic life. Leipzig, 1873.

50 (i. 178). The Process of Fertilization has been very variously viewed, and was formerly often regarded as an especially mysterious process, or even as a supernatural miracle. Now appears no more "wonderful or supernatural" than the process of digestion, of muscular movement, or of any other biological function. For the earlier views, cf. Leuckart, article "Zeugung" (generation) in R. Wagner's "Dictionary of Physiology," 1850.

51 (i. 179). Monerula. The simple, very transient, kernel-condition, which we briefly call the "monerula," and, in accordance with the fundamental law of Biogeny, regard as an ontogenetic reproduction of the phylogenetic Moneron parent-cell, appears to vary to some extent in different organisms, especially in the matter of duration. In those cases in which it no longer occurs, and in which the kernel of the fertilized cell persists wholly or partially, we may regard this phenomenon as a later, kenogenetic curtailment of Ontogeny.

52 (i. 181). The Plasson of the monerula appears, morphologically, a homogeneous and structureless substance, like that of the Moneron. This is not contradicted by the fact that we ascribe a very complex molecular structure to the plastidules, "plasson-molecules," of the monerula; this latter will naturally be more complex in proportion as the organism which ontogenetically constitutes it is higher, and as the ancestral series of that organism is longer, in proportion as the preceding processes of Heredity and Adaptation are more numerous.

53 (i. 182). The Fundamental Significance of the Parent-cell, Zytula, as the foundation-stone of the young organism in the course of development, can only be rightly appreciated, if the part taken in its constitution by the two generating cells is fully appreciated, the part taken by the male sperm-cell and the female egg-cell.

54 (i. 183). The One-celled Germ-organism, like the act of fertilization from which it results, has been very variously

viewed. Cf. on this subject, in addition to the four important works, here quoted, by Auerbach, Bütschli, Hertwig, and Stemberger, the most recent annals of the progress of the history of evolution (Waldeyer in Virchow-Hirsch's "*Jahresberichte*," Berlin; Hertwig in Hofmann-Schwalbe's "*Jahresberichte*," Leipzig).

55 (i. 185). Protozoa and Metazoa. Cf. vol. i. p. 248; ii. 248. The Protozoa and Metazoa are genetically and anatomically very distinct, that the former, as Protista, may even be excluded entirely from the animal kingdom, and may be regarded as a neutral intermediate kingdom between the plant and animal kingdoms.—"*Generelle Morphologie*," vol. i. pp. 191-230. According to this view the Metazoa alone are really animals.

56 (i. 186). The Unity of the Zoogenetic Conception, resting on the Gastræa-theory, has as yet not been destroyed by the numerous attacks directed against that theory: for none of these attacks have succeeded in substituting anything positive by pure negation no advance can be made in this dark and difficult subject.

57 (i. 187). The Egg-cleavage and Gastrulation of Man represented diagrammatically in Figs. 12-17 of Plate II., is probably in no essential way different from that of the Rat, which has as yet been most closely examined in this point.

58 (i. 188). Ernst Haeckel, "Arabian Corals" ("*Arabische Korallen*"). "A Journey to the Coral Banks of the Red Sea, a Glimpse into the Life of Coral Animals. A popular lecture with scientific explanations." With 5 coloured plates, and woodcuts. Berlin, 1876.

59 (i. 189). The Number of the Segmentella, or cleavage cells, increases, in the original, pure forms of palingenetic cleavage, in regular geometric progression. But the point at which this proceeds varies in the various archiblastic animals so that the Morula, as the final result of the cleavage-process, consists sometimes of 32, sometimes of 64, sometimes of 128 cells, and so on.

60 (i. 189). The Mulberry-germ, or Morula. The

entella, or cleavage-cells, which constitute the Morula at the close of palingenetic egg-cleavage, generally appear entirely similar, with morphological difference in size, form, or constitution. This does not, however, hinder the fact that these cells have separated, even during cleavage, into animal and vegetable cells, have differentiated physiologically, as is indicated in Figs. 2 and 3, Plate II., as probable.

61 (i. 189). The Bladder-germ of Archiblastic Animals (*Blastula*, or *blastosphæra*), which is now commonly known as the germ-vesicle, or, more accurately, as the "germ-membrane vesicle," must not be confused with the essentially different "germ-vesicle" of amphiblastic mammals, which is better called the "intestinal-germ vesicle" (*gastrocystis*). The *gastrocystis* and the blastula are still often united under the name of "germ-vesicle, or *vesicula blastodermica*." Cf. vol. i. p. 290.

62 (i. 192). The Definition of the Gastrula was first established by me in 1872, in my "Monograph of Chalk-sponges" (vol. i. pp. 333, 345, 466). There I already gave due weight to the "extremely great significance of the gastrula in reference to the general Phylogeny of the animal kingdom" (p. 333). The fact that these larval forms re-occur in the most different animals, cannot, I think, be sufficiently estimated, and bears plain witness to the former common descent of all from the *Gastræa*" (p. 345).

63 (i. 194). The Uniaxial Outline of the Gastrula is, on account of the two different poles of the axis, more accurately described as a diplopoleic uniaxial form (a sternometric outline: conoid-form, or cone). Cf. my "Promorphology" ("Generelle Morphologie," vol. i. p. 426).

64 (i. 194). Primitive Intestine and Primitive Mouth. My distinction of the primitive intestine and primitive mouth (*protogaster* and *protostoma*) from the later, permanent intestine and mouth (*metagaster* and *metastoma*) has been variously attacked; it is, however, as much justified as the distinction of the primitive kidney from the permanent kidney, of the primitive vertebrae from the permanent vertebrae. The primitive intestine

forms but a part of the permanent intestine, and the primi mouth (at least in the higher animals) does not become permanent mouth.

65 (i. 196). Primitive germ-layers (*blastophylla*). As two primary germ-layers (*entoderma* and *exoderma*) originate the sole histogenetic rudiment of the whole body, and the *mesoderma*, the nutritive yolk, and all other accessory parts of the germ have developed only secondarily from the former. I consider it very important to distinguish between the primary and secondary germ-layers. The latter, to distinguish them from the former, might be called "after germ-layers" (*telasma*).

66 (i. 201). Unequal Cleavage and Hood-gastrula (*mentatio inæqualis* et *Amphigastrula*). Next to Amphibia the most accessible examples for observation of unequal cleavage and the Amphigastrula are afforded by the indigenous bodied Animals (*Mollusca*) and Worms (Snails and Marine Earth Worms and Leeches).

67 (i. 202). The Colour of Amphibian-eggs is occasioned by the accumulation of dark colouring-matter at the animal pole of the egg. In consequence of this the animal-cells of the egg appear darker than the vegetative cells of the entoderm. In most animals the reverse is the case; the protoplasm of the entoderm cells being usually darker and more coarsely granular (vol. i. p. 197).

68 (i. 207). Hood-gastrula of Amphibia. Cf. Robert Ruedemann "On the Evolution of Batrachia" ("Ueber die Entwicklung der Batrachier," p. 126; Plate XII. Figs. 3-7). Stricker's "Manual of Tissues" ("Handbuch der Gewebelehre," vol. ii. p. 1202; Figs. 399-402). Goette, "History of the Evolution of the Bombinator" ("Entwicklungsgeschichte der Unke," p. 120; Plate II. Figs. 32-35).

69 (i. 214). Hood-gastrula of Mammals. Eduard Beneden, "La maturation de l'œuf, la fécondation et les premières phases du développement embryonnaire des Mammifères, d'après des recherches faites chez le lapin." Brussels, 1875. No figures.

given with these "Communication préliminaire;" Van Beneden's description is, however, so clear, so thorough and careful, that they afford an entirely satisfactory insight into unequal egg-cleavage and the formation of the Hood-gastrula in Mammals. Other observers, who have studied the germination of Mammalian eggs (among the most recent Kölliker, Rauber, and Newson may be especially mentioned), have overlooked or failed to recognize the important features discovered by Van Beneden.

70 (i. 218). The Disc-gastrula (*Disco-gastrula*) of Osseous Fishes (*Teleostei*). Van Bambeke, "Recherches sur l'embryologie des poissons osseux." Brussels, 1875. The transparent fish-eggs, in which I observed discoid cleavage (*Segmentatio discoidalis*) and the formation of the Disc-gastrula by invagination, are accurately described in my article on "The Gastrula and Egg-cleavage of Animals" ("Jen. Zeitschrift für Naturwissenschaft," 1875, vol. ix. p. 432-444; Plates IV., V.). On the disc-gastrula of *Selachii*, cf. Balfour, "The Development of the Placozo branch Fishes."—"Journ. of Anat. and Physiol.," vol. x. 517; Plates XX., XXIII.

71 (i. 221). Yelk-cells of Birds. The cell-like constituent parts, which occur in great number and variety in the nutritive yelk of Birds and Reptiles, as in most Fishes, are nothing less than true cells, as His and others have asserted. This does not mean that in this matter a distinct limit everywhere exists between the nutritive and the formative yelks, as in our oceanic fish-eggs (Figs. 42, 43, note 70). On the contrary, *originally* (phylogenetically) the nutritive yelk originated from part of the entoderm.

72 (i. 223). Egg-cells of Birds. Notwithstanding the large nutritive yelk, the "after-egg" (*metovum*) of Birds and Reptiles is, in form-value, a single cell. The very small, active protoplasm of the "tread" does, however, indeed fall far short, in volume, of the huge mass of the yellow yelk-ball. The bird's eggs are absolutely the largest cells of the animal body. Cf. note 43, and Eduard van Beneden, "Recherches sur la composition et la

signification de l'œuf." Brussels, 1870. Hubert Ludwig Egg-structure in the Animal Kingdom" ("Ueber die Eib in Thierreiche." Würzburg, 1874).

73 (i. 226). Discoidal cleavage (*Segmentatio discoida* Bird's eggs. Cf. Kölliker, "History of the Evolution and the Higher Animals" ("Entwicklungsgeschichte des sachen und der höheren Thiere." 2nd edition, 1876, pp. Figs. 16-22).

74 (i. 227). Disc-gastrula (*Disco-gastrula*) of Bird Rauber, "On the Place of the Chick in the System of tion" ("Ueber die Stellung des Hühnchens im Entwick plan"). Leipzig, 1876. Foster and Balfour, "The Elen Embryology." London, 1874.

75 (i. 231). Bladder-gastrula (*Perigastrula*) of Art Animals (*Arthropoda*). Cf. Bobretzky, "Russian Essay Germ-history of *Astacus* and *Palæmon*." Kiew, 1873. own article on the gastrula and egg-cleavage.—"Jen. Ze für Naturwissenschaft." Vol. ix. pp. 444-452, Plate VI.

76 (i. 234). The Four-layer Theory, which was first stated by Baer in 1837 ("Entwicklungsgeschichte der" vol. ii. pp. 46, 68), and which we have here carried out k yet appears the only form of the germ-layer theory, on comparative observation of all higher animals, su universal law of germination for all and at the same tim the inconsistent reputations of many observers.

77 (i. 239). Caspar Friedrich Wolff first indicated th layer Theory (note 76). Cf. the remarkable sentence, qu vol. i. p. 45, from his pregnant work on the formation intestinal canal (note 19).

78 (i. 240). The Four Main Types of Gastrulation are diagrammatically distinguished in Plates II. and III. Tables III. and IV. (vol. i. pp. 241, 242), are of course co by intermediate forms. These are transitions both betw primordial and the unequal forms, and between the primor the superficial forms; similarly, the unequal form of egg-c is connected by twixt-forms with the discoidal forms, whic

ain, perhaps, connected in the same way with the superficial

9 (i. 241). The Gastrulation of the various classes of animals has been far too little studied to enable us thoroughly summarize the distribution of the various forms within the separate classes. Yet it is already evident that primordial egg-cleavage and the formation of the Archigastrula occur in the same classes of each tribe.

10 (i. 243). The Rhythm of egg-cleavage is by no means as regular as might appear from the four first examples in the five series. There are, on the contrary, many variations, and not infrequently an entirely irregular and very variable sequence of cleavages occurs (especially in discoidal cleavage).

11 (i. 246). Definition of the Type. Cf. Gegenbaur, *Elements of Comparative Anatomy*, 1874, p. 59.

12 (i. 246). Types and Phyla. According to the prevailing "epitheory," the types of the animal kingdom are *parallel*, entirely independent; according to my "Gastræa-theory," on the contrary, they are divergent tribes, connected at the base; according to the view of Claus and other opponents, there is no essential distinction.

13 (i. 248). The one-celled condition of Infusoria entirely invalidates their morphological comparison with Metazoa. Cf. my article "On the Morphology of Infusoria" (*Jen. Zeitschrift Naturwissenschaft* 1873, vol. vii. p. 516-568).

14 (i. 257). The axes of the Vertebrate outline. Cf. my *Stereomorphology* (Stereometry of Organisms).—"Generelle Morphologie," vol. i. pp. 374-574. "Singly double-outlines" (*pleura*), p. 519. "Bilateral-symmetrical" forms in the fourth classification of the word.

15 (i. 255). The Primitive Vertebrate Type, as it is represented in Figs. 52-56, is a hypothetic diagram, which is principally based on the outline of the *Amphioxus*, but in which the comparative Anatomy of *Ascidia* and *Appendicularia* on the one side, of *Cyclostomi* and *Selachii* on the other, is regarded. The diagram is by no means meant to be an "exact figure," but

a provisional stage in the hypothetical reconstruction of the unknown, long extinct parent-form of Vertebrates, an "Amphioxus type."

86 (i. 258). Only very uncertain assumptions can be made as to the sense-organs of the hypothetical parent-form, for the organs, more than any others, have been subject to adaptation and in *Ascidia*, as in the *Amphioxus*, have probably been much atrophied. The earliest Vertebrates probably inherited a pair of eyes of very simple character and a pair of simple ear-vesicles from Worms.

87 (i. 267). The primitive kidneys were perhaps already metameric in the hypothetical parent-form of Vertebrates, so that in addition to the two longitudinal main canals (primitive kidney ducts) numerous transverse tubes (segmental canals) were connected with these main canals, a pair in each metamere of the middle part of the body. Perhaps these already opened through ciliated funnels into the body-cavity (*coeloma*), as is now the case in Annelids, and, according to Balfour, in the embryos of Selachii. Cf. Balfour, "Development of Elasmobranch Fishes."—"Quarterly Journal of Microscopical Science New Series, vol. xiv. p. 323; "Journal of Anat. and Physiol. vol. x.

88 (i. 273). The germination of Primitive Vertebrates. (with Table VI., Table VII. (vol. i. p. 327), Table XI. (p. 46) also the diagrammatic figures in Plates IV. and V. with explanation (p. 321).

89 (i. 276). The Germ-forms of the earliest Vertebrates they are represented in diagrammatic cross sections in Figs. 62-69, can only, of course, be approximately guessed, and with the aid of Comparative Anatomy and Ontogeny. The hypothetical diagrams, therefore, by no means claim to be accepted dogmatically, any more than do those in Figs. 52-59. (Cf. note 85.)

90 (i. 280). Main incidents in Vertebrate germination. The main palingenetic incidents here enumerated, perhaps sixth, ninth, and tenth originally occurred in a very

ent form. The other seven now appear to be pretty well established.

91 (i. 285). The flat germ-disc of Birds, which even now, in opinion of most embryologists, represents the first starting-point in the formation of the embryo, and to which all other germ-forms have been referred, is, on the contrary, a late and much modified germ-form, which has arisen in consequence of extension of the gastrula over the greatly enlarging nutritive k.

92 (i. 289). Site of Fertilization. In Man, as in other mammals, fertilization of the eggs probably usually takes place in the oviduct: here, the eggs which, at the rupture of the ovarian follicles, have emerged from the female ovary and passed into the outer opening of the oviduct, meet with the live sperm-cells of the male seed, which, during copulation, penetrated into the uterus, and from there passed into the inner lining of the oviduct. Rarely, fertilization occurs even on the ovary, or not till within the uterus. (Cf. Chapter XXV.)

93 (i. 293). The origin of the mesoderm in Mammals, as in other animals, is, at present, among the most obscure and concealed points of Ontogeny. Remak, Balfour, and others derive it from the entoderm, Kölliker and others from the exoderm. Hildeyer, His, and others assert that both primary germ-layers take part in the formation of the mesoderm. The last assumption is, I believe, correct. (Cf. notes 76, 77.)

94 (i. 297). The Germ-shield (*Notaspis*). The ordinary view, that the germ-shield (= Remak's "Doppelschild") is the earliest rudiment of the actual embryo, results in many erroneous conclusions. It is, therefore, necessary to point out especially that the germ-shield represents the first well-defined central axial part of the embryo.

95 (i. 317). Body Wall and Intestinal Wall. The morphological distinction between the body wall and the intestinal wall, certainly primordial, is probably referable to the simple primary germ-layers of the Gastræa. If the skin-fibrous layer is derived from the exoderm, and the intestinal-fibrous layer from the

entoderm, this most simply explains the progressive development of this distinction, which may be traced through the series Worms, and up to Vertebrates.

96 (i. 320). Palingenetic and Kenogenetic germination. In the germ-history of Vertebrates no clear conception of embryological process has yet been attained, because all authors have started from the higher Vertebrates (usually from Chick) and have assumed that the form of evolution occurring in this case is original and typical. It is only since the germ-history of the *Amphioxus* has taught us the palingenetic, the original form of germination of Vertebrate organisms, that we have been enabled, by Comparative Ontogeny (and especially the principles of the *Gastræa* theory), rightly to understand to explain phylogenetically the kenogenetic forms of germination of higher Vertebrates.

97 (i. 321). The Diagrams in Plates IV. and V. are as simple and abstract as possible, in order to render the desired explanation as easy as possible.

98 (i. 346). Primitive Vertebrae and Metamers. For a right conception of "primitive vertebral" structure it is especially necessary to point out that the primitive vertebrae are much more than their name indicates. They must, in fact, be conceived as individual, consecutive sections of the trunk, which have arisen one after the other, as true "metameric" consecutive pieces ("*Generelle Morphologie*," vol. i. p. 100). Each primitive vertebra of a Vertebrate, like each trunk-segment or metameron of an Annelid or Arthropod, contains an essential, morphological constituent parts, characteristic of the corresponding animal-tribe.

99 (i. 349). Origin of the Primitive Vertebrae. My conception of these as individual, morphological "consecutive pieces," which, like the metamers of Cestods and Annelids, have arisen by terminal budding from a single unarticulated piece, has been much attacked. I therefore emphatically remark that we must understand this process in the widest sense. In both cases it is certainly a reproduction of individual, like parts, which originated (in time and space) consecutively.

100 (i. 361). The agreement among the germ-forms of various Mammals is instructive especially because it shows us, by diversity in the mode of evolution, the most diverse structures can originate from one and the same form. As we usually see this in germ-forms, we may hypothetically assume the same to have occurred among tribe-forms. Moreover, this agreement is never absolute identity, but always only the very greatest similarity. Even the germs of the various individuals of a species are never actually identical.

101 (i. 366). The law of the ontogenetic connection of systematically allied animal-forms has many apparent exceptions. These are, however, fully explained by the adaptation of the germ to kenogenetic conditions of existence. Where the palinogenetic form of evolution of the germ has been accurately transmitted by heredity, that law is always in force. Cf. Fritz Müller, "Für Darwin" (note 111).

102 (i. 367). Earliest human germs. Cf. Kölliker, "History of the Evolution of Man" ("Entwicklungsgeschichte des Menschen." 2nd edition, 1876, pp. 303-319). Also Ecker, "Icones physiologicæ." Leipzig, 1859. Plates XXV.-XXXI. The earliest human germs which have yet been certainly recognized, were from twelve to fourteen days old, and were observed by Prof. Allen Thomson, of Glasgow. No opportunity has ever occurred for the observation of earlier germs.

103 (i. 369). Human germs of three weeks (twenty to twenty-five days) exhibit in their whole structure that phylogenetic stage of evolution which, among extant Vertebrates, is represented by the Cyclostomi (Lampreys and Hags, vol. ii. p. 103), and which must be referable to extinct Monorhine ancestors of similar structure.

104 (i. 370). Human germs of four weeks (twenty-five to thirty days), on the whole, exhibit in their whole structure that phylogenetic stage of evolution, which is exhibited in Sharks and Rays, among extant Vertebrates, and which is referable to similar extinct Primitive Fish ancestors (*Proselachii*). Of course this comparison is affected by various kenogenetic modifications

(both heterotopic and heterochronic), just as in the frog (Cf. note 108.)

105 (i. 374). The nose of Nosed-apes is much more different from that of other Apes than from that of Man. Moreover even the extreme variety and variability in the external form of the human nose shows how small is the morphological value of this organ, so important to the physiognomy.

106 (i. 383). The bladder-like form of the human Allantois. Cf. W. Krause, "On the Allantois in Man" ("Ueber die Allantois des Menschen."—"Archiv für Anat. u. Physiol.," 1875, p. 211 Plate VI.).

107 (i. 400). The navel-cord (*funiculus umbilicalis*), and the placenta, is an organ shared by Man exclusively with Placental Animals. Cf. Chap. XIX. pp. 155-168, and Figs. 200, 201. On the more minute structure of this organ, and on the special features of the embryonic blood-circulation, cf. Kölliker, "History of the Evolution of Man." 2nd edition, 1876, pp. 319-320.

108 (i. 401). The Kenogeny of Man. In pointing out the phylogenetic significance of the separate incidents and periods of human germ-history, and in explaining them by reference to corresponding processes and stages in the tribal history of our ancestors, we must always bear in mind that in Man, as in higher animals, the original palingenetic cause of germination has undergone much kenogenetic modification in consequence of many adaptations to the very various conditions of embryonic life, that it has thus been much violated and contracted. The higher the organism develops, the more are especially the earliest stages of evolution abbreviated.

109 (i. 404). The sections of human germ-history, of which only four larger and ten smaller are mentioned here in reference to their phylogenetic significance, allow of much more division if their comparative Ontogeny is minutely examined. Their phylogenetic significance may also be very well explained by a fitting reference to kenogenetic displacements in place and time (vol. i. p. 13).

110 (i. 405). Figures of human embryos in all stages

am-history were given in very beautiful detail by M. P. Erdl
ty years ago: "The Evolution of Man, and of the Chick in
Egg" ("Die Entwicklung des Menschen, und des Hühnchens
Ei." Leipzig, 1845).

111 (i. 409). Fritz Müller, "Für Darwin." Leipzig, 1864.
very excellent little book, in which the modification of the
fundamental law of Biogeny (with reference to the Phylogeny of
Insecta) are explained for the first time.

112 (i. 413). The Method of Phylogeny is of the same
Morphological value as the well-known method of Geology, and
may, therefore, claim exactly the same scientific acceptance.
See the excellent discourse by Eduard Strasburger, "On the
Importance of Phylogenetic Methods in the Study of Living
Organisms."—"Jenaische Zeitschrift für Naturwissenschaft," 1874,
vol. viii. p. 56.

113 (i. 415). Johannes Müller, "On the Structure and Vital
Phenomena of *Amphioxus lanceolatus*."—Transactions of the
Berlin Academy, 1844.

114 (i. 415). Recent works on the Amphioxus. W. Rolph
and E. Ray Lankester especially have recently added to our
knowledge of the organology of the Amphioxus, Wilhelm Müller
and P. Langerhans to that of its histology. The literature of
this subject is fully represented by W. Rolph, in his "Researches
on the Structure of the Amphioxus" ("Untersuchungen über
den Bau des Amphioxus."—"Morpholog. Jahrb.," vol. ii. p. 87,
plates V. and VII.), and in P. Langerhans, "On the Anatomy of
the Amphioxus" ("Zur Anatomie des Amphioxus."—"Archiv.
für Mikr. Anat.," vol. xii. p. 290, Plates XII.-XV.).

115 (i. 416). Acrania and Craniota. The separation of
Vertebrates into Skull-less Animals (*Acrania*) and Skulled
Animals (*Craniota*), which I first indicated in 1866 in my
"Generelle Morphologie," appears to me absolutely essential for
a phylogenetic explanation of the Vertebrate-tribe.

116 (i. 428). Max Schultze, "History of the Evolution of
the Petromyzon" ("Entwicklungsgeschichte von *Petromyzon*." Haar-
lem, 1856). The Ontogeny of the Hags, which promises very
important results, is yet, unfortunately, entirely unknown.

117 (i. 430). Savigny, "Mémoires sur les Animaux Vertébrés." Vol. ii., Ascidiæ, 1816. Giard, "Recherches sur les Synascidiæ."—"Archives de Zoologie Experimentale," 1872.

118 (i. 435). Syn-ascidia and Echinoderms. The Correlation of Echinoderms, which I explained in 1866 ("General Zoology," vol. ii. p. lxiii), and which has been much attacked, "paradoxical," is as yet the sole theory attempting the explanation of this remarkable group of animals.

119 (i. 442). Kowalevsky, "History of the Evolution of Amphioxus and of Simple Ascidiæ" ("Mémoires de l'Académie des Sciences de St. Petersburg." 7 Serie. Tom. x. and xi. 1867-8).

120 (i. 450). The metameric structure of the Amphioxus, which is indicated in its nerve and muscle systems, undoubtedly shows that the notochord exists in Vertebrates previous to the metameric structure, and consequently that it is inherited from unarticulated Chorda Animals.

121 (i. 454). The Metamorphosis of the Amphioxus, which the larva passes into the adult form, is not yet known in all its details. This does not, however, detract from the extraordinarily important bearing of the thoroughly established fact of the earliest incidents in its germination on the palingogenesis of the Vertebrates.

122 (i. 455). Fertilization of Ascidia (*Phallusia manubriata*). Eduard Strasburger, "On Cell-structure and Cell-division. Studies of Fertilization." 2nd edition. Jena, 1876, Plate VIII.

123 (i. 462). Kupffer. The tribal relation of Amphioxus to the Vertebrates ("Archiv für Mikros. Anat.," 1870, vol. i. pp. 115-170). Oscar Hertwig, "Researches into the Structural Evolution of the Cellulose Mantles of Tunicata" ("Monographien über den Bau und die Entwicklung des Tunicatens Mantels der Tunicaten"). Richard Hertwig, "Contributions to the Knowledge of Ascidian Structure" ("Beiträge zur Kenntnis des Baues der Ascidiæ."—"Jenaische Zeitschrift für Naturwissenschaft," 1873, vol. vii.).

124 (i. 464). The Phylogenetic Importance of the Amphioxus cannot be too highly insisted on. Without knowledge of Anatomy and Ontogeny, the origin of Vertebrates would be entirely dubious, and their descent from Worms would appear incredible.

125 (i. 467). The Ontogenetic Cell-pedigree, as it is represented, with reference to the Amphioxus, in Table XI., probably adds good, in its most important features, for all Vertebrates, and, therefore, also for Man. For, more than any other form, the Amphioxus by strict Heredity has accurately retained its lineage. This histogenetic cell-pedigree is apparently well established as regards most and the chief features; on the other hand, it yet appears doubtful with regard to the origin of the primitive kidneys, the testes, and ovaries.

126 (ii. 4). Milne-Edwards, "Leçons sur la Physiologie comparée," vol. ix.

127 (ii. 6). Eternity of Organic Life. According to the monistic view, organic life is a further form of evolution of the inorganic word-processes, and had a beginning in time on our planet. In opposition to this, A. Fechner, among others, in his "Thoughts on the Creation and Evolution of Organisms," has stated certain opposed "kosmorganic phantasys" which appear entirely irreconcilable with the ontogenetic facts given here.

128 (ii. 18). Bernhard Cotta ("Geologie der Gegenwart," 1866; 4th edition, 1874) and Karl Zittel ("Aus der Urzeit," München, 1875, 2nd edition) have made some excellent remarks on the duration and the whole course of the organic history of the world.

129 (ii. 21). August Schleicher, "The Darwinian Theory and Philology" ("Die Darwin'sche Theorie und die Sprachwissenschaft." Weimar, 1863. 2nd edition, 1873).

130 (ii. 25). At first sight, most polyphyletic hypotheses appear more simple and easy than do monophyletic, but the former always present more difficulties the more they are considered.

131 (ii. 25). Those physiologists who desire an experi-

mental proof of the theory of descent, merely thereby proving extraordinary ignorance of the morphological scientific facts relating to this matter.

132 (ii. 30). Spontaneous generation.—“Generelle Morphologie,” vol. i. pp. 167–190. “Monera and Spontaneous generation.”—“Jenaische Zeitschrift für Naturwissenschaft,” vol. vi. pp. 37–42.

133 (ii. 33). The Absence of Organs in Monera. In that Monera are “organisms without organs,” we understand the definition of organs in a morphological sense. In a physiological sense, on the other hand, we may call the variable physiological processes of the body of the Moneron the “*pseudopodia*” of the organism.

134 (ii. 36). Induction and Deduction in Anthropology.—“Generelle Morphologie,” vol. i. pp. 79–88; vol. ii. p. 357. “History of Creation,” vol. ii. p. 357.

135 (ii. 42). Animal Ancestors of Man. The number of species (or, more accurately, form-stages, which are distinguished as “species”) must, in the human ancestral line (in the course of many millions of years!), have amounted to many thousands; the number of genera to many hundreds.

136 (ii. 47). Following Elsberg, we give the name of “plastidules” to the “molecules of plaston,” to the smallest like units of that albuminous substance which, according to the “physical theory,” is the material substratum of all the active phenomena of life. Cf. my work on “The Perigenesis of Plastidules” (“Perigenesis der Plastidule oder Wellenzeugung der Plastidule.” Berlin, 1876). This is an attempt to explain mechanically the elementary processes of evolution.

137 (ii. 49). Bathybius and the free protoplasm of the deep. Cf. my “Studies on Monera and other Protoplasmic Organisms,” Leipzig, 1870, p. 86. The most recent observations on Bathybius are those of Dr. Emil Bessel, who found this form on the coast of Greenland (in Smith’s Sound), at a depth of 550 ft. He noticed very active amœboid movements in the organism as well as the assumption of foreign particles (carminum). “It consists of nearly pure protoplasm, tinged most inter-

solution of carmine in ammonia. It contains fine gray granules of considerable refracting power, and besides the latter a great number of oleaginous drops, soluble in ether. It manifests very marked amœboid motions, and takes up particles of carmine, etc." Packard, "Life Histories of Animals, including Man." New York, 1876.

138 (ii. 50). The Philosophical Importance of Monera in explaining the most obscure biological questions cannot be sufficiently emphasized. Monograph of Monera.—"Jenaische Zeitschrift für Naturwissenschaft," vol. iv., 1868, p. 64.

139 (ii. 54). The Nature and Significance of the Egg-cell can only be philosophically understood by means of phylogenetic elimination.

140 (ii. 58). Synamœba. Cienkowski, "On the Structure and Evolution of Labyrinthula" ("Über den Bau und die Entwicklung der Labyrinthuleen."—Arch. für Mikrosk. Anat., 1870, vol. iii. p. 274). Hertwig, "Microgromia Socialis."—*Ibid.*

141 (ii. 61). Catallacta, a new Protista-group (*Magosphæra rutila*). See "Jenaische Zeitschrift für Naturwissenschaft," vol. vi., 1871, p. 1.

142 (ii. 66). Haliphysema and Gastrophysema. Extant starfishes. See "Jenaische Zeitschrift für Naturwissenschaft," vol. xi., 1876, p. 1, Plates I.–VI.

143 (ii. 70). The five first stages in the evolution of the animal body, which are compared in Table XVII., and which are common to Man and all higher Animals, are established beyond all doubt as existing in the Ontogeny of most extant animals. As Comparative Anatomy shows that corresponding stages yet exist in the system of the lower animals, we may assume, in accordance with the fundamental law of Biogeny, that similar forms existed phylogenetically as most important ancestral forms.

144 (ii. 77). On the distinction of the axes, and on the symmetric outline of the animal body, see "*Promorphologie*" "*Generelle Morphologie*," vol. i. pp. 374–574).

145 (ii. 87). The hermaphrodite structure of our ancestral

series was perhaps transmitted from the Chorda Animals even far as the lower stages of Vertebrate ancestors. Cf. Chapter X

146 (ii. 89). I am inclined to regard the Appendicular living Chorda Animals of the present day; they are the Invertebrates permanently possessing a notochord, and thus by many other peculiarities, distinguished from genuine Tates.

147 (ii. 105). Metamorphosis of Lampreys. That the Ammocoetes change into Petromyzon was known two hundred years ago (1666) to the fisherman Leonhard Baldner of Samburg; but this observation remained unrecognized, and modification was first discovered by August Müller in ("Archiv für Anat.," 1856, p. 325). Cf. Siebold, "The Freshwater Fishes of Central Europe" ("Die Süßwasserfische Mittel-Europa," 1863).

148 (ii. 114). Selachii as Primitive Fishes. The old disputes as to the systematic position and kindred of Selachii were definitely settled by Gegenbaur, in the introduction to his work on "The Head-skeleton of Selachii."

149 (ii. 118). Gerard Krefft, "Description of a Gigantic phibian;" and Albert Günther, "Ceratodus, and its Systematic Position."—"Archiv für Naturgeschichte," 37, 1871, vol. 321; also "Phil. Trans.," 1871, Part II. p. 511, etc.

150 (ii. 129). The duration of metamorphosis of Amphibia varies much in the different forms of Frogs and Toads, the former forming a complete phylogenetic series from the original, complete form, to the later, much shortened and vitiated form of modification.

151 (ii. 129). "All the histological features of the Salamander (*Salamandra maculata*) force the impression that it belongs to an entirely different epoch of terrestrial life than that of the Water Salamander (*Triton*), externally so similar."—J. Remak ("Entwicklung der Wirbelthiere," p. 117).

152 (ii. 130). Siredon and Amblystoma. Very various views have lately been expressed as to the phylogenetic significance to be attributed to the much-discussed modification of the M

into an Amblystoma. Cf. on this subject especially Weismann, in "Zeitsch. für wissensch. Zoologie," vol. 1, pp. 297-334.

(ii. 131). The Leaf-frog of Martinique (*Hylodes martinicensis*) loses its gills on the seventh day, its tail and yolk-sac on the eighth day of egg-life. On the ninth or tenth day after hatching the complete frog emerges from the egg.—Béarn, "Hylodes Martinicensis et ses Métamorphoses." "Journal. de l'Académie des Sciences de Paris," par Grevais, vol. ii. 1873, p. 13.

(ii. 133). "Homo diluvii testis" = Andreas Schenckner, the bone of an ancient evil-doer; Soften, stone, the heart of the children of evil" (Dionysius Miller). "Quarantennary and Now" ("Sonst und Jetzt," 1856, p. 239).

(ii. 133). The Amnion-structure of the three highest vertebrate-classes, wanting in all lower Vertebrates, has no connection with the similar, but independently acquired Amnion-structure (analogous, but not homologous) of higher Arthropods.

(ii. 138). The former existence of a Protamnion, the parent-form of all Amniota, is undoubtedly shown by comparative Anatomy and Ontogeny of Reptiles, Birds, and Mammals. No fossil remains of such a Protamnion have, however, been discovered. They must be sought in the Permian carboniferous formation.

(ii. 147). The former organization of the Protamnion is hypothetically reconstructed from the comparative anatomy of the Salamander, Lizards, and Embryos (*Amphioxus*).

(ii. 153). The Ectodermic structure of the Protamnion is very different from all known Fossil Animals (*Amphioxus*), but possessed all the essential internal characters of the latter.

(ii. 163). The phylogeny of the Protamnion, as the ancestral placental parent-group, is not influenced by any fossil Protamnion, but it is most able to determine the phylogeny of the latter.

160 (ii. 168). On the structure of the Decidua very various theories have been given. Cf. Kölliker, "History of the Evolution of Man" ("Entwicklungsgeschichte des Menschen." 2nd edition, 1871, pp. 319-376). Ercolani (Giambattista), "Sul processo formativo della placenta." Bologna, 1870. "Le glandole otricolari del'utero." Bologna, 1868, 1873. Huxley, "Lectures on the Elements of Comparative Anatomy," 1864, pp. 101-112.

161 (ii. 172). Huxley, "Anatomy of Vertebrates," 1873, p. 382. Previously Huxley had separated the "Primates" into seven families of nearly equal systematic value. (See "Man's Place," etc., p. 119.)

162 (ii. 179). Darwin. Sexual selection in Apes and Man.—"Descent of Man," vol. ii. pp. 210-355.

163 (ii. 180). Man-like Holy Apes. Of all Apes, some Holy Apes (*Semnopithecus*) most resemble Man, in the form of their nose and the character of their hair (both that on the head and that on the beard).—Darwin, "Descent of Man," vol. i. p. 335; vol. ii. p. 172.

164 (ii. 182). Friedrich Müller ("Allgemeine Ethnographie." Vienna, 1873, p. 29), on the supposed age of man. Families of languages (pp. 5, 15, etc.).

165 (ii. 183). The plate (XV.) representing the migrations, given in the "History of Creation," merely claims the value of a first attempt, is an hypothetic sketch, as I there expressly said, and as, in consequence of repeated attacks, I must here insist.

166 (ii. 201). The Leather-plate. The phylogenetic distinction of a special leather-plate, the outermost lamella separating from the skin-fibrous layer, is justified by Comparative Anatomy.

167 (ii. 204). Milk-glands. Huss, "Contributions to the History of the Evolution of the Milk-glands" ("Beiträge zur Entwicklungsgeschichte der Milchdrüsen"); and Gegenbaur, "On the Milk-gland Papillæ" ("Jenaische Zeitschrift für Naturwissenschaft," 1873, vol. vii. pp. 176, 204).

168 (ii. 208). On the hairy covering of Man and Apes, see Darwin, "Descent of Man," vol. i. pp. 20, 167, 180; vol. ii. pp. 280, 298, 335, etc.

169 (ii. 217). Dorsal side and ventral sides are homologous in Vertebrates, Articulated Animals (*Arthropoda*), Soft-bodied Animals (*Mollusca*), and Worms, so that the dorsal marrow and the ventral marrow are not comparable. Cf. Gegenbaur, "Morph. Jahrbuch," vol. i. pp. 5, 6.

170 (ii. 228). The unknown ontogenetic origin of the sympathetic nerve-system must probably, for phylogenetic reasons, be sought chiefly in the intestinal layer, *not* in the skin-layer.

171 (ii. 248). On the cavities connected with the nose, see Gegenbaur, "Elements of Comparative Anatomy," p. 580.

172 (ii. 260). The analogies in the germination of the higher sense organs were rightly grasped even by the earlier natural philosophers. The first more accurate sketches of the very obscure germ-history of the sense-organs, especially of the eye and ear, were given (1830) by Emil Huschke, of Jena (*Isis*, *Meckel's Archiv*, etc.).

173 (ii. 265). Hasse, "Anatomical Studies" ("Anatomische Studien"), chiefly of the organ of hearing. Leipzig, 1873.

174 (ii. 269). Johannes Rathke, "On the Gill-apparatus and the Tongue-bone" ("Ueber den Kiemen-apparat und des Zungenbein," 1832). Gegenbaur, "On the Head-skeleton of *Selachii*," 1872. (See note 124.)

175 (ii. 272). On the Rudimentary Ear-shell of Man, cf. Darwin, "Descent of Man," vol. i. pp. 17-19.

176 (ii. 276). Scarcely anywhere does Comparative Anatomy prove its high morphological value as with reference to the skeleton of Vertebrates: in this matter it accomplishes much more than Ontogeny. There is all the more reason to insist on this here, as Goette, in his gigantic history of the evolution of Bombinator, has recently denied all scientific value to Comparative Anatomy, and asserted that Morphology is explained solely by Ontogeny. Cf. my "Aims and Methods of the Recent History of Evolution" ("Ziele und Wege der heutigen Entwicklungsgeschichte," 1875, p. 52, etc.).

177 (ii. 283). The Human Tail, like all other rudimentary organs, is very variable in point of size and development. In

rare cases it remains permanently, projecting freely: usual disappears at an early period, as in Anthropoid Apes.

178 (ii. 284). On the Number of Vertebrae in different Mals, cf. Cuvier, "Leçons d'Anatomie Comparée." 2nd ed. tome i., 1835, p. 177.

179 (ii. 293). On the earlier Skull-theory of Goethe and O cf. Virchow, "Goethe as a Naturalist" ("Goethe als Naturforscher," 1861, p. 103).

180 (ii. 295.). Karl Gegenbaur, "The Head-skeleton of Selachii" ("Das Kopfskelet der Selachier"). As the foundation of a study of the head-skeleton of Vertebrates (1872).

181 (ii. 301). Karl Gegenbaur, "On the Archipterygi"—"Jenaische Zeitschrift für Naturwissenschaft," vol. vii. p. 131.

182 (ii. 304). Gegenbaur, "Researches into the Comparative Anatomy of Vertebrates" ("Untersuchungen zur Vergleichenden Anatomie der Wirbelthiere"). Part I. Carpus and Tibia (1864). Part II. The shoulder girdle of Vertebrates. Pectoral fins of Fishes (1866).

183 (ii. 305). Charles Martins, "Nouvelle comparaison des membres pelviens et thoraciques chez l'homme et chez les mammifères."—"Memoires de l'Acad. de Montpellier," vol. vi. 1857.

184 (ii. 308). Ossification. Not all bones of the human body are first formed of cartilage. Cf. Gegenbaur, "On Primary and Secondary Bone-formation, with special reference to the modern Skull Theory."—"Jenaisch. Zeitschrift für Naturwissenschaft," 1867, vol. iii. p. 54.

185 (ii. 308). Johannes Müller, "Comparative Anatomy of Myxinoidea."—"Transactions of the Berlin Academy," 1834.

186 (ii. 314) The Homology of the Primitive Intestine. The two primary germ-layers is the postulate for morphological comparison of the various Metazoa-tribes.

187 (ii. 322). In the Evolution of the Intestine, Amphibians and Ganoids have, by heredity, retained the original Craniotome more accurately than have Selachii and Osseous Fishes (Teleostei).

palaeogenetic germination of *Selachii* has been much altered by ontogenetic adaptations.

88 (ii. 323). On the Homology of Scales and Teeth, cf. Gegenbaur, "Comparative Anatomy" ("Grundriss der vergleichenden Anatomie," 1874, pp. 426, 582); also Oscar Hertwig, "Jenaische Zeitschrift für Naturwissenschaft," 1874, vol. viii. On the important distinction of homology (morphological resemblance) and Analogy (physiological resemblance), see Gegenbaur, as above, p. 63; also my "Generelle Morphologie," vol. i. p. 313.

89 (ii. 337). Wilhelm Müller, "On the Hypobranchial Valve in Tunicates, and its Presence in the Amphioxus and Ostomi."—"Jenaische Zeitschrift für Naturwissenschaft," vol. viii. p. 327.

90 (ii. 358). The Nerve-muscular Cells of the Hydra throw earliest light on the simultaneous, phylogenetic differentiation of nerve and muscle tissue. Cf. "Klemenberg, Hydra." Leipzig,

91 (ii. 383). The germ-history of the human heart accurately reproduces in all essential points its tribal history. This palaeontological reproduction is, however, much contracted in particular details and vitiated by kenogenetic modifications of the original line of evolution, displacements partly in time, partly in place, which are the result of embryonic adaptations.

92 (ii. 383). On the Special Germ-history of the Human Nervous system, cf. Kölliker, "History of the Evolution of Man" ("Entwicklungsgeschichte des Menschen." 2nd edition, 1876); Rathke's excellent work on Ontogeny.

93 (ii. 387). The Homologies of the Primitive Organs, as they are here provisionally described in accordance with the recapitulation-theory (note 24), can only be established by further co-ordination between Comparative Anatomy and Ontogeny. Cf. Gegenbaur on Comparative Anatomy ("Grundriss der vergleichenden Anatomie").

94 (ii. 390). The Mechanism of Reproduction. As the conditions of reproduction and of heredity, connected with reproduction, are referable to growth, so the former as well as the

latter are finally explicable as the results of the attraction and rejection of homogeneous and heterogeneous particles.

195 (ii. 397). Eduard van Beneden, "De la Distinction originelle du Testicule et de l'Ovaire." Brussels, 1874.

196 (ii. 399). On the Original Hermaphrodite Structure of Vertebrates, cf. Waldeyer, "Ovary and Egg" ("Eierstock und Ei," 1872, p. 152); also Gegenbaur ("Grundriss der vergleichenden Anatomie," 1874, p. 615). On the origin of the eggs from the ovary-epithelium, cf. Pflüger, "On the Ovaries of Mammals and Man" ("Die Eierstöcke der Säugethiere und des Menschen," 1863).

197 (ii. 423). On the special germ-history of the urinary and sexual organs, cf. Kölliker, "History of the Evolution of Man." On the homologies of these organs, see Gegenbaur ("Grundriss der vergleichenden Anatomie," 1874, pp. 610-628).

198 (ii. 449). Wilhelm Wundt, "Lectures on the Human and Animal Mind" ("Vorlesungen über die Menschen- und Thierseele." 1863). W. Wundt, "Outlines of Physiological Psychology" ("Grundzüge der Physiologischen Psychologie," 1874).

199 (ii. 457). On Active (actual) and Latent (preterital) forces, cf. Hermann Helmholtz, "Interoperation of Natural Forces" ("Wechselwirkung der Naturkräfte," Part II., 1871).

200 (ii. 457). "Anthropology as Part of Zoology."—"Generelle Morphologie," vol. ii. p. 432. "History of Creation," vol. i. 7; vol. ii. 347.

INDEX.

A

- ax*, ii. 73, 92
i, ii. 75, 92
forms, ii. 86
i, i. 116; ii. 97
apple, ii. 336
ion, i. 158
arth, i. 400
 thoughts on creation, i. 116
(trachea), ii. 330, 333
 182
s, i. 380; ii. 135, 411
 period, ii. 12
 ii. 146, 204
 i. 314, 386
animals, ii. 120, 133
 sheaths of, i. 387
 water, i. 314
 ii. 120, 133
 i. 142; ii. 152
 false feet of, i. 142
 d egg-cells, i. 144; ii. 53
 movements, i. 142; ii. 53
 states, ii. 56
a, ii. 120, 122
astrula, i. 200, 241
nia, i. 160
cus, i. 413; ii. 98
 blastula of, i. 443
 body-form of, i. 417
 cells of, their pedigree,
 i. 467
 chorda of, i. 417
 distribution of, i. 415
 gastrula of, i. 444
 germ-layers of, i. 447
Amphioxus, medullary tube of, i. 418
 place of, in natural
 system, i. 416
 sexual organs of, i. 425
 side canals of, i. 423
 significance of, i. 254,
 427
Amphirhina, ii. 97, 101
Analogy, ii. 412
Anamnia, ii. 97, 120
Ancestral series of man, ii. 44, 184
Animalculists, i. 37
Animal germ-layer, i. 194, 327
 organs, ii. 192, 194
Anorgana, i. 156; ii. 30
Anthropocentric conception, ii. 457
Anthropoids, ii. 177, 189
Anthropolithic epoch, ii. 11, 16
Anthropozoic periods, ii. 12, 17
Antimera, i. 257
Anus, i. 339; ii. 323, 345
Anus-groove, i. 339
Auvil (Incus of ear), ii. 261, 268
Ape-men, ii. 44, 181
Apes, ii. 165, 189
 eastern, ii. 172, 189
 flat-nosed, ii. 172, 189
 narrow-nosed, ii. 172, 189
 question as to descent of, ii.
 165, 441
 tailed, ii. 172, 189
 western, ii. 172, 189
Aorta, i. 265; ii. 378
 roots of, ii. 375
 stem of, ii. 375
Aortal arches, ii. 375, 378
Appendicularia, i. 459; ii. 90

Archelminthes, ii. 76
Archigastrea, i. 198, 241
 Archilithic epoch, ii. 9, 19
Archipterygium, ii. 301
 Archizoic periods, ii. 9, 11
Area generativa, i. 292
 — *opaca*, i. 296
 — *pellucida*, i. 296
 Aristotle, i. 27; ii. 368
 — epigenesis, i. 29
 — heart formation, ii. 368
 — his history of evolution, i. 27
 Arm, lower, ii. 278, 304
 — upper, ii. 278, 304
Arteria omphalo-mesenterica, i. 395
 — *umbilicales*, i. 400
 — *vertebrales*, i. 395
 — *vitellinae*, i. 395
 Arteries, i. 393
 Artery-arches, ii. 377
 — stalk, ii. 380
Arthropoda, ii. 92, 94
 Articulated animals, ii. 92, 94
 Articulation in man, i. 346
Ascidia, i. 429; ii. 90
 — *blastula* of, i. 455
 — *chorda* of, i. 456
 — communities of, i. 455
 — *gastrula* of, i. 455
 — gill-sac of, i. 431
 — heart of, i. 433
 — homologies of, i. 465, 466
 — intestine of, i. 432
 — mantle of, i. 430, 461
 — medullary tube of, i. 458
 — sexual organs of, i. 434
 — tail of, i. 456
Ascula, ii. 68
Atrium, ii. 374, 381
 Auditory nerve, ii. 262
 — organ, ii. 260
 — passage, ii. 269
 — vesicles, ii. 262
 Auricular processes of heart, ii. 381
 Axes of the body, i. 255; ii. 77
 Axial cord, i. 301
 — rod (notochord), i. 302
 — skeleton, ii. 280, 299
 Axis-plate, i. 299
Axolotl, ii. 126

B

BAER, KARL ERNST, i. 50
 — his germ-layer theory, i. 51
 — his law, i. 58
 — life of, i. 52
 — on the bladder-like outgrowth, ii. 62
 — on the human egg, i. 55; 424
 — on the notochord, i. 55
 — on type theory, i. 54
Balanoglossus, ii. 85
 Bathybius, ii. 49
Batrachia, ii. 131
 Bats, ii. 169, 187
 Beaked animals, ii. 147, 187
 Bell-gastrula, i. 198
 Bilateral outline, i. 257; ii. 74
Bimana, ii. 169
 Biogeny, i. 24; ii. 434
 — fundamental law of, 24; ii. 434
 Birds, ii. 120, 138
 — *gastrula* of, i. 223
 Bischoff, Wilhelm, i. 59
 Bladder-gastrula, i. 229, 241
Blastæa, ii. 61
Blastocœloma, i. 189
Blastodermis, i. 189
Blastodiscus, i. 227
 Blastogeny, i. 24
Blastophylla, i. 195
 Blastophyly, i. 24
Blastosphæra, i. 191
Blastula, i. 191, 242
 Blind-intestine, ii. 330, 343
 Blood-cells (corpuscles), i. 15; 366
 — relationship, i. 112
 — vessels, ii. 370
 Bloodless worms, ii. 75
 Bonnet, i. 40
 Brain, i. 212, 232
 — bladders of, i. 343; ii.
 — parts of, ii. 212
 — skull of, ii. 292
 Breast-body, ii. 282
 — bone, ii. 282
 — cavity, i. 261
 — *vertebrae*, ii. 282

gemmation), ii. 391
seriosus, ii. 374
ali, ii. 250

C

- to epoch, ii. 11, 15
 period, ii. 15, 19
 ii. 278, 304
 period, ii. 9, 19
aricularis, ii. 381
 rous period, ii. 10, 19
reins, i. 391
 . 278
 , ii. 176, 189
 hes, theory of, i. 76
cientes, i. 16, 80; ii. 455
ules, i. 16, 80; ii. 455
mpani, ii. 261, 270
on, i. 124
el (nucleus), i. 125
s, i. 124
 tance, i. 125
 25
le, i. 171; ii. 392
s, i. 171; ii. 392
 ry of, i. 60, 121
 eart, ii. 120
edulla, ii. 210, 232
 erve-system, ii. 210
keleton, ii. 280, 299
 of sight," ii. 252
Fosteri, ii. 119
n, ii. 212, 232
 , ii. 212, 232
 i. 187
ha, ii. 160, 187
iod, ii. 14, 19
mges, i. 117
 portance of, i. 31
ee, ii. 178, 180
imals, ii. 84, 87
rsalis, i. 255, 301
death, ii. 286
ssue of, ii. 286
tebralis, i. 255, 301
s, i. 84, 87
a, ii. 252, 258
i, 387; ii. 158
frondosum, ii. 160
lave, ii. 160
- Chorion, smooth, ii. 160
 ——— tufted, ii. 160
 Chorology, i. 113
 Chyle vessels, ii. 374
Cicatricula, i. 138
 Circulation in *Amphioxus*, i. 423
 ——— *Ascidia*, i. 433
 ——— Fishes, ii. 375
 ——— germ-area, i. 397
 ——— Mammals, ii. 378
 Clavicula, ii. 278, 304
 Cleavage cells, i. 185
 ——— forms of, i. 242
 ——— of egg, i. 185, 241
 ——— partial, of bird's egg, i. 224
 ——— rhythm, i. 243
 ——— superficial, i. 200, 241
 ——— unequal, i. 200, 241
 Clitoris, ii. 423, 431
 Cloaca, ii. 145, 418
 Cloacal animals, ii. 145, 187
 Coalescence, i. 164
 Coal period, ii. 11, 19
Coccyx, ii. 282
Cochlea, ii. 263, 268
Cæliterata, ii. 73
Cæloma, i. 260; ii. 75
Cælomati, ii. 75, 92
Columna vertebralis, i. 349; ii. 285
 Comparative Anatomy, i. 107, 245
 Concrescence, i. 164
Conjunctiva, ii. 259
 Connective membrane of eye, ii. 251
 ——— tissue, ii. 363
Connectivum, ii. 361, 366
 Convolutions of brain, ii. 226
 Copulation organs, ii. 421
Copulativa, ii. 421
Coracoideum, ii. 278, 304
Corium, ii. 200, 232
 Cormogeny, i. 24
 Cormophyly, i. 24
Cornea, ii. 251, 258
Costæ, ii. 278, 282
 Covering tissue, ii. 361
Craniota, ii. 100, 120
Cranium, ii. 291
 Creation, i. 74, 79; ii. 183
 Crooked intestine, ii. 319, 330
 Cross-vertebræ, ii. 282
 Crystalline lens, ii. 253, 258
 Culture period, ii. 11

Curves of embryo, i. 369
Cutis, ii. 200, 232
 Cuvier, theory of catastrophes, i. 76
 ——— theory of types, i. 57
Cyclostoma, ii. 101, 120
Cytods, i. 103
Cytula, i. 176
Cytococcus, i. 176

D

DALTON, i. 51
 Darwin, Charles, i. 96
 ——— descent of man, i. 103
 ——— selection, theory of, i. 95
 ——— sexual selection, i. 103
 ——— Erasmus, i. 96
 Darwinism, i. 95
Decidua, ii. 161, 165
 ——— animals, ii. 161
Decidua-less animals, ii. 161
Deciduata, ii. 161, 187
 Deduction, i. 104; ii. 37
 Degree of development, i. 58
Derma, ii. 232
 Descent, theory of, i. 84
 ——— of man, i. 104
 Devonian period, ii. 10, 19
Didelphia, ii. 149, 187
 Differentiation, i. 152, 159
 Digestive intestine, ii. 330
 Digits, ii. 278
 Diluvial period, ii. 12, 15
Dipneusta, ii. 115, 120
Discognathula, i. 219, 241
Discoidal cleavage, i. 225, 242
Discoplacentalia, ii. 162, 187
Discus blastodermicus, i. 139, 226
 Doellinger, i. 50
 Dorsal furrow, i. 302
 ——— marrow, ii. 221
 ——— swellings, i. 303
 Double-breathers, ii. 117, 120
 ——— nostrils, ii. 97, 101
 "Double-shield," i. 297
 Dualism, i. 17; ii. 456
 Dualistic philosophy, i. 17
Ductus Gartneri, ii. 416, 431
 ——— *Mülleri*, ii. 415, 431
 ——— *Rathkei*, ii. 415, 431
 ——— *Wolffi*, ii. 415, 431
 Dysteleology, i. 109

E

EAR, BONELETS OF, ii. 268
 ——— labyrinth of, ii. 262, 263
 ——— muscles of, ii. 271
 ——— nerve of, ii. 266
 ——— pouch (*utricle*), ii. 262
 ——— sac (*sacculus*), ii. 262
 ——— shell of, ii. 269
 ——— snail of, ii. 263, 268
 ——— trumpet, ii. 260
 ——— vesicles, ii. 265
 ——— wax, glands of, ii. 263
Echidna, ii. 147
Echinoderm, i. 435; ii. 92
 Egg-cell, i. 132
 ——— of Chick, i. 139
 ——— Bird, i. 139
 ——— Mammal, i. 137
 ——— Man, i. 137; ii. 425
 ——— sponge, i. 144
 ——— cleavage, i. 185, 242
 ——— holoblastic, i. 215
 ——— human, i. 137; ii. 425
 ——— membranes, i. 375; ii. 158
 ——— meroblastic, i. 216
 Elbow, ii. 278, 304
 Elementary organism, i. 124
 Embryo of Vertebrates, i. 360
 Embryology, i. 3
 Empty intestine, ii. 319
Encephalon, ii. 232
Endocoelarium, ii. 366, 400
Enteropneusta, ii. 86
Entoderma, i. 206, 236
 Eocene period, ii. 11, 15
Epidermis, ii. 200, 232
Epididymis, ii. 417, 428
Epigenesis, i. 39, 41
Epigenesis, theory of, i. 39, 41
 Epithelial tissue, ii. 361
Epithelium, ii. 361, 366
 Epochs, duration of, ii. 3
 Evolution of forms, i. 19
 ——— of functions, i. 19
 ——— history of, i. 1, 24
 ——— theory of, i. 34
 Excretory organs, i. 267; ii. 403
Exocoelarium, ii. 369, 400
Exoderma, i. 195, 236

ies, ii. 111, 306
 50, 258
 nective membrane of, ii.
 1
 , ii. 259
 ed membrane of, ii. 252,
 58
 ective membrane of, ii. 251
 il of, ii. 250
 bow membrane of, ii. 252
 ular membrane of, ii. 252
 cles, i. 357; ii. 253

F

S AB AQUAPENDENTE, i. 31
 elopment of, ii. 245, 346
 all of, ii. 278, 294
 canals, ii. 431
 hydatids, ii. 431
 er of corium, ii. 232
 reast, ii. 202
 ells, i. 171, 392
 ppulatory organs, ii. 423
 cretory ducts, ii. 415, 431
 erm-glands, ii. 398
 erm-layer, ii. 398
 ilk-glands, ii. 202
 hallus (*Clitoris*), ii. 423
 xual organs, ii. 423
 xual plate, ii. 401
 terus, ii. 417
 . 278, 304
 ion, i. 169, 176
 . 278, 304
 ral rod of, ii. 302
 of, ii. 302
 eton of, ii. 302
 ses, i. 16
 ii. 278
 . 109, 120
 as of, ii. 111
 astrula of, i. 219
 ales of, ii. 331
 ed foot, ii. 123
 ns, ii. 76
 259
 er, i. 236
 70
 l matter, ii. 456

Forces, active, ii. 457
 ——— latent, ii. 457
 Formative functions, i. 15
 ——— yolk, i. 216
 Forms, science of, i. 20
 Frog-Batrachia, ii. 131
 Frogs, ii. 131
 ——— egg-cleavage of, i. 203
 ——— gastrula of, i. 207
 ——— larva of, ii. 127
 ——— metamorphosis of, ii. 126
 Frontal process, ii. 244
 Functions of evolution, ii. 155
 ——— science of, i. 19
Funiculus genitalis, ii. 418
 ——— *umbilicalis*, i. 383; ii. 168

G

GALL-BLADDER, ii. 341
 ——— ducts, ii. 341
 ——— intestine, ii. 317, 330
 Ganoid Fishes, ii. 112, 120
 Gartnerian duct, ii. 416, 431
Gastræa, i. 232; ii. 66
 ——— theory of, i. 247; ii. 195
Gastræads, ii. 62, 70
Gastrocystis, i. 291
Gastrodiscus, i. 292
Gastrula, i. 192; ii. 65
 ——— Bell-, i. 198
 ——— Bladder-, i. 200
 ——— Disc-, i. 200
 ——— Hood-, i. 200
 Gegenbaur, i. 108; ii. 96
 ——— on Comparative Ana-
 tomy, ii. 96
 Gegenbaur on head-skeleton, ii. 293
 ——— on theory of descent,
 i. 108
 ——— skull theory, ii. 293
 ——— theory of limbs, ii. 299
Generatio spontanea, ii. 30
 "Generelle Morphologie," i. 102
 Geological hypotheses, i. 410
 Germ, i. 3
 Germ-area, i. 292
 ——— dark, i. 297
 ——— light, i. 297
 ——— cavity, i. 189
 ——— disc, i. 139, 226

Germ-epithelium, ii. 401
 — glands, ii. 398
 — history, i. 6, 24
 — layer, middle, i. 13
 — membrane, i. 189
 — membrane vesicle, i. 189
 — plate, ii. 401
 — point, i. 135
 — shield, i. 297
 — spot, i. 135
 — vesicle, i. 179, 291
 Gibbon, ii. 178, 181
 Glacial period, ii. 11
 Glands of intestine, ii. 320
 — akin, i. 201
Glans phalli, ii. 422
Glomeruli renales, ii. 407
Gnathostomi, ii. 109
 Goethe, Wolfgang, i. 88
 — his skull theory, ii. 293
 — morphology, i. 88
 — on metamorphosis, i. 90
 — on reason, ii. 453
 — on specification, i. 90
 Goette, Alexander, i. 65
Gonades, ii. 398
Gonochorismus, ii. 69, 395
Gonophori, ii. 402
 Gorilla, i. 178, 180
 Graafian follicles, ii. 424
Gubernaculum Hunteri, ii. 431

H

HAIR, ii. 205, 232
 Hair-animals, ii. 205
 Hairy covering, ii. 206
Haliphysema, ii. 66
 Haller, Albrecht, i. 38
 Hand, ii. 169
 — skeleton of, ii. 302
 Hare-lip, ii. 246
 Harvey, i. 31
 Head-cap, i. 386
 — marrow, ii. 210
 — plate, i. 335
 — ribs, ii. 298
 — sheath, i. 387
 Heart, auricle of, ii. 381
 — auricular processes of, ii. 381

Heart cavity, i. 394
 — development of, ii. 385
 — human, ii. 379, 382
 — mesentery, i. 394
 — ventricle of, ii. 381
*Hecopithec*i, ii. 172
 Heredity, i. 161
 — vitiated, i. 408
 Hermaphrodites, ii. 395
 Hermaphrodite gland, ii. 401
 — Vertebrates, ii. 4
Hermaphroditismus, ii. 69, 395
*Hesperopithec*i, ii. 172
 Heterochronism, i. 13
 Heterotopism, i. 13
 Hind-brain, ii. 221, 232
 — intestine, ii. 343
 — limbs, ii. 111
 Hip-bone, ii. 278
 His, Wilhelm, i. 64
 Histogeny, i. 24
 Histology, i. 24
 Histophyly, i. 24
 Hollow-worms, ii. 76
 Holoblastic eggs, i. 215
Hologastrula, i. 241
 Homology of primitive intestine
 247; ii. 321
 — of the animal tribe
 387
 — germ-layers, i.
 — sexes, ii. 431
 Hood-gastrula, i. 200, 241
 Hoofed animals, ii. 160, 188
 Horn-plate, i. 307
 Horn-stratum of *Epidermis*, ii. 1
Humerus, ii. 278, 304
 Huxley, i. 101; ii. 294
 — germ-layer theory, i. 6
 — his Evidences, i. 101
 — Man and Ape, i. 101
 — primates, law of, ii. 17
 — skull theory, ii. 294
Hylobates, ii. 181, 189
Hypospadia, ii. 423

I

IMMACULATE CONCEPTION, i. 170
Indecidua ii. 159, 187
 Individuality, i. 123

lity of cells, i. 123
 — of metamera, i. 348
 nanic pedigree, ii. 23
 , i. 104; ii. 35
 history of earth, ii. 5
 mental capacity of, ii. 448
 ntum, ii. 199
 germ-disc, i. 291
 — vesicle, i. 291
 head-cavity, i. 335
 after, ii. 321
 blind, ii. 330
 crooked, ii. 319, 330
 digestive, ii. 330
 empty, ii. 330
 middle, ii. 330
 stomach, ii. 330
 ates, i. 414
 2, 258

J

JUSTAY, i. 101
 es, ii. 102
 r, ii. 102
 er, ii. 102
 period, ii. 14, 19

K

KAMMEL, i. 79
 ii. 403, 412
 ystem, ii. 403
 rimitive, ii. 306, 410
 sis, i. 12
 tic cleavage, i. 231
 cell (nucleus), i. 127
 rg, Nicolaus, ii. 358
 Albert, i. 59, 62
 ky, August, i. 59, 441

L

LAMINA, ii. 260
 nula, ii. 58
 Jean, i. 82
 his life, i. 82
 Man and Ape, i. 85
 "Philosophie Zoologique,"
 i. 83
 IL

Lamina dermalis i. 273, 327
 — *gastralis*, i. 273, 327
 — *inodermalis*, i. 327
 — *inoastralis*, i. 327
 — *myogastralis*, i. 327
 — *neurodermalis*, i. 327
 Lampreys, ii. 101, 121
 Lancelet, i. 253, 413
 Lankester, Ray, i. 60
Lanugo, ii. 206
 Larynx, ii. 330
Latebra (of bird's egg), i. 138
 Laurentian period, ii. 9, 19
 Layers (*Laminæ*), i. 273, 327
 Leather-plate, i. 327
 — skin, ii. 200, 232
 Leeuwenhoek, i. 37
 Leg, lower, ii. 278
 — upper, ii. 278
 Leibnitz, i. 39
 Lemuria, ii. 183
 Lemurs, ii. 164
 Lens, ii. 251, 254
Lepidosiren paradoxa, ii. 119
Leptocardia, ii. 120
 Limbs, ii. 111, 306
 — fore, ii. 302
 — hind, ii. 306
 — skeleton of, ii. 305
 — theory of, ii. 305
 Linnæus, Karl, i. 73
 Lip-cartilage, ii. 245
 — fissure, ii. 246
 Liver, ii. 330, 341
 Lizards, ii. 120, 129
Locomotorium, ii. 194, 274
 Lori, ii. 163
 Lyell, Charles, i. 77
 Lymph-cells, ii. 366
 — vessels, ii. 373

M

Macula germinativa, i. 133
Magosphara planula, ii. 60
 Male breast, ii. 204
 — cells, i. 171; ii. 392
 — copulatory organs, ii. 423
 — excretory ducts, ii. 414, 431
 — germ-glands, ii. 398
 — germ-layer, ii. 398

2 K

- Male milk-glands**, ii. 204
 — phallus (*Penis*), ii. 423
 — sexual organs, ii. 431
 — sexual plate, ii. 401
 — uterus, ii. 419
Malpighi, i. 31
Malthus, i. 98
Mamilla, ii. 204
Mamma, li. 204
Mammalia, ii. 141, 187
Mammals, egg-cleavage of, i. 210
 — gastrula of, i. 213
 — mental capacities of, ii. 448
Man-apes, ii. 178
Mantle animals, ii. 83
Marsupobranchii, ii. 104
Marsupialia, ii. 149, 187
Martins, Charles, ii. 304
Materialism, ii. 456
Maternal placenta, ii. 160
Matter, ii. 457
Mechanism in nature, i. 80
Meckel's cartilage, ii. 298
Medulla, ii. 211, 232
 — capitis, ii. 211
 — centralis, ii. 211
 — oblongata, ii. 211
 — spinalis, ii. 211
Medullary furrow, i. 302
 — membranes, ii. 228
 — plate, i. 327
 — swellings, i. 303
 — tube, i. 305
Meninges, ii. 228, 232
Meroblastic eggs, 216
Merogastrula, i. 241
Mesentery, ii. 320
Mesoderma, i. 236, 278
Mesolithic epoch, ii. 14, 19
Mesozoic periods, ii. 12, 14
Metacarpus, ii. 278, 304
Metagaster, ii. 321
Metogastrula, i. 199
Metamers, i. 346
Metameric structure, i. 347
Metanephra, ii. 412
Metatarsus, ii. 278, 304
Metazoa (intestinal animals), i. 243 ;
 ii. 92
Microlestes, ii. 149
Mid-brain, ii. 221, 232
Middle germ-layer, i. 278
 — intestine, ii. 330
 — layers, i. 278
 — part of foot, ii. 278, 304
 — hand, ii. 278, 304
Migration, theory of, i. 114
Milk, ii. 202
 — glands, ii. 143, 202
Mind, ii. 226, 447
 — activity of, ii. 210
 — cells of, i. 129
 — development of, ii. 450
 — heredity of, ii. 452
Mollusca, ii. 92, 94
Monads, i. 39
Monera, i. 180 ; ii. 43
Monerula, i. 179
Monistic philosophy, i. 16 ; ii. 456
Monocondyles, ii. 138
Monodelphia, ii. 151, 187
Monogeny, i. 160
Monophyletic origin, ii. 277
Monorhina, ii. 101, 120
Monotrema, ii. 145, 187
Monstrous evolution, i. 168
Morphogeny, i. 21, 24
Morphology, i. 21
Morphophyly, i. 24
Morula, i. 189
Motor apparatus, ii. 194, 274
 — germinative layer, i. 330
Mouth, i. 338 ; ii. 315, 330
 — cavity, ii. 315, 330
 — groove, i. 338
Mud-fishes, ii. 115, 120
Mulberry-germ, i. 189
Müller, Fritz, i. 59, 408
 — Hermann, i. 170
 — Johannes, i. 59 ; ii. 96, 414
Müllerian duct, ii. 414, 431
Muscles, i. 259 ; ii. 364
Muscle-plate, i. 353
 — system, ii. 308
Myxinoidea, ii. 101, 120

N

- Nails**, ii. 204, 232
Natural history of creation, i. 104
 — philosophy, i. 82
Navel, i. 315, 335

series, i. 400
 rd, i. 384; ii. 168
 esentery arteries, i. 395
 ——— veins, i. 399
 sins, i. 399
 ssicle, i. 337, 377
 rature, i. 369
 rrow, ii. 211, 222
 tebræ, ii. 281
 ls, i. 126
 tem, ii. 211, 232
 scular cells, ii. 232, 236
 g membrane, ii. 259
 f milk-glands, ii. 202
 ss animals, ii. 146, 204
 74; ii. 247
 pe, ii. 175
 ties, ii. 245
 , ii. 243
 ow, ii. 244
 ves, ii. 244
 esses, ii. 242
 s, ii. 242
 i. 297
 , i. 133
 i. 125
 , i. 158
 yolk, i. 216

O

, i. 114
 renz, i. 49
 grooves, ii. 240
 nerve, ii. 239
 organ, ii. 239
 is (evolution of the germ).
 tic fission of the layers,
 i. 238
 — unity, i. 366
 , i. 5
 ii. 398, 429
 ve, ii. 250
 tang, ii. 178, 181
 ii. 399, 429
 istory of the earth, ii. 7
 s without organs, ii. 45
 ny, i. 24
 gy, ii. 192
 yly, i. 24

Organ-systems, age of, ii. 357, 367
 ——— human, ii. 194
 Original cleavage, i. 198, 241
Ornithodelphia, ii. 145, 187
Ornithorhynchus, ii. 147
Ornithostoma, ii. 147, 187
Os ilium, ii. 278, 304
Os ischiæ, ii. 278, 304
Os pubis, ii. 278, 304
 Outer skin, ii. 200, 232
 Ovary, ii. 398, 430
 ——— plate, ii. 401
 Oviduct, ii. 403, 429
Ovococcus, i. 183
Ovoplasma, i. 183
Ovula holoblasta, i. 215, 241
 ——— meroblasta, i. 216, 241
 Ovulists, i. 37
Ovulum, i. 171, 183

P

Pachycardia, i. 120
 Palate, ii. 330
 ——— roof, ii. 330
 ——— soft, ii. 330
 Palæolithic epoch, ii. 10, 19
 Palæontology, i. 106
 Palæozoic periods, ii. 12, 13
Palingenesis, i. 10
 Palingenetic cleavage, i. 211
 Palingeny, i. 11
Pancreas, ii. 330, 343
 Pander, Christian, i. 51
 Paradise, ii. 183
 Parallelism in evolution, ii. 70
 Parent-cell, i. 176
 ——— kernel, i. 176
Parovarium, ii. 417, 431
Parthenogenesis, i. 40, 170
 Partial cleavage, i. 216, 240
 ——— of bird's egg, i. 224
 Pastrana, Julia, i. 374
 Pedigree, i. 112
 ——— of animals, ii. 93
 ——— Apes, ii. 189
 ——— cells, i. 467
 ——— Indo-Germanic lan-
 guages, ii. 23
 ——— Mammals, ii. 188
 ——— man, ii. 23

- Pedigree of Vertebrates, ii. 121
 Pelvic girdle, ii. 278, 304
 — intestinal cavity, i. 335
Penis, ii. 423, 431
Pentadactylia, ii. 123, 302
Perigastrula, i. 230, 241
 Peripheric nerve-system, ii. 228
 Permian period, ii. 10
Petromysontes, ii. 101, 120
Phallus, ii. 422, 431
Phallusia, i. 454
Pharynx, ii. 316, 330
 Philology, ii. 20
 — comparative, ii. 21
 Philosophy, i. 17; ii. 456
Phylogenesis (evolution of the tribe),
 i. 12
 Phylogenetic fission of the layers,
 i. 238
 — hypotheses, i. 413
 Phylogeny, i. 5, 72
 Physiogeny, i. 21, 24
 Physiology, i. 20
 — comparative, i. 20
 Physiophyly, i. 24
 Pig, i. 362
 Pigment-membrane, ii. 252
Pithecanthropi, ii. 181
 Pithecooid theory, ii. 441
Placenta, i. 383; ii. 155, 168
 — disc-shaped, ii. 162, 187
 — embryonic, ii. 160
 — *fatalis*, ii. 160
 — girdle-shaped, ii. 162, 187
 — maternal, ii. 160
 — *uterina*, ii. 160
 Placental animals, ii. 153, 187
 Placentalia, ii. 153, 187
Planæa, ii. 61
 Planæads, ii. 61
 Plant-animals,
Planula, ii. 59
Plasson, i. 130; ii. 43
 Plastids, i. 130; ii. 45
 Plastid ancestors, ii. 184
 Plastid theory, i. 180
 Plastidules, ii. 47
Plates (lamellæ), i. 303
Plathelminthes, ii. 76
Platyrrhinæ, ii. 175, 189
 Pleuro-peritoneal cavity, i. 260
 Pliocene period, ii. 11, 15
Polydactylia, ii. 123
Porus genitalis, ii. 402
 Post-glacial period, ii. 11
 Pouch-bones, ii. 151
 Pouched animals, ii. 146, 157
 Prædelineation theory, i. 37
 Præformation, i. 34
 — theory, i. 34
Præputium, ii. 423, 431
 Pressure, sense of, ii. 238
 Primary axial skeleton, ii. 235
 — age, ii. 10, 11
 — germ-layers, i. 196
 Primates, ii. 169
 Primitive amnion animals, ii. 133
 — animal ancestors, ii. 184
 — animals, or Protosoa, i. 248
 — clavicle, ii. 278
 — egg (*Protovum*), i. 134
 — fins, ii. 303
 — Fishes, ii. 112, 120
 — furrow, i. 226
 — germ-layers, i. 195
 — groove, i. 335
 — intestine, i. 444; ii. 313
 — kidney, i. 306; ii. 410
 — ducts, ii. 406
 — Mammal, ii. 142, 187
 — Man, ii. 182
 — mouth, i. 444; ii. 313
 — skull, ii. 296
 — slime, ii. 43
 — streak, i. 299
 — urine-bladder, ii. 411
 — urine-sac, i. 379; ii. 411
 — vertebrae, i. 346
 — vertebral cords, i. 305
 — vertebral plates, i. 346
 — Vertebrate (ideal), i. 256
 — (real), ii. 98
 — Worm, ii. 74, 80
 Primordial cleavage, i. 198, 241
 — kidneys, i. 307; ii. 410
 — skull, ii. 297
 — times, ii. 9, 11
 Prochorion, ii. 157
Procoracoideum, ii. 278, 304
Promammalia, ii. 142, 187
Prosimia, ii. 163, 187
Protamæba, ii. 46
Protamnion, ii. 133
Prothelmis, ii. 76

er, i. 444; ii. 314
 ra, ii. 46
 bra, ii. 410
 ma, i. 131
us annectens, ii. 119
 , ii. 248
 er, ii. 406
 dia of Amœbæ, i. 142
 i. 225, 446
 gy, ii. 448
 ne, ii. 278
germinativum, i. 135
 252, 258
 membrane, ii. 254

R

i. 278, 304
 Heinrich, i. 59
 duct, ii. 415, 431
 i. 453
 Bogulaus, i. 61
 Robert, i. 62
 412
 ction, i. 159
 ctive organs, ii. 392, 413
 ii. 120, 138
 dry intestine, i. 262; ii. 330
 — organs, i. 262; ii. 333
 i. 252, 258
 285
 iedrich, i. 101
 ouths, ii. 101, 120
 tary organs, i. 109
 ne, ii. 282
 rtebræ, ii. 282
 anus of, i. 206
 nutritive cavity, i. 207

S

DER, ii. 127
 glands,
 da, ii. 138
 ii. 278, 303
 n, M. J., i. 60, 123
 , Theodor, i. 60
 , i. 251, 258

Scolecida, ii. 86
Scrotum, ii. 423, 431
 Sea-nettles, ii. 73, 92
 Secondary age, ii. 11, 14
 — axial skeleton, ii. 291
 — germ-layers, i. 235, 273
 — kidneys, ii. 412
 — sexual character, ii. 396
 — strata of earth, ii. 12
 Seed (male), i. 36
 — animalcules, i. 173
 — cells, i. 173
 — duct, ii. 403, 429
 Segmental canals, ii. 406
 Segmentation, i. 186
 Segmentella, i. 186
Selachii, ii. 112, 121
 Selection, theory of, i. 95
 Semi-apes, ii. 164, 187
 Semicircular canals of ear, ii. 263,
 268
 Semper, Karl, i. 91, 426
 Sense of pressure, ii. 238
 — of warmth, ii. 238
 Sense-organs, ii. 238
Sensorium, ii. 194
 Sensory apparatus, ii. 194
 — functions, ii. 238
 — layer, i. 236, 229
 — nerves, ii. 238
 Sexes, separation of, ii. 69, 396
 Sexual cells, origin of,
 — cord, ii. 418
 — ducts, ii. 402, 429
 — folds, ii. 422, 431
 — furrow, ii. 422, 431
 — glands, ii. 398
 — nerves, ii. 238
 — organs, i. 266
 — plates, ii. 399
 — selection, i. 103; ii. 394
 — sense, ii. 238
 Sheath (*Vagina*), ii. 417, 431
 — of amnion, i. 387
 Shin-bone, ii. 278, 304
 Shoulder-blade, ii. 278, 304
 — girdle, ii. 278, 304
 Side-layers, i. 303
 — plates, i. 303
 — sheath, i. 303
 Silurian period, ii. 9, 19
Simia, ii. 165

Single-nostrils, ii. 101, 120
Sinus urogenitalis, ii. 419, 421
Siredon, ii. 129
 Skeleton, ii. 278
 Skeleton-forming cell-layer, ii. 287
 Skeleton, muscles of, ii. 194
 — plate, ii. 287
 Skin, ii. 195, 229
 — covering, ii. 195, 232
 — fibrous layer, i. 236
 — glands, ii. 201, 232
 — layer, i. 236
 — muscle layer, i. 236
 — muscles, ii. 194
 — navel, i. 317
 — nerves, ii. 238
 — sensory layer, i. 236
 — stratum, i. 236
 Skull, ii. 292
 — floor, ii. 292
 — roof, ii. 292
 — vertebræ, ii. 294
 — theory of, ii. 295
 Skulled Animals, ii. 100
 Skull-less Animals, i. 416; ii. 99
 Small brain, ii. 213, 232
 Soft-bodied Animals, ii. 92, 94
 Soft worms (*Scolecida*), ii. 86
Soxobanchia, ii. 129
Soxura, ii. 129
 Species (idea of), i. 73, 115
Sperma, i. 171
Spermaductus, ii. 403, 429
 Spermalists, i. 37
 Spermatzoa, i. 172
 Sperm-cells, i. 172
Spermococcus, i. 183
Spermoplasma, i. 183
Spermulum, i. 183
 Spine, ii. 280
 Spiritualism, ii. 456
 Spoke-bone (*Radius*), ii. 278, 304
 Sponges, ii. 73, 92
 Spontaneous generation, ii. 30
 Star-animals, i. 435
Stenops, ii. 164
Sternum, ii. 278
 Stomach, ii. 330
 — intestine, ii. 330
 Struggle for existence, i. 95
Subcutis, ii. 232
 Sucking worms, ii. 76

Superficial cleavage, i. 229
 Sweat-glands, ii. 202
 Swimming-bladder, ii. 111, 336
 Sylvian aqueduct, ii. 221
Synmacrobium, ii. 56
 System of animals, ii. 92
 — germ-layers, i. 273, 3
 — mammals, ii. 188
 — organs, ii. 194
 — tissues, ii. 366
 — vertebrates, ii. 120

T

TADPOLES, ii. 128
 Tail, human, i. 372; ii. 233
 Tail-cap, i. 387
 — curvature, i. 369
 — sheath, i. 387
 — vertebræ, ii. 283
 Tailed apes, ii. 180
 — *Batrachia*, ii. 129
Tarsus, ii. 278
 Taste, nerve of, ii. 238
 — sense of, ii. 238
 Teeth, ii. 173, 331
Tegumentum, ii. 199
 Teleology, i. 16, 109
Teleostei, ii. 115, 120
 Terminal budding, i. 349
 Tertiary age, ii. 11, 15
 Testes, ii. 399, 429
 — change of place of, ii. 41
 — sac, ii. 423, 431
Testiculi, ii. 399, 429
Theoria Generationis, i. 41
Thorax, ii. 282
 Thyroid gland, ii. 336
Tibia, ii. 278, 304
 Tissues, ii. 362, 366
 — age of, ii. 361, 366
 — connective, ii. 363
 — covering, ii. 361, 366
 — vascular, ii. 361, 366
 Total cleavage, i. 217, 242
 Tongue, ii. 331
 — arch, ii. 296
 — bone, ii. 298
 Tortoise, ii. 120
 Touch bodies, ii. 220

gan of, ii. 199, 238
 a forms, i. 117
 138
 period, ii. 14, 19
 story, i. 7, 24
 germ-layer, i. 239
topia, ii. 431
 rts, ii. 120
 , ii. 83, 92
 ia, ii. 79
 ain, ii. 220, 232
 aw, ii. 246
 o cavity, ii. 261, 268
 -membrane, ii. 261, 268
 em, ii. 261, 268
 animal kingdom, i. 56, 246
 eory of, i. 56, 246

U

278, 304
 , ii. 160, 188
 conception of the world, i.
 456
 , ii. 413
 i. 413
 ii. 423, 431
 bladder, ii. 413
 ducts, ii. 406
 organs, ii. 403
 sac, i. 379
 sexual cavity, ii. 419
 sexual duct, ii. 403
 system, ii. 403
 ii. 417, 431
icornis, ii. 418
nasculus, ii. 419
 i. 430

V

ii. 417, 431
 ella, ii. 48
 neden, Eduard, i. 60, 209;
 3
ferentia, ii. 403, 429
bilicalia i. 399
 r system, ii. 384

Vegetative germ-layer, i. 196, 327
 ——— organs, ii. 193, 194
 Veins, i. 393
 Ventral cavity, i. 316
 ——— plates, i. 316
 ——— vessel, i. 423
 ——— wall, i. 316
 Ventricle of heart, ii. 374
 Vermal appendage of cœcum, ii. 344
 Vertebrae, ii. 280
 ——— number of, ii. 283
 Vertebral arches, ii. 284
 ——— bodies, ii. 284
 ——— canal, ii. 284
 ——— column, ii. 280
Vertebrarium, ii. 278
 Vertebrates, ii. 92
 Vertebrates, ancestors of, ii. 185
 ——— mental capacities of,
 ii. 446
 ——— pedigree of, ii. 93
 ——— system of, ii. 97
Vesicula blastodermica, i. 290
 ——— *germinativa*, i. 133
 ——— *prostatica*, ii. 419
 ——— *umbilicalis*, i. 377
Vestibulum vaginæ, ii. 431
 Virginal generation, i. 170
Vitellus, i. 135

W

WAGNER, MORITZ, i. 114
 Wallace, Alfred, i. 98, 99
 Water, amount of in body, ii. 7
 Whale-like Animals, ii. 160, 189
 Whales, ii. 160
 Wolff, Caspar Friedrich, i. 40
 ——— his life, i. 41
 ——— his Natural Philosophy, i. 47
 ——— on formation of intestine, i. 44
 ——— on germ-layers, i. 45
 ——— *Theoria Generationis*, i. 41
 Wolffian bodies, ii. 411
 ——— duct, ii. 414, 431
 Wolff's primitive kidneys, ii. 411, 431
 Woolly hair of embryo, ii. 206
 Worms, i. 246; ii. 74
 ——— ancestors of, ii. 73
 ——— tribe of, ii. 73
 Wrist, ii. 278

Y

YELK, i. 135

— arteries, i. 395

— cavity, i. 138

— duct, i. 338; ii. 168

— formative, i. 216

— membrane, i. 138

— nutritive, i. 216

— sac, i. 337

Yolk veins, i. 395

— vessels, i. 395

Z

Zona pellucida, i. 135*Zonaplacentalia*, ii. 162, 187*Zoophyta*, i. 246; ii. 73

THE END.









STANFORD
LIBRARIES

STANFORD
LIBRARIES

STANFORD
LIBRARIES

STANFORD
LIBRARIES



STANFORD
LIBRARIES

STANFORD
LIBRARIES

STANFORD
LIBRARIES

STANFORD
LIBRARIES



STANFORD
LIBRARIES

STANFORD
LIBRARIES

STANFORD
LIBRARIES

STANFORD
LIBRARIES



Stanford University Libraries



3 6105 019 664 882

575
H133e
v.2

STANFORD UNIVERSITY LIBRARIES
STANFORD AUXILIARY LIBRARY
STANFORD, CALIFORNIA 94305-6004
(415) 723-9201

All books may be recalled after 7 days

DATE DUE

DOC FEB 08 1996

SEP 1 1999

JUN 14 2000

